

Taylor series (with radii of convergence given):

$$\begin{aligned}\frac{1}{1-x} &= 1 + x + x^2 + \cdots = \sum_0^{\infty} x^n, \quad |x| < 1 \\ e^x &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots = \sum_0^{\infty} \frac{x^n}{n!}, \quad |x| < \infty \\ \cos x &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots = \sum_0^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}, \quad |x| < \infty \\ \sin x &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots = \sum_0^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}, \quad |x| < \infty\end{aligned}$$

The Gamma function. For $x > 0$, $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$.

If x is not 0 or a negative integer, $\Gamma(x+1) = x\Gamma(x)$.

If n is a non-negative integer, $\Gamma(n+1) = n!$. $\Gamma(1/2) = \sqrt{\pi}$.

The Method of Frobenious—solution forms:

$$\begin{aligned}y_1(x) : \quad &x^{r_1} \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) : \quad &x^{r_2} \sum_{n=0}^{\infty} b_n x^n \quad \text{or} \quad y_1(x)(\ln x) + x^{r_2} \sum_{n=1}^{\infty} c_n x^n \quad \text{or} \quad C y_1(x)(\ln x) + x^{r_2} \sum_{n=0}^{\infty} d_n x^n.\end{aligned}$$

The Method of Frobenious—useful formula:

$$u'' + p(x)u' + q(x)u = \frac{C}{x^2} [y_1(x) - xp(x)y_1(x) - 2xy'_1(x)].$$

Bessel Functions.

A. The Bessel equation of order ν : $x^2 y'' + xy' + (x^2 - \nu^2)y = 0$.

B. Bessel functions:

$$\begin{aligned}J_\nu(x) &= \left(\frac{x}{2}\right)^\nu \sum_{k=0}^{\infty} \frac{(-1)^k}{k!\Gamma(\nu+k+1)} \left(\frac{x}{2}\right)^{2k} \\ J_{-\nu}(x) &= \left(\frac{x}{2}\right)^{-\nu} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!\Gamma(k-\nu+1)} \left(\frac{x}{2}\right)^{2k} \\ Y_\nu(x) &= \frac{(\cos \nu\pi)J_\nu(x) - J_{-\nu}(x)}{\sin \nu\pi}, \quad \text{if } \nu \neq 0, 1, 2, \dots \\ Y_n(x) &= \lim_{\nu \rightarrow n} Y_\nu(x), \quad \text{if } n = 0, 1, 2, \dots\end{aligned}$$

On the exam, the remaining part of this formula sheet will be Appendix C, the Laplace transform tables, from the text.