

LaTeX Math for Undergrads

Rule One Any mathematics at all, even a single character, goes in a mathematical setting. Thus, for “the value of x is 7” enter ‘the value of $\langle x \rangle$ is $\langle 7 \rangle$ ’.

Template Your document should contain at least this.

```
\documentclass{article}
\usepackage{amsmath, amssymb, amsthm}
\usepackage{utf8}{inputenc}

\begin{document}
--document body here--
\end{document}
```

Common constructs

```
x^2 x~2 \sqrt{2}, \sqrt[3]{ } \sqrt{2}, \sqrt[n]{3}
x_{i,j} x_{i,j} \frac{2}{3}, 2/3 \frac{2}{3}, 2/3
```

Calligraphic letters Use as $\langle \mathcal{A} \rangle$.

```
ABCDEFGHIJKLMN O PQRSTU VWX YZ
```

Greek

```
\alpha \alpha \xi, \Xi \xi, \Xi
\beta \beta o o
\gamma, \Gamma \gamma, \Gamma \pi, \Pi \pi, \Pi
\delta, \Delta \delta, \Delta \varpi \varpi
\epsilon \epsilon \rho \rho
\varepsilon \varepsilon \varrho \varrho
\zeta \zeta \sigma, \Sigma \sigma, \Sigma
\eta \eta \varsigma \varsigma
\theta \Theta \theta, \Theta \tau \tau
\vartheta \vartheta \upsilon, \Upsilon \upsilon, \Upsilon
\iota \iota \phi, \Phi \phi, \Phi
\kappa \kappa \varphi \varphi
\lambda \Lambda \lambda, \Lambda \chi \chi
\mu \mu \psi, \Psi \psi, \Psi
\nu \nu \omega, \Omega \omega, \Omega
```

Sets and logic

```
\cup \cup \mathbb{R} \mathbb{R} \forall \forall
\cap \cap \mathbb{Z} \mathbb{Z} \exists \exists
\subset \subset \mathbb{Q} \mathbb{Q} \neg \neg
\subseteq \subseteq \mathbb{N} \mathbb{N} \vee \vee
\supset \supset \mathbb{C} \mathbb{C} \wedge \wedge
\supseteq \supseteq \emptyset \emptyset \vdash \vdash
\in \in \emptyset \emptyset \models \models
\ni \ni \aleph \aleph \Rightarrow \Rightarrow
\notin \notin \setminus \setminus \nRightarrow \nRightarrow
\in \in \equiv \equiv
```

Negate an operator, as in $\not\subset$, with $\not\subset$. For the set complement, get A^c with \mathcal{A}^c , get A^c with \mathcal{A}^c with \mathcal{A}^c , or get \bar{A} with \bar{A} .

Decorations

```
f' f' \dot{a} \dot{a} \tilde{x} \tilde{x}
f'' f'' \ddot{a} \ddot{a} \bar{x} \bar{x}
\Sigma^* \Sigma^* \hat{x} \hat{x} \vec{x} \vec{x}
```

If the decorated letter is i or j then some decorations need \imath or \jmath , as in $\vec{\imath}$. Some authors use boldface for vectors: \boldsymbol{x} .

Entering $\overline{x+y}$ produces $\overline{x+y}$, and $\widehat{x+y}$ gives $\widehat{x+y}$. Comment on an expression as here (there is also $\overbrace{\dots}$).

```
x+y \underbrace{x+y}_{|A|}
```

Dots Use low dots in a list $\{0, 1, 2, \dots\}$, entered as $\{0, 1, 2, \dots\}$. (If you use \dots in plain text as London, Paris, \dots , note the thinspace $\,$, before the period.) Use centered dots in a sum or product $1 + \dots + 100$, entered as $1+\cdots+100$. You can also get vertical dots \vdots and diagonal dots \ddots .

Roman names Enter $\tan(x)$, with a backslash, instead of $\tan(x)$. These get the same treatment.

```
\sin \sin \sinh \sinh arcsin \arcsin
\cos \cos \cosh \cosh arccos \arccos
\tan \tan \tanh \tanh arctan \arctan
\sec \sec \coth \coth \min \min
\csc \csc \det \det \max \max
\cot \cot \dim \dim \inf \inf
\exp \exp \ker \ker \sup \sup
\log \log \deg \deg \liminf \liminf
\ln \ln \arg \arg \limsup \limsup
\lg \lg \gcd \gcd \lim \lim
```

Other symbols

```
< < \angle \angle \cdot \cdot
\leq \leq \sphericalangle \sphericalangle \pm \pm
> > \ell \ell \mp \mp
\geq \geq \parallel \parallel \times \times
\neq \neq 45^\circ 45^\circ \div \div
\ll \ll \cong \cong * *
\gg \gg \ncong \ncong | |
\approx \approx \sim \sim \dagger \dagger
\asymp \asymp \simeq \simeq n! n!
\equiv \equiv \nsim \nsim \partial \partial
\prec \prec \oplus \oplus \nabla \nabla
\preceq \preceq \ominus \ominus \hbar \hbar
\succ \succ \odot \odot \circ \circ
\succeq \succeq \otimes \otimes * *
\propto \propto \oslash \oslash \surd \surd
\doteq \doteq \upharpoonright \upharpoonright \checkmark \checkmark
```

Enter $a|b$ for the divides relation $a|b$. Use \mid as in $\{a \in S \mid \text{text}(a=0) \text{ or } (a \text{ is odd})\}$ for the set $\{a \in S \mid a = 0 \text{ or } a \text{ is odd}\}$.

Variable-sized operators The summation $\sum_{j=0}^3 j^2$ $\sum_{j=0}^3 j^2$ and the integral $\int_{x=0}^3 x^2 dx$ $\int_{x=0}^3 x^2 dx$ expand when displayed.

$$\sum_{j=0}^3 j^2 \quad \int_{x=0}^3 x^2 dx$$

These do the same.

```
\int \int \iint \iiint \oint \bigcup \bigcap
```

Arrows

\rightarrow	<code>\rightarrow, \to</code>	\mapsto	<code>\mapsto</code>
\rightrightarrows	<code>\rightrightarrows</code>	\longmapsto	<code>\longmapsto</code>
\longrightarrow	<code>\longrightarrow</code>	\leftarrow	<code>\leftarrow</code>
\Rightarrow	<code>\Rightarrow</code>	\Leftrightarrow	<code>\Leftrightarrow</code>
\nrightarrow	<code>\nrightarrow</code>	\downarrow	<code>\downarrow</code>
\Longrightarrow	<code>\Longrightarrow</code>	\uparrow	<code>\uparrow</code>
\rightsquigarrow	<code>\rightsquigarrow</code>	\Updownarrow	<code>\Updownarrow</code>

The right arrows in the first column have matching left arrows, such as `\leftarrow`, and there are some other matches for down arrows, etc.

Fences

$()$	$()$	$\langle \rangle$	<code>\langle \rangle</code>	$ $	$ $
$[]$	$[]$	$\lfloor \rfloor$	<code>\lfloor \rfloor</code>	$ $	$\ \ $
$\{ \}$	$\{ \}$	$\lceil \rceil$	<code>\lceil \rceil</code>		

They will grow with the enclosed formula using `\left` and `\right`.

$$\left\langle i, 2^{2^i} \right\rangle \left\langle i, 2^{2^i} \right\rangle$$

Every `\left` must match a `\right` and they must end on the same line in the output. For a one-sided fence put a period `\left.` or `\right.` on the other side.

$$\left. \frac{df}{dx} \right|_{x_0}$$

Fix the size with `\big`, `\Big`, `\bigg`, or `\Bigg`.

$$\left[\sum_{k=0}^n e^{k^2} \right]$$

Arrays, Matrices Make an array of mathematical text as you make a table of plain text.

0	\leftrightarrow	0	<code>\begin{array}{rcl}</code>
1	\leftrightarrow	1	<code>0 \&\leftarrow &0 \\\</code>
2	\leftrightarrow	4	<code>1 \&\leftarrow &1 \\\</code>
			<code>2 \&\leftarrow &4 \\\</code>
			<code>\vdots & \vdots</code>
			<code>\end{array}</code>

Definition by cases is an array with two columns.

$$f_n = \begin{cases} a & \text{if } n = 0 \\ r \cdot f_{n-1} & \text{else} \end{cases}$$

A matrix is another array variant. With this abbreviation you need not specify that columns are centered.

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

For the determinant use `|A|` inline and `\vmatrix` in display.

Spacing in mathematics

$\rightarrow \leftarrow$	$\backslash,$	$\rightarrow \leftarrow$	<code>\quad</code>
$\rightarrow \leftarrow$	$\backslash:$	$\rightarrow \leftarrow$	<code>\quad</code>
$\rightarrow \leftarrow$	$\backslash;$	$\rightarrow \leftarrow$	<code>\!</code>

The left column spaces are in ratio 3 : 4 : 5. The last in the right column is a negative space, opposite to `\,`. Get arbitrary space as in `\hspace{0.5cm}`.

Displayed equations Put equations on a separate line with the `equation*` environment.

$$S = k \log W$$

You can break into multiple lines.

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

Align using the `align*` environment

$$\begin{aligned} \nabla \cdot \mathbf{D} &= \rho \\ \nabla \cdot \mathbf{B} &= 0 \end{aligned}$$

(you can have an empty left or right side of the alignment). For each environment, get a numbered version by omitting the asterisk, as with `align` in place of `align*`.

Calculus examples The last three here are display style.

$$f: \mathbb{R} \rightarrow \mathbb{R}$$

$$9.8 \text{ m/s}^2$$

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\int x^2 dx = x^3/3 + C$$

$$\nabla = i \frac{d}{dx} + j \frac{d}{dy} + k \frac{d}{dz}$$

Discrete mathematics examples There are four modulo forms: $m \bmod n$ is from `m\bmod n`, and $a \equiv b \pmod m$ is from `a\equiv b\pmod m`, and $a \equiv b \pmod m$ is from `a\equiv b\pmod m`, and $a \equiv b \pmod m$ is from `a\equiv b\pmod m`.

For combinations the binomial symbol $\binom{n}{k}$ is from `\binom{n}{k}`. This resizes to be bigger in a display (to require the display version use `\dbinom{n}{k}` and for the inline version use `\tbinom{n}{k}`).

For permutations use n^r from `n^{\underline{r}}` (some authors use $P(n, r)$, or ${}_nP_r$ from `\{}_nP_r`).

Statistics examples

$$\sigma^2 = \sqrt{\sum (x_i - \mu)^2 / N}$$

$$E(X) = \mu_X = \sum (x_i \cdot P(x_i))$$

The probability density of the normal distribution

$$\frac{1}{\sqrt{2\sigma^2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

comes from this.

$$\frac{1}{\sqrt{2\sigma^2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

For more See also the Comprehensive L^AT_EX Symbols List at mirror.ctan.org/info/symbols/comprehensive and DeT_EXify at detexify.kirelabs.org/classify.html.