

CORRECTIONS TO

REPRESENTATIONS AND INVARIANTS OF THE CLASSICAL GROUPS

by Roe Goodman and Nolan R. Wallach

(1999 paperback printing)

Revised August 30, 2002

**p.v** (Table of Contents): §1.2 begins on page 18

**p.v** (Table of Contents): §1.3 begins on page 35

**p.25, 1.10** REPLACE: We denote by  $s_0$

BY: We denote by  $s_l$

**p.25, 1.11 (display)** REPLACE:  $s_0$  BY:  $s_l$

**p.25, 1.13** REPLACE:

$$J_+ = \begin{bmatrix} 0 & s_0 \\ s_0 & 0 \end{bmatrix}, \quad J_+ = \begin{bmatrix} 0 & s_0 \\ -s_0 & 0 \end{bmatrix},$$

BY:

$$J_+ = \begin{bmatrix} 0 & s_l \\ s_l & 0 \end{bmatrix}, \quad J_+ = \begin{bmatrix} 0 & s_l \\ -s_l & 0 \end{bmatrix},$$

**p.25, 1.-10** REPLACE:  $s_0 a^t s_0$  BY:  $s_l a^t s_l$

**p.25, 1.-7** REPLACE:

$$A = \begin{bmatrix} a & b \\ c & -s_0 a^t s_0 \end{bmatrix},$$

BY:

$$A = \begin{bmatrix} a & b \\ c & -s_l a^t s_l \end{bmatrix},$$

**p. 25, 1.-6** REPLACE: such that  $b^t = -s_0 b s_0$  and  $c^t = -s_0 c s_0$

BY: such that  $b^t = -s_l b s_l$  and  $c^t = -s_l c s_l$

**p.25, l.-3** REPLACE:

$$A = \begin{bmatrix} a & b \\ c & -s_0 a^t s_0 \end{bmatrix},$$

BY:

$$A = \begin{bmatrix} a & b \\ c & -s_l a^t s_l \end{bmatrix},$$

**p. 25, l.-2** REPLACE: such that  $b^t = s_0 b s_0$  and  $c^t = s_0 c s_0$

BY: such that  $b^t = s_l b s_l$  and  $c^t = s_l c s_l$

**p.26, l.6** REPLACE:

$$S = \begin{bmatrix} 0 & 0 & s_0 \\ 0 & 1 & 0 \\ s_0 & 0 & 0 \end{bmatrix}.$$

BY:

$$S = \begin{bmatrix} 0 & 0 & s_l \\ 0 & 1 & 0 \\ s_l & 0 & 0 \end{bmatrix}.$$

**p.26, l.12** REPLACE:

$$A = \begin{bmatrix} a & w & b \\ u & 0 & -w^t s_0 \\ c & -s_0 u^t & -s_0 a^t s_0 \end{bmatrix},$$

BY:

$$A = \begin{bmatrix} a & w & b \\ u & 0 & -w^t s_l \\ c & -s_l u^t & -s_l a^t s_l \end{bmatrix},$$

**p.26, l.13** REPLACE: such that  $b^t = -s_0 b s_0$  and  $c^t = -s_0 c s_0$

BY: such that  $b^t = -s_l b s_l$  and  $c^t = -s_l c s_l$

**p.30, l.14** REPLACE:  $f_C(g)$  BY:  $f_C^\pi(g)$

**p.30, l.16** REPLACE:  $f_{d\pi(A)C}(g)$  BY:  $f_{d\pi(A)C}^\pi(g)$

**p.31, l.-7 and -8** REPLACE:

$$X_{d\sigma(A)}f = X_A(f \circ \sigma) = X_{d\pi(A)}(f \circ \rho) = X_{d\rho(d\pi(A))}f.$$

Hence  $d\sigma(A) = d\rho(d\pi(A))$ .

BY:

$$X_{d\sigma(A)}f^\sigma = X_A(f \circ \sigma) = X_{d\pi(A)}(f \circ \rho) = X_{d\rho(d\pi(A))}(f \circ \rho).$$

Now evaluate at  $I$  to see that  $d\sigma(A) = d\rho(d\pi(A))$ .

**p.32, 1.7** REPLACE:  $\{(A, -A) : A \in \mathfrak{g} \cap \mathfrak{h}\}$

BY:  $\{(A, A) : A \in \mathfrak{g} \cap \mathfrak{h}\}$

**p.32, 1.24** REPLACE: Now assume  $d\sigma(\mathfrak{g}) = \mathfrak{h}$ .

BY: Now assume  $d\sigma(\mathfrak{g}) = \mathfrak{h}$  and  $H$  is connected.

**p.40, end of line 1** The last word should be HINT:

**p.40, end of line 3** The last word should be “that”

**p.45, 1.-10** REPLACE:  $\{g \in G : gg^t = I\}$

BY:  $\{g \in \mathrm{GL}(n, \mathbb{C}) : gg^t = I\}$

**p.48, 1.14** REPLACE: Theorem D.2.7 BY: Theorem D.2.8

**p.48, 1.-3** ADD SENTENCE: Hence  $\det g \in \mathbb{R}$ .

**p.49, 1.-10** REPLACE:

Exercis (REMAINDER OF LINE DISAPPEARED IN TYPSETTING)

BY:

Exercises A.1.7, # 6). However, in the case of algebraic groups, Theorem

**p.94, 1.6** REPLACE: Let  $s_0 \in \mathrm{GL}(2l, \mathbb{C})$

BY: Let  $s_l \in \mathrm{GL}(l, \mathbb{C})$

**p.94, 1.10** REPLACE:

$$\pi(\sigma) = \begin{bmatrix} s_\sigma & 0 \\ 0 & s_0 s_\sigma s_0 \end{bmatrix},$$

BY:

$$\pi(\sigma) = \begin{bmatrix} s_\sigma & 0 \\ 0 & s_l s_\sigma s_l \end{bmatrix},$$

**p.95, 1.8** REPLACE:

$$\phi(\sigma) = \begin{bmatrix} s_\sigma & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & s_0 s_\sigma s_0 \end{bmatrix},$$

BY:

$$\phi(\sigma) = \begin{bmatrix} s_\sigma & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & s_l s_\sigma s_l \end{bmatrix},$$

**p.155, 1.10** REPLACE: for all  $x \in F$ .

BY: for all  $x \in F$ . (HINT: Use Proposition 3.4.7.)

**p.170, 1.12** REPLACE: if  $\phi \in \mathcal{J}$  then there exist

BY: if  $\phi \in \mathcal{J}_+$  then there exist

**p.174, 1.-7** REPLACE: induction that  $\mathcal{H} \cdot (\mathcal{P}\mathcal{J}_+)$  contains all polynomials

BY: induction that  $\mathcal{H} \cdot (1 + \mathcal{P}\mathcal{J}_+)$  contains all polynomials

**p.175, 1.7** REPLACE: 4.1.4(1), which contradicts

BY: 4.1.4, which contradicts

**p.183, 1.-7 display** REPLACE:

$$uZw = \begin{bmatrix} I_r & O_{r,m-r} \\ O_{m-r,r} & O_{m-r} \end{bmatrix}$$

BY:

$$uZw = \begin{bmatrix} I_r & O_{r,m-r} \\ O_{k-r,r} & O_{k-r,m-r} \end{bmatrix}$$

**p.183, 1.-5 display** REPLACE:

$$X = \begin{bmatrix} I_r & O_{r,m-r} \\ O_{m-r,r} & O_{k-r} \end{bmatrix}, \quad Y = \begin{bmatrix} I_r & O_{r,n-r} \\ O_{n-r,r} & O_{n-r} \end{bmatrix},$$

BY:

$$X = \begin{bmatrix} I_r & O_{r,n-r} \\ O_{k-r,r} & O_{k-r,n-r} \end{bmatrix}, \quad Y = \begin{bmatrix} I_r & O_{r,m-r} \\ O_{n-r,r} & O_{n-r,m-r} \end{bmatrix},$$

**p.184, l.-8 display** REPLACE:

$$X = \begin{bmatrix} J_r & O_{r,k-r} \\ O_{k-r,r} & O_{n-r,k-r} \end{bmatrix} g.$$

BY:

$$X = \begin{bmatrix} J_r & O_{r,k-r} \\ O_{n-r,r} & O_{n-r,k-r} \end{bmatrix} g.$$

**p.189, l.10 display** REPLACE:  $\prod_{j=1}^k y_j^{q_j}$  BY:  $\prod_{j=1}^m y_j^{q_j}$

**p.189, l.13** REPLACE:  $z = (v_1, \dots, v_k, v_1^*, \dots, v_k^*)$

BY:  $z = (v_1, \dots, v_k, v_1^*, \dots, v_m^*)$

**p.198, l.-7** REPLACE: representation on  $\mathbb{C}^n$  BY: representation on  $V$

**p.198, l.-5** REPLACE: space  $\mathcal{P}^{[p,q]}(V^k \otimes (V^*)^m)^{\text{GL}(V)}$

BY: space  $\mathcal{P}^{[p,q]}(V^k \oplus (V^*)^m)^{\text{GL}(V)}$

**p.198, l.-3** REPLACE: acts on  $\mathcal{P}^{[p,q]}(V^k \otimes (V^*)^m)$

BY: acts on  $\mathcal{P}^{[p,q]}(V^k \oplus (V^*)^m)$

**p.198, l.-1 display** REPLACE:  $\mathcal{P}^{[p,q]}(V^k \otimes (V^*)^m)^{\text{GL}(V)} = 0$

BY:  $\mathcal{P}^{[p,q]}(V^k \oplus (V^*)^m)^{\text{GL}(V)} = 0$

**p.199, l.2 display** REPLACE:  $\mathcal{P}^{[p,q]}(V^k \otimes (V^*)^m)^{\text{GL}(V)}$

BY:  $\mathcal{P}^{[p,q]}(V^k \oplus (V^*)^m)^{\text{GL}(V)}$

**p.199, l.4** REPLACE: complete contractions  $C_s$  BY: complete contractions  $\lambda_s$

**p.199, l.6 display** REPLACE:  $C_s$  BY:  $\lambda_s$

**p.199, l.9 display** REPLACE:  $C_s$  BY:  $\lambda_s$

**p.227, l.-1** REPLACE: general Capelli problem.”

BY: general “Capelli problem.”

**p.254, l.-17 to l.-14** REPLACE:

Suppose for the sake . . . impossible. Hence  $V_k = \mathcal{H}^k(\mathbb{C}^n)$ .

BY:

We next prove (3). The intertwining property is easily checked (here it is understood that in  $V^k \otimes \mathcal{H}^k(\mathbb{C}^n)$ ,  $\mathfrak{g}'$  acts only on  $V^k$  and  $G$  acts only on  $\mathcal{H}^k(\mathbb{C}^n)$ ). Suppose  $\sum \psi_j(r^2)f_j = 0$ . We may assume that  $\{f_j\}$  is linearly independent and by homogeneity that  $\psi_j(r^2) = c_j r^2$ . This immediately implies  $c_j = 0$ , proving injectivity of the map. From (3) and Theorem 4.5.12 it follows that  $\mathcal{H}^k(\mathbb{C}^n)$  is an irreducible  $G$ -module. Hence  $V_k = \mathcal{H}^k(\mathbb{C}^n)$ .

**p.254, l.-7** REPLACE:

Statement (3) follows from (1) and (2) and Theorem 4.5.12. To prove (4) we only

BY:

To prove (4), we only

**p.257, l.-3** REPLACE: *of size  $r$  such that* BY: *of size  $2r$  such that*

**p.258, l.18** REPLACE: *it has degree  $|\mu|$*  BY: *it has degree  $|\mu|/2$*

**p.259, l.5** REPLACE: *such that  $|\mu| = r$  and* BY: *such that  $|\mu| = 2r$  and*

**p.272, l.-1** REPLACE:  $\epsilon(x^*)\epsilon(y^*) = -\epsilon(x^*)\epsilon(y^*)$  BY:  $\epsilon(x^*)\epsilon(y^*) = -\epsilon(y^*)\epsilon(x^*)$

**p.273, change displayed formula numbers:**

(4.5.5)  $\longrightarrow$  (6.5.5)

(4.5.6)  $\longrightarrow$  (6.5.6)

(4.5.7)  $\longrightarrow$  (6.5.7)

(4.5.8)  $\longrightarrow$  (6.5.8)

**p.275, change displayed formula number:**

(4.5.9)  $\longrightarrow$  (6.5.9)

**p.276, change displayed formula numbers:**

$$(4.5.10) \longrightarrow (6.5.10)$$

$$(4.5.11) \longrightarrow (6.5.11)$$

**p.304, l.-9** REPLACE:  $\pi$  BY:  $\pi^\lambda$

**p.340, l.-16** REPLACE:  $\gamma s_0 \gamma^t = I_{2l}$  BY:  $\gamma s_{2l} \gamma^t = I_{2l}$

**p.340, l-15** REPLACE: where  $s_0$  is the matrix

BY: where  $s_{2l}$  is the matrix

**p.340, l-14** REPLACE: corresponding to  $s_0$  as in

BY: corresponding to  $s_{2l}$  as in

**p.340, l-12** REPLACE:

$$\gamma g \gamma^{-1} (\gamma g \gamma^{-1})^t = \gamma g s_0 g^t \gamma^t = \gamma s_0 \gamma^t = I_{2l}.$$

BY:

$$\gamma g \gamma^{-1} (\gamma g \gamma^{-1})^t = \gamma g s_{2l} g^t \gamma^t = \gamma s_{2l} \gamma^t = I_{2l}.$$

**p.340, l-9** REPLACE: defined by the equation  $g^t g = I$ .

BY: defined by the equation  $g^t g = I_{2l}$ .

**p.342, l.-12** REPLACE: invariant volume forms on  $K/T$ ,  $T$ , and  $K$ , respectively

BY: invariant volume forms on  $T$ ,  $K/T$ , and  $K$ , respectively

**p.371, l.6** REPLACE: (published by CAN, Amsterdam, 1992).

BY: (available at <http://young.sp2mi.univ-poitiers.fr/~marc/LiE>)

**p.376, l.9** REPLACE: representation  $(\sigma_k^\lambda, G^\lambda)$  BY: representation  $(\sigma^\lambda, G^\lambda)$

**p.377, l.-3** REPLACE:  $\lambda^*$  BY:  $\lambda^T$

**p.377, l.-2** REPLACE:  $\lambda_1^* \geq \lambda_2^*$  BY:  $\lambda_1^T \geq \lambda_2^T$

**p.377, l.-1** REPLACE:  $\lambda_j^*$  BY:  $\lambda_j^T$

**p.378, l.1, l.2, and l.4** REPLACE:  $\lambda^*$  BY:  $\lambda^T$  (4 REPLACEMENTS)

**p.384, l.10** REPLACE:

(1) The multiplicities  $m_{\pi, \pi'}$  in (9.2.1) are either 0 or 1,

BY:

(1) The multiplicities  $m_{\pi, \pi'}$  are either 0 or 1 and each  $\pi \in \widehat{K}$  (resp.  $\pi' \in \widehat{K}'$ ) occurs at most once in (9.2.1).

**p.384, l.-3** REPLACE:

$G \times G'$  on  $Y$  is also multiplicity free.

BY:

$(\rho(G), \rho(G'))$  on  $Y$  is a dual reductive pair.

**p.385, l.8** REPLACE: for  $f \in \mathcal{P}(X)$  BY: for  $f \in \mathcal{P}^d(X)$

**p.385, l.-1** REPLACE:

This shows that  $\mathcal{P}^d(X)$  is multiplicity-free for  $G \times G'$

BY:

This shows that  $(G, G')$  acts as a dual pair on  $\mathcal{P}^d(X)$ .

**p.386, l.5** REPLACE: Thus  $Y$  is multiplicity free for  $K \times K'$ .

BY: Thus  $(K, K')$  acts as a dual pair on  $Y$ .

**p.386, l.10 and l.11** REPLACE:

As a  $(G \times G')$ -module,  $Y$  is also multiplicity free, since

BY:

We have  $(G, G')$  acting as a dual pair on  $Y$ , since

**p.386, l.-1** REPLACE:  $V_{(\tau)}$  BY:  $V_\tau$



**p.394, 1.7** REPLACE:  $\lambda^*$  BY:  $\lambda^T$

**p.394, 1.9** REPLACE:

$$e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_1^*} \otimes e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_2^*} \otimes \cdots \otimes e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_q^*}$$

BY:

$$e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_1^T} \otimes e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_2^T} \otimes \cdots \otimes e_1 \otimes e_2 \otimes \cdots \otimes e_{\lambda_q^T}$$

**p.394, 1.10** REPLACE:  $\lambda_1^*$  BY:  $\lambda_1^T$

**p.394, 1.11** REPLACE:  $\lambda_1^* + 1, \dots, \lambda_2^*$

$$\text{BY: } \lambda_1^T + 1, \dots, \lambda_2^T$$

**p.394, 1.12** REPLACE:  $\omega_{\lambda_1^*} \otimes \omega_{\lambda_2^*} \otimes \cdots \otimes \omega_{\lambda_q^*}$ ,

$$\text{BY: } \omega_{\lambda_1^T} \otimes \omega_{\lambda_2^T} \otimes \cdots \otimes \omega_{\lambda_q^T},$$

**p.403, 1.7** REPLACE:  $\lambda^*$  BY:  $\lambda^T$  (2 REPLACEMENTS)

**p.403, 1.8** REPLACE:  $\text{Col}(A^*)$  BY:  $\text{Col}(A^T)$

**p.403, 1.12 and 1.13** REPLACE:

suppose  $\lambda = \lambda^*$ . Prove that  $\sigma^\lambda(s) = 0$  for all odd permutations  $s \in \mathfrak{S}_k$ .

BY:

suppose  $\lambda = \lambda^T$ . Prove that  $\chi^\lambda(s) = 0$  for all odd permutations  $s \in \mathfrak{S}_k$ , where  $\chi^\lambda$  is the character of  $\sigma^\lambda$ .

**p.418, 1.-3** REPLACE:  $\mu_1 \geq \cdots \geq \mu_n$

$$\text{BY: } \mu_1 \geq \cdots \geq \mu_n \geq 0$$

**p.434, 1.11** REPLACE:

$$\mathcal{HT}_r^{\otimes k} = \{u \in \mathcal{T}_r^{\otimes k} : u \cdot u = 0 \text{ for all } u \in \mathcal{B}_{k,r+1}(V, \omega)\}$$

BY:

$$\mathcal{HT}_r^{\otimes k} = \{u \in \mathcal{T}_r^{\otimes k} : z \cdot u = 0 \text{ for all } z \in \mathcal{B}_{k,r+1}(V, \omega)\}$$

**p.436, equation (10.3.4)** REPLACE:

$$1 \leq m(r, \lambda) \leq \dim(G^\lambda) |\mathcal{M}(k, r)|$$

BY:

$$\dim(G^\lambda) \leq m(r, \lambda) \leq \dim(G^\lambda) |\mathcal{M}(k, r)|$$

**p.436, l.-8** REPLACE: Let  $r \geq 0$  BY: Let  $r > 0$

**p.487, l.-5 and l.-6** REPLACE:

and

$$\frac{d}{dt}(y^{-1}\theta(y)(I + t\theta(B))y(I + tB))|_{t=0} = \text{Ad}(y^{-1})\theta(B) + B.$$

BY:

whereas the curve  $t \mapsto y(I + tB)$  is tangent to  $Q$  at  $y$  provided

$$0 = \frac{d}{dt}(y^{-1}\theta(y)(I + t\theta(B))y(I + tB))|_{t=0} = \text{Ad}(y^{-1})\theta(B) + B.$$

**p.502, l.17** REPLACE:

algebraic groups by Theorem ??.

BY:

algebraic groups by Corollary 11.1.3.

**p.502, l.-5** REPLACE:

$$(\exp yX_0)g(\exp -yX_0) = t \exp[(t^{-\alpha} - 1)y + zX_0].$$

BY:

$$(\exp yX_0)g(\exp -yX_0) = t \exp[((t^{-\alpha} - 1)y + z)X_0].$$

**p.544, l.-4** REPLACE:  $G = \text{SO}(\mathbb{C}_n, B)$

BY:  $G = \text{SO}(\mathbb{C}^n, B)$

**p.561, l.18** REPLACE:  $\theta(g) = JgJ^{-1}$ .

BY:  $\theta = \theta_{n,n}$  as in Type AIII.

**p.566, l.-15** REPLACE: set  $\mathcal{J} = \{f_{\text{top}} : f \in \mathcal{I}\}$ .

BY: set  $\mathcal{J} = \text{span}\{f_{\text{top}} : f \in \mathcal{I}\}$ .

**p.566, l.-11** REPLACE: It is even easier to prove that  $\mathcal{J}$  is closed under addition.

BY: By definition  $\mathcal{J}$  is closed under addition.

**p.567, l.13** REPLACE:  $\{f_{\text{top}} : f \in \mathcal{I}_c\}$ . BY:  $\text{span}\{f_{\text{top}} : f \in \mathcal{I}_c\}$ .

**p.597, change displayed formula numbers:** (4.5.1)  $\longrightarrow$  (A.3.1)

(4.5.2)  $\longrightarrow$  (A.3.2)

**p.598, l.15** CHANGE DISPLAYED FORMULA NUMBER : (4.5.3) TO: (A.3.3)

**p.599, l. -12** CHANGE DISPLAYED FORMULA NUMBER : (4.5.4) TO: (A.3.4)

**p.599, l.-7** REPLACE: Theorem 13.3.1 BY: Theorem A.3.1

**p.599, l.-4** CHANGE DISPLAYED FORMULA NUMBER : (4.5.5) TO: (A.3.5)

**p.600, l.5** REPLACE: /6, Theorem 14 BY: §6, Theorem 14

**p.600, l.11** CHANGE DISPLAYED FORMULA NUMBER : (4.5.6) TO: (A.3.6)

**p.600, l.-8** CHANGE DISPLAYED FORMULA NUMBER : (4.5.7) TO: (A.3.7)

**p.600, l.-2** REPLACE: Proposition 13.3.2 BY: Proposition A.3.2

**p.601, l.12** CHANGE DISPLAYED FORMULA NUMBER : (4.5.8) TO: (A.3.8)

**p.601, l.-12** REPLACE: Theorem 13.3.3 BY: Theorem A.3.3

**p.601, l.-5** REPLACE: Lemma 13.3.4 BY: Lemma A.3.4

**p.602, l.7** CHANGE DISPLAYED FORMULA NUMBER : (4.5.9) TO: (A.3.9)

**p.602, l.12** CHANGE DISPLAYED FORMULA NUMBER : (4.5.10) TO: (A.3.10)

**p.603, l.-15** REPLACE: Corollary 13.3.5 BY: Corollary A.3.5

**p.604, 1.5** CHANGE DISPLAYED FORMULA NUMBER : (4.5.11) TO: (A.3.11)

**p.604, 1.9** REPLACE: Proposition 13.3.6 BY: Proposition A.3.6

**p.605, 1.-3** REPLACE: Lemma 13.4.1 BY: Lemma A.4.1

**p.606, 1.-3** REPLACE: view  $\mathbb{C}^r = M_{1 \times r}$  as row vectors. We

BY: view  $\mathbb{C}^r$  as  $r \times 1$  matrices. We

**p.607, 1.5** CHANGE DISPLAYED FORMULA NUMBER : (4.5.1) TO: (A.4.1)

**p.607, 1.9** REPLACE: on  $\mathbb{C}^m$  and  $\mathbb{C}^n$  that BY: on  $\mathbb{C}^{m+1}$  and  $\mathbb{C}^{n+1}$  that

**p.607, 1.-10** REPLACE: Lemma 13.4.2 BY: Lemma A.4.2

**p.608, 1.-12** REPLACE: Lemma 13.4.3 BY: Lemma A.4.3

**p.609, 1.17** REPLACE: Lemma 13.4.4 BY: Lemma A.4.4

**p.609, 1.21** REPLACE: Proposition 13.4.5 BY: Proposition A.4.5

**p.609, 1.-3** REPLACE: Corollary 13.4.6 BY: Corollary A.4.6

**p.610, 1.5** REPLACE: Theorem 13.4.7 BY: Theorem A.4.7

**p.610, 1.-16** REPLACE: Corollary 13.4.8 BY: Corollary A.4.8

**p.611, 1.11** REPLACE: Ch. I, / 5.2, Theorem 3 BY: Ch. I.5.2, Theorem 3

**p.611, 1.4** REPLACE: called *simple* if BY: called *smooth* if

**p.611, 1.5** REPLACE: the simple points BY: the smooth points

**p.611, 1.-7** REPLACE: Theorem 13.4.9 BY: Theorem A.4.9

**p.611, 1.-4** REPLACE: Corollary 13.4.10 BY: Corollary A.4.10

**p.620, 1.-11** REPLACE:  $\rightarrow W$  BY:  $\rightarrow W^{\otimes p}$

**p.649, 1.-10** REPLACE: (Section A.1.5) BY: (Section A.1.6)

**p.649, l.-9** REPLACE: set of regular points BY: set of smooth points

**p.650, l.10** REPLACE: (Corollary A.3.3) BY: (Theorem A.1.18)

**p.651, l.4** REPLACE: (**Submanifolds**) BY: (**Open submanifolds**)

**p.655, l.-6** REPLACE: (Example 5 of Section D.1.1)

BY: (Example 6 of Section D.1.1)

**p.658, formula D.1.4** REPLACE:  $|\varphi|\omega$  BY:  $\varphi\omega$

**p.661, l.2** REPLACE: is regular (Proposition 1.2.3)

BY: is smooth (Theorem 1.2.3)

**p.661, l.14** REPLACE: D.1.2, #4). BY: D.1.4, #4).

**p.663, l.-1** REPLACE: chart containing (1) BY: chart containing 1)

**General Typesetting Error:** The ligature “ff” (as in the words: define, finite, field, fixed, satisfied, first, suffices, find) was omitted on the following pages:

15-32, 226, 227, 273-275, 278, 279, 598-611

The ligature “ff” was also omitted on some pages (as in the word *influential* on page 227, line -9).