

**A version of this problem was on exams A and B (May 5).**

- (15) 1. Suppose  $L_1$  is the straight line described parametrically by  $\begin{cases} x = 3t + 1 \\ y = 7t + 2 \\ z = -t + 1 \end{cases}$  and  $L_2$  is the straight line described parametrically by  $\begin{cases} x = 2s - 2 \\ y = s + 6 \\ z = -2s + 6 \end{cases}$ .
- a) The lines  $L_1$  and  $L_2$  intersect in a point. Find the point and verify that the lines intersect there.
- b) Find an equation for the plane containing  $L_1$  and  $L_2$ .

**A version of this problem was on exams C and D (May 10).**

- (15) 1. An equation for the plane  $P_1$  is  $2x - 2y + 3z = 3$  and an equation for the plane  $P_2$  is  $6x + y - 2z = 5$ . These planes are not parallel.
- a) What is the angle between  $P_1$  and  $P_2$ ? (The answer should be an “unsimplified” value of arccos.)
- b) Find a point which is on both  $P_1$  and  $P_2$ .
- c) Find parametric equations for the line which is the intersection of  $P_1$  and  $P_2$ .
- (15) 2. Suppose  $f(x, y, z) = x^2 e^{4y-z}$ .
- a) Compute  $\nabla f$ .
- b) Suppose  $p = (2, 1, 4)$ . Find the maximum rate of change of  $f$  at  $p$  and find a vector in the direction of this maximum increase.

Maximum rate of change: \_\_\_\_\_

Vector in the direction of that rate: \_\_\_\_\_


**This appeared on exams A and B (May 5).**

- c) Write parametric equations for the line normal to the level surface of  $f$  at  $p$ .

**This appeared on exams C and D (May 10).**

- c) Write an equation for the plane tangent to the level surface of  $f$  at  $p$ .

- (15) 3. Compute  $\iint_D x^4 y \, dA$  where  $D$  is the triangular region in the plane whose vertices are at  $(0, 0)$ ,  $(2, 2)$ , and  $(3, 2)$ .



- (20) 4. Suppose  $J = \int_0^1 \int_0^\pi \int_0^{\sqrt{1-z}} (z^3 \sin \theta) r \, dr \, d\theta \, dz$ , a triple integral of the function  $z^3 \sin \theta$  over a region in  $\mathbb{R}^3$  described using cylindrical coordinates.
- a) Sketch the solid region in  $\mathbb{R}^3$  over which the integral  $J$  is computed, and accompany your sketch with equations describing the boundary surfaces of the region using the standard rectangular coordinates  $x$ ,  $y$ , and  $z$ .
- b) Compute  $J$ .

- (15) 5. Suppose  $f(x, y) = (2y + 1)e^{(x^2 - y)}$ .  
 a)  $f(x, y)$  has one critical point. Find this point.

**You must check that you have found all the critical points, and that there is exactly one.**

b) Use the Second Derivative Test to determine the nature of the one critical point.

- (25) 6. The vector field  $\mathbf{T}$  is defined below. Find the flux of  $\mathbf{T}$  through the upper half ( $z \geq 0$ ) of the sphere of radius 5, centered at the origin and oriented outward.

$$\mathbf{T}(x, y, z) = \sin(z)\mathbf{i} + e^{\sqrt{z^4 + x^2}}\mathbf{j} + (x^2 + y^2 + z^2)\mathbf{k}$$

**Note** You may want to use the Divergence Theorem on a “simple” solid region to change the desired computation to the computation of a triple integral and a simpler flux integral.

- (20) 7. Suppose  $\mathbf{F}(x, y, z) = (y^2 z)\mathbf{i} + (2xyz)\mathbf{j} + (xy^2 + 4z)\mathbf{k}$ , a vector field defined and continuously differentiable throughout space.

a) Determine whether there is a scalar function  $P(x, y, z)$  defined everywhere in space such that  $\nabla P = \mathbf{F}$ . If there is such a  $P$ , find it; if there is not, explain why not.


b) Compute the integral  $\int_C \mathbf{F} \cdot \mathbf{t} \, ds$ , where  $C$  is the circular helix whose position vector is given by  $\mathbf{R}(t) = (\cos t)\mathbf{i} + (\sin t)\mathbf{j} + t\mathbf{k}$  for  $0 \leq t \leq 2\pi$ . Use information from your answer to a) to help if you wish.

- (15) 8. Evaluate the line integral  $\int_R (2xy^2 - 5 \cos x) \, dx + (-3x^2 - 7e^y) \, dy$  where  $R$  is the boundary of the rectangle with vertices  $(0, 1)$ ,  $(2, 1)$ ,  $(2, 2)$ , and  $(0, 2)$  oriented in the usual (counterclockwise) fashion.

**Note** Use *any* method to evaluate this integral. Some methods are easier than others.

- (20) 9. Sketch the three level curves of the function  $W(x, y) = x - y^2$  which pass through the points  $P = (3, -1)$  and  $Q = (0, 1)$  and  $R = (2, 1)$ . Be sure to label each curve with the appropriate function value and be sure that your drawing is clear and unambiguous.

Also sketch on the same axis the vectors of the gradient vector field  $\nabla W$  at the points  $P$  and  $Q$  and  $R$  and  $S$  and  $T$ . The point  $S = (0, -2)$  and  $T = (-2, 0)$ .



- (20) 10. Suppose  $R$  is the bounded region in the  $xy$ -plane defined by  $y = x^2$  and  $y = x + 2$ .  
 a) Sketch the region  $R$ .  
 b) If  $f(x, y)$  is a function defined in the region  $R$ , describe how to write  $\iint_R f(x, y) \, dA$  as a sum of one or more iterated  $\mathbf{dx} \, \mathbf{dy}$  integrals.  
 c) If  $f(x, y)$  is a function defined in the region  $R$ , describe how to write  $\iint_R f(x, y) \, dA$  as a sum of one or more iterated  $\mathbf{dy} \, \mathbf{dx}$  integrals.

- (12) 11. Find the curvature of the ellipse  $\begin{cases} x = 3 \cos t \\ y = 4 \sin t \end{cases}$  at the points  $(3, 0)$  and  $(0, 4)$ .
- (8) 12. Suppose  $z$  is defined implicitly as a function of  $x$  and  $y$  by the equation  $z^2x + 3xy^2 + e^{(y^2z)} = 4$ . Find  $\frac{\partial z}{\partial x}$ .

# Final Exam for Math 251, sections 5–10

May 5, 2006 or May 10, 2006

NAME \_\_\_\_\_

Please circle your section number: 5 6 7 8 9 10

Do all problems, in any order.

Show your work. An answer alone may not receive full credit.

No notes other than the distributed formula sheet may be used on this exam.

No calculators may be used on this exam.

Problem Number	Possible Points	Points Earned:
1	15	
2	15	
3	15	
4	20	
5	15	
6	25	
7	20	
8	15	
9	20	
10	20	
11	12	
12	8	
Total Points Earned:		