












Function	Domain	Range	Graph
x^2	all x	$y \geq 0$	
x^3	all x	all y	
\sqrt{x}	$x \geq 0$	$y \geq 0$	
$ x $	all x	$y \geq 0$	
$1/x$	$x \neq 0$	$y \neq 0$	
$\sin x$	all x	$-1 \leq y \leq 1$	
$\cos x$	all x	$-1 \leq y \leq 1$	
$\tan x$	$x \neq (\text{odd int})\frac{\pi}{2}$	all y	
$\ln x$	$x > 0$	all y	
e^x	all x	$y > 0$	
$\arctan x$	all x	$-\frac{\pi}{2} < y < \frac{\pi}{2}$	
$\arcsin x$	$-1 \leq x \leq 1$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$	

Function Derivative

C	0
x^n	nx^{n-1}
e^x	e^x
$\ln x$	$1/x$
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$(\sec x)^2$
$\arctan x$	$\frac{1}{1+x^2}$
$\arcsin x$	$\frac{1}{\sqrt{1-x^2}}$

Function Derivative

$Kf(x)$	$Kf'(x)$
$f(x)+g(x)$	$f'(x)+g'(x)$
$f(x) \cdot g(x)$	$f'(x) \cdot g(x) + f(x) \cdot g'(x)$
$\frac{f(x)}{g(x)}$	$\frac{f'(x)g(x) - g'(x)f(x)}{g(x)^2}$
$f(g(x))$	$f'(g(x)) \cdot g'(x)$


Logarithmic properties

$$\begin{aligned}\ln(a \cdot b) &= \ln a + \ln b & \ln(a^b) &= b \ln(a) \\ \ln(a/b) &= \ln(a) - \ln(b) & \ln(\frac{1}{b}) &= -\ln(b) \\ \ln(e^a) &= a & \ln(1) &= 0 & \ln(e) &= 1\end{aligned}$$

Exponential properties

$$\begin{aligned}a^{b+c} &= a^b \cdot a^c & a^{-b} &= 1/a^b \\ (a^b)^c &= a^{bc} & e &\approx 2.718 \\ a^0 &= 1 & e^{\ln a} &= a \text{ if } a > 0\end{aligned}$$

Triangle things



$$\begin{aligned}360^\circ (\text{degrees}) &= 2\pi \text{ radians} \\ \sin \theta &= \frac{OPP}{HYP} & \cos \theta &= \frac{ADJ}{HYP} & \tan \theta &= \frac{OPP}{ADJ} \\ \text{Pythagoras} & & (ADJ)^2 + (OPP)^2 &= (HYP)^2 & (\sin \theta)^2 + (\cos \theta)^2 &= 1\end{aligned}$$

θ	$\sin \theta$	$\cos \theta$	$\tan \theta$
0	0	1	0
$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$\frac{\pi}{2}$	1	0	NONE
π	0	-1	0

More formulas

The **roots** of $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Distance from (a, b) to (c, d) : $\sqrt{(a-c)^2 + (b-d)^2}$

Circle center (h, k) & radius r : $(x-h)^2 + (y-k)^2 = r^2$

Line $y = mx + b$ and $m = \frac{y_2 - y_1}{x_2 - x_1}$ (**slope** of the line)

Addition formulas $\sin(A+B) = \sin A \cos B + \cos A \sin B$

$\cos(A+B) = \cos A \cos B - \sin A \sin B$

Periodicity $\sin(x+2\pi) = \sin x$ and $\cos(x+2\pi) = \cos x$ and $\tan(x+\pi) = \tan x$ for all x

Area and Volume Formulas

Triangle $A = \frac{1}{2} \text{ BASE} \cdot \text{HEIGHT}$

Rectangle $A = \text{LENGTH} \cdot \text{WIDTH}$

Circle $A = \pi \text{ RADIUS}^2$

Circle $C = 2\pi \text{ RADIUS}$



Box $V = \text{LENGTH} \cdot \text{WIDTH} \cdot \text{HEIGHT}$

Cylinder $V = \pi \text{ RADIUS}^2 \cdot \text{HEIGHT}$



Sphere $A = 4\pi \text{ RADIUS}^2$

Sphere $V = \frac{4}{3}\pi \text{ RADIUS}^3$


w in f 's domain is a **critical number** if **either** $f'(w) = 0$: could look like



or $f'(w)$ doesn't exist: might look like \pm



or even like \pm if f isn't continuous at w .



The First Derivative Test A critical number w is a **relative max** if f' (left of w) > 0 & f' (right of w) < 0 ; **relative min** if f' (left of w) < 0 & f' (right of w) > 0 .

Important No other critical numbers should be between w and where the sign of f' is checked!



If both are positive or both are negative, then w is an **inflection point** of f .

The Second Derivative Test A critical number w is a **relative min** if $f''(w) > 0$ & **relative max** if $f''(w) < 0$.

Finding max/min on a closed interval

If f is continuous on $a \leq x \leq b$ then f 's max/min values must occur either at a or at b or at a critical number inside the interval.

f has an **inflection point** at w if w is in f 's domain and if the concavity of f 's graph is different on either side of w :

\pm  (here $f''(w) = 0$) or \pm  (here $f''(w)$ doesn't exist).

f is **continuous** at w if $\lim_{x \rightarrow w} f(x)$ exists and equals $f(w)$
or check $\lim_{x \rightarrow w^+} f(x)$ and $\lim_{x \rightarrow w^-} f(x)$ both exist and $= f(w)$.

f is **differentiable** at w if $\lim_{h \rightarrow 0} \frac{f(w+h)-f(w)}{h}$ exists. This is $f'(w)$: **the rate of change of f with respect to w** or **the slope of the tangent line** to $y = f(x)$ at $x = w$.

Implicit differentiation/related rates

Key point Differentiate a whole equation. Don't forget what's varying, chain rule, product rule, etc.

Example If $xy^2 = \sin(x+y) + 3x$ then $\frac{d}{dx}$ the equation. Get $1 \cdot y^2 + x \cdot 2yy' = \cos(x+y)(1+y') + 3$.

Solve for y' .

f defined in $a < x < b$ has a **relative maximum** at w in the interval if $f(w) \geq f(x)$ for x 's near w on both sides.

f defined in $a < x < b$ has a **relative minimum** at w in the interval if $f(w) \leq f(x)$ for x 's near w on both sides.

Relative max and min must occur **at critical numbers**.

Differential or tangent line approximation

$f(w+\Delta w) \approx f(w) + f'(w)\Delta w$. The graph's bending causes **error**: the true value is larger when the graph is concave up and smaller when the graph is concave down.

Intermediate Value Theorem If f is continuous in $a \leq x \leq b$, f 's values include all numbers between $f(a)$ and $f(b)$: a continuous function's graph has no jumps.

Mean Value Theorem If f is differentiable in $a \leq x \leq b$, there are some c 's in the interval with $f'(c) = \frac{f(b)-f(a)}{b-a}$: some tangent lines of a differentiable function's graph must be parallel to any chord.

Rolle's Theorem MVT with $f(a) = f(b) = 0$.


Fund. Thm. of Calculus If $F' = f$ then $\int_a^b f(x) dx = F(b) - F(a)$; $\frac{d}{dx} \int_a^x f = f(x)$.

f is **increasing** in $a < x < b$ if $f(x_1) \leq f(x_2)$ for any $x_1 \leq x_2$ in the interval. If $f'(x) > 0$ always in $a < x < b$ then f is increasing there.

f is **decreasing** in $a < x < b$ if $f(x_1) \geq f(x_2)$ for any $x_1 \leq x_2$ in the interval. If $f'(x) < 0$ always in $a < x < b$ then f is decreasing there.

f is **concave up** if lines connecting the graph are above the graph: it bends *up*. If $f''(x) > 0$ always in $a < x < b$, f is concave up.

f is **concave down** if lines connecting the graph are below the graph: it bends *down*. If $f''(x) < 0$ always in $a < x < b$, f is concave down.




Function	Antiderivative
$f(x)$	$\int f(x) dx$
$Kf(x)$	$K \int f(x) dx$
$f(x)+g(x)$	$\int f(x) dx + \int g(x) dx$
x^n	$\frac{1}{n+1}x^{n+1} + C, n \neq -1$
$\frac{1}{x}$	$\ln x + C \ (x > 0)$
e^x	$e^x + C$
$\sin x$	$-\cos x + C$
$\cos x$	$\sin x + C$
$\tan x$	$\ln(\sec x) + C$
$\int f(u) du$	$\int f(g(x)) g'(x) dx$
when $u = g(x)$ (Substitution)	

Sample points $\{c_1, c_2, c_3\}$ **Partition** $\{a = x_0 \leq x_1 \leq x_2 \leq x_3 = b\}$


Riemann sum $f(c_1)(x_1 - x_0) + f(c_2)(x_2 - x_1) + f(c_3)(x_3 - x_2)$

With many sample points and small differences in the partition, the sum will closely approximate the **definite integral** $\int_a^b f(x) dx$.




If $\lim_{x \rightarrow a(\pm)} f(x) = \pm\infty$ then $x = a$ is a **vertical asymptote** of $y = f(x)$ and if $\lim_{x \rightarrow \pm\infty} f(x) = b$ then $y = b$ is a **horizontal asymptote** of $y = f(x)$.

If $g(x) \leq f(x)$ in $a \leq x \leq b$, $\int_a^b g(x) dx \leq \int_a^b f(x) dx$.



$$\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx$$



$\int_a^b f(x) dx$ is **signed area**: area I - area II + area III.

