

Annals of Mathematics

Nonsymmetric Koornwinder Polynomials and Duality

Author(s): Siddhartha Sahi

Source: *The Annals of Mathematics*, Second Series, Vol. 150, No. 1 (Jul., 1999), pp. 267-282

Published by: *Annals of Mathematics*

Stable URL: <http://www.jstor.org/stable/121102>

Accessed: 11/06/2011 15:32

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=annals>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Annals of Mathematics is collaborating with JSTOR to digitize, preserve and extend access to *The Annals of Mathematics*.

<http://www.jstor.org>

Nonsymmetric Koornwinder polynomials and duality

By SIDDHARTH SAHI*

1. Introduction

In the fundamental work of Lusztig [L] on affine Hecke algebras, a special role is played by the root system of type \tilde{C}_n . The affine Hecke algebra is a deformation of the group algebra of an affine Weyl group which usually depends on as many parameters as there are distinct root lengths, i.e. one or two for an irreducible root system. However in the \tilde{C}_n case, the Hecke algebra H has three parameters, corresponding to the fact that there is a simple coroot which is divisible by 2.

Recently, Cherednik [C1]–[C3] has introduced the notion of a double affine Hecke algebra, and has used it to prove several conjectures on Macdonald polynomials. These polynomials, and Cherednik’s double affine Hecke algebra, involve two or three parameters, i.e., one more than the number of root lengths.

In this paper, motivated by the work of Noumi [N], we define a double affine Hecke algebra for the \tilde{C}_n case, which depends on three additional parameters, making six altogether. The associated orthogonal polynomials are precisely the six-parameter family of polynomials P_λ introduced by Koornwinder in [Ko].

These polynomials are themselves quite remarkable. *Every* symmetric Macdonald polynomial [M] associated to a classical root system (i.e. those of types A , B , C , D , and the two classes of type BC , not considered by Cherednik) can be obtained from the P_λ by a suitable limiting procedure [D]. Moreover, for $n = 1$, the P_λ become the Askey-Wilson polynomials, which sit atop an impressive hierarchy of orthogonal polynomials in one variable [AW].

Koornwinder and Macdonald have formulated several conjectures for these polynomials, which are analogous to those proved by Cherednik for Macdonald polynomials. These are the “constant term,” “norm,” “evaluation,” and

*This work was supported by an NSF grant, and carried out in part during the author’s visit to Japan. The author wishes to thank M. Wakayama, H. Ochiai, K. Mimachi for their hospitality, and M. Noumi for explaining the results of [N].

“duality” conjectures. For a certain five-parameter subfamily of the P_λ , these conjectures were proved by van Diejen in [D].

In the general (six-parameter) setting, van Diejen has shown that either of the last two conjectures implies the other three. One of the results in this paper is a proof of the duality conjecture which implies all the rest by van Diejen’s work.

Here is an outline of the paper: After a brief summary of the relevant results of Koornwinder and Noumi, we define the six-parameter double affine Hecke algebra \mathcal{H} and establish its basic properties, including the existence of an involution. Next, we introduce certain commutators S_i in \mathcal{H} , called the *intertwiners*, and use them to construct a family of polynomials $\{E_\alpha\}$. We call these the nonsymmetric Koornwinder polynomials, and we describe their relationship to the P_λ . Finally, we establish the duality conjecture for P_λ together with its analog for E_α .

A substantial part of this paper is directly motivated by the results of Cherednik in the two-/three-parameter setting. The idea of using intertwiners as creation operators was introduced in [K], [KS] and [S] for GL_n , and in [C4] for other root systems.

We have avoided one layer of notational complexity by identifying the coroot lattice of C_n with \mathbb{Z}^n . Thus we have suppressed explicit reference to roots and weights. Implicitly, though, these are ubiquitous.

We have also obtained fairly precise results concerning the orthogonality and triangularity of the nonsymmetric Koornwinder polynomials, which we shall report elsewhere.

Finally, we remark that according to the note added in proof to [D], Macdonald has informed van Diejen that he has proved the evaluation conjecture. By van Diejen’s work, this would also imply the duality conjecture for the P_λ , though not for the E_α .

2. Preliminaries

In this section we briefly recall some results of Koornwinder, Lusztig, and Noumi which we shall need. For more details the reader should consult [Ko], [L], and [N].

We fix six indeterminates q, t, t_0, t_n, u_0, u_n , and let \mathbb{F} be the field of rational functions in their *square roots*. We also define

$$(1) \quad a = t_n^{1/2} u_n^{1/2}, b = -t_n^{1/2} u_n^{-1/2}, c = q^{1/2} t_0^{1/2} u_0^{1/2}, d = -q^{1/2} t_0^{1/2} u_0^{-1/2}.$$

Let $\mathcal{R} = \mathbb{F}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ be the ring of Laurent polynomials in n variables over the field \mathbb{F} , and let \mathcal{S} be the subring consisting of *symmetric* polynomials, i.e. those which are invariant under permutations and inversions of the variables.

2.1. *Koornwinder polynomials.* In [Ko], Koornwinder defined a basis $\{P_\lambda\}$ of \mathcal{S} which is indexed by $\lambda \in \mathbb{Z}^n$ with $\lambda_1 \geq \dots \geq \lambda_n \geq 0$, and defined as follows:

Let T_{q,x_i} denote the i^{th} q -shift operator acting on $\mathcal{R} := \mathbb{F}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ by

$$T_{q,x_i} f(x_1, \dots, x_i, \dots, x_n) := f(x_1, \dots, qx_i, \dots, x_n).$$

Consider the following q -difference operator

$$D := \sum_{i=1}^n \Phi_i(x)(T_{q,x_i} - 1) + \sum_{i=1}^n \Phi_i(x^{-1})(T_{q,x_i}^{-1} - 1)$$

where $\Phi_i(x)$ is a rational function in x_1, \dots, x_n defined by

$$\Phi_i(x) := \frac{(1 - ax_i)(1 - bx_i)(1 - cx_i)(1 - dx_i)}{(1 - x_i^2)(1 - qx_i^2)} \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(1 - tx_i x_j^{-1})(1 - tx_i x_j)}{(1 - x_i x_j^{-1})(1 - x_i x_j)}.$$

Koornwinder showed that D preserves \mathcal{S} and is diagonalizable with distinct eigenvalues

$$d_\lambda = \sum_{i=1}^n \left[q^{-1} abcd t^{2n-i-1} (q^{\lambda_i} - 1) + t^{i-1} (q^{-\lambda_i} - 1) \right].$$

The Koornwinder polynomial P_λ is characterized by the equation

$$(2) \quad DP_\lambda = d_\lambda P_\lambda,$$

together with the condition that the coefficient of $x^\lambda := x_1^{\lambda_1} \cdots x_n^{\lambda_n}$ in P_λ is 1.

It turns out that Koornwinder's operator D is one among a commuting family of difference operators, all of which are simultaneously diagonalized by the P_λ . These higher operators were constructed abstractly by Noumi, and explicitly by van Diejen.

To describe the results of Noumi, we need to introduce some additional notation.

2.2. *The affine Weyl group.* The affine Weyl group W of type \tilde{C}_n is generated by elements s_0, s_1, \dots, s_n which satisfy $s_i^2 = 1$ and, for $n > 1$, also satisfy the braid relations

$$s_i s_j s_i \cdots = s_j s_i s_j \cdots$$

with two, three, or four terms on each side accordingly as i and j are connected by zero, one or two lines in the Coxeter graph

$$0 = 1 - 2 - \dots - (n-2) - (n-1) = n.$$

The (finite) Weyl group W_0 of type C_n is the subgroup generated by s_1, \dots, s_n .

W has a natural faithful affine action on $V = \mathbb{R}^n$ in which

$$(3) \quad s_n \cdot v = (v_1, \dots, v_{n-1}, -v_n), \quad s_0 \cdot v = (-v_1 - 1, v_2, \dots, v_n)$$

and the other s_i act by interchanging v_i and v_{i+1} . Indeed W is the semidirect product of W_0 and $\tau(\mathbb{Z}^n)$, where τ^v denotes the translation by v . In terms of the generators,

$$(4) \quad \tau^v = \tau_1^{v_1} \cdots \tau_n^{v_n}, \quad \tau_i = (s_i \cdots s_{n-1})(s_n \cdots s_0)(s_1^{-1} \cdots s_{i-1}^{-1}).$$

Since the W -action on V is affine, we get a representation of W on the space \tilde{V} of affine linear functionals on V , which we identify with $V \times \mathbb{R}\delta$ as follows:

$$(5) \quad \langle v + r\delta, v' \rangle := v'_1 v_1 + \cdots + v'_n v_n + r; \quad \langle w(v + r\delta), v' \rangle := \langle v + r\delta, w^{-1} \cdot v' \rangle.$$

This representation is given explicitly by

$$(6) \quad \begin{aligned} s_i(v + r\delta) &= s_i v + r\delta, \quad i \neq 0 \\ s_0(v + r\delta) &= (-v_1, v_2, \dots, v_n) + (r - v_1)\delta. \end{aligned}$$

We define an exponential map from the lattice $\mathbb{Z}^n \times \mathbb{Z}\delta \subset \tilde{V}$ to \mathcal{R} by

$$(7) \quad x^{v+k\delta} := q^{-k} x_1^{v_1} \cdots x_n^{v_n}, \quad v \in \mathbb{Z}^n, \quad k \in \mathbb{Z}.$$

Since the lattice is preserved under (6), we get a representation of W on \mathcal{R} by putting

$$(8) \quad w(x^{\tilde{v}}) := x^{w(\tilde{v})}; \quad \tilde{v} \in \mathbb{Z}^n \times \mathbb{Z}\delta.$$

Then W acts by algebra homomorphisms and $\mathcal{S} = \mathcal{R}^{W_0}$. Explicitly we have

$$(9) \quad \begin{aligned} s_0 f(x) &= f(qx_1^{-1}, x_2, \dots, x_n) \\ s_i f(x) &= f(x_1, \dots, x_{i+1}, x_i, \dots, x_n) \quad i \neq 0, n \\ s_n f(x) &= f(x_1, \dots, x_{n-1}, x_n^{-1}) \\ \tau_i &= T_{q,x_i}. \end{aligned}$$

2.3. The affine Hecke algebra. The affine Hecke algebra H of type \tilde{C}_n is generated over \mathbb{F} by elements T_0, T_1, \dots, T_n which satisfy the same braid relations as the s_i , and also satisfy

$$T_i - T_i^{-1} = t_i^{1/2} - t_i^{-1/2}$$

where $t_1 = \cdots = t_{n-1} = t$.

The elements T_1, \dots, T_n generate the (finite) Hecke algebra H_0 of type C_n . H and H_0 have natural bases $\{T_w\}$ consisting of w in W and W_0 , respectively, where

$$(10) \quad T_w = T_{i_1} \cdots T_{i_l}$$

if $w = s_{i_1} \cdots s_{i_l}$ is a *reduced* (i.e. shortest) expression of w in terms of the s_i .

The analogs of the translations τ_i in (4) are the elements

$$(11) \quad Y_i = (T_i \cdots T_{n-1})(T_n \cdots T_0)(T_1^{-1} \cdots T_{i-1}^{-1}), \quad i = 1, \dots, n.$$

Lusztig [L] has shown that the Y_i commute pairwise and generate a sub-algebra

$$\mathcal{R}_Y \approx \mathbb{F}[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}],$$

and that multiplication gives us a vector space isomorphism

$$(12) \quad H_0 \otimes \mathcal{R}_Y \approx H.$$

2.4. The Noumi representation. Let s_i act on \mathcal{R} by (9); then Noumi in [N] has shown that the following map π extends to a representation of H on \mathcal{R} :

$$(13) \quad \begin{aligned} \pi(T_0^{\pm 1}) &:= t_0^{\pm 1/2} + t_0^{-1/2} \frac{(1 - cx_1^{-1})(1 - dx_1^{-1})}{1 - qx_1^{-2}} (s_0 - 1) \\ \pi(T_i^{\pm 1}) &:= t_i^{\pm 1/2} + t_i^{-1/2} \frac{(1 - t_i x_i x_{i+1}^{-1})}{(1 - x_i x_{i+1}^{-1})} (s_i - 1) \quad i \neq 0, n \\ \pi(T_n^{\pm 1}) &:= t_n^{\pm 1/2} + t_n^{-1/2} \frac{(1 - ax_n)(1 - bx_n)}{1 - x_n^2} (s_n - 1), \end{aligned}$$

where a, b, c, d are as in (1).

Moreover if f is in \mathcal{S} then $\pi(f) := \pi(f(Y_1, \dots, Y_n))$ preserves \mathcal{S} . Noumi showed that the restriction of $\pi(Y_1 + \cdots + Y_n + Y_1^{-1} + \cdots + Y_n^{-1})$ to \mathcal{S} is a linear combination of the Koornwinder operator D and a scalar. This means that the Koornwinder polynomials are simultaneous eigenfunctions for the $\pi(f)$. More precisely, P_λ is characterized by

$$(14) \quad \pi(f)P_\lambda(x) = f(q^{\lambda+\rho})P_\lambda(x),$$

where $q^{\lambda+\rho}$ means $(q^{\lambda_1+\rho_1}, \dots, q^{\lambda_n+\rho_n})$ and ρ is defined by

$$(15) \quad q^{\rho_i} = st^{n-i}, \quad \text{with } s := (t_0 t_n)^{1/2} = \sqrt{q^{-1}abcd}.$$

Remark. The fact that π extends to a representation can be derived from Proposition 3.6 in [L], along the lines suggested in Proposition 4.6 of [M].

3. The double affine Hecke algebra

We now introduce the algebra \mathcal{H} which will be the principal object of study in this paper. For convenience, we will write $Z \sim z$ as an abbreviation for the relation

$$Z - Z^{-1} = z^{1/2} - z^{-1/2}.$$

Definition. Let \mathcal{H} be the algebra generated over \mathbb{F} by elements $T_i^{\pm 1}$, $i = 0, \dots, n$, and commuting elements $X_i^{\pm 1}$, $i = 1, \dots, n$, subject to the relations:

- (i) $T_i \sim t_i$,
- (ii) the \tilde{C}_n braid relations for the T_i 's,
- (iii) $T_i X_j = X_j T_i$ if $|i - j| > 1$, or if $i = n$ and $j = n - 1$,
- (iv) $T_i X_i = X_{i+1} T_i^{-1}$, $i = 1, \dots, n - 1$,
- (v) $X_n^{-1} T_n^{-1} \sim u_n$,
- (vi) $U_0 \equiv q^{-1/2} T_0^{-1} X_1 \sim u_0$.

If we set $u_0 = u_n = 1$ and $t_0 = t_n$, then \mathcal{H} specializes to the three-parameter double affine Hecke algebra considered in [C1] for the affine root system \tilde{C}_n .

Our definition is motivated by the following considerations:

Define a map π from the generators of \mathcal{H} to $\text{End}(\mathcal{R})$ by letting $\pi(T_i^{\pm 1})$ be as in (13), and letting $\pi(X_i^{\pm 1})$ be the operator of multiplication by $x_i^{\pm 1}$.

3.1. THEOREM. *The map π extends to a representation of \mathcal{H} on \mathcal{R} .*

Proof. We need only verify that (i)–(vi) hold for $\pi(T_i^{\pm 1})$ and $\pi(X_i^{\pm 1})$.

The relations (i) and (ii) follow from Noumi's result, and (iii) and (iv) are easily verified using the formulas. For (v) we have

$$\begin{aligned} X_n^{-1} T_n^{-1} &\mapsto t_n^{-1/2} x_n^{-1} + t_n^{-1/2} \frac{(1 - ax_n)(1 - bx_n)}{1 - x_n^2} (x_n^{-1} s_n - x_n^{-1}) \\ T_n X_n &\mapsto t_n^{1/2} x_n + t_n^{-1/2} \frac{(1 - ax_n)(1 - bx_n)}{1 - x_n^2} (s_n x_n - x_n). \end{aligned}$$

Since $s_n x_n = x_n^{-1} s_n$ and $x_n - x_n^{-1} = -x_n^{-1}(1 - x_n^2)$, we get

$$\begin{aligned} X_n^{-1} T_n^{-1} - T_n X_n &\mapsto t_n^{-1/2} x_n^{-1} - t_n^{1/2} x_n - t_n^{-1/2} x_n^{-1} (1 - ax_n)(1 - bx_n) \\ &= -(t_n^{1/2} + t_n^{-1/2} ab)x_n + t_n^{-1/2}(a + b). \end{aligned}$$

Substituting $a = t_n^{1/2} u_n^{1/2}$ and $b = -t_n^{1/2} u_n^{-1/2}$, this becomes $u_n^{1/2} - u_n^{-1/2}$ proving (v).

Relation (vi) can be proved similarly using $s_0 x_1 = qx_1^{-1} s_0$. \square

3.2. THEOREM. *The representation π is faithful.*

Proof. We first note that in any word in \mathcal{H} involving the generators, the relations (i)–(vi) allow us to commute the T_i 's past the X_j 's. Thus every element of \mathcal{H} can be written as a linear combination of $X^\alpha T_w$, where $X_\alpha = X_1^{\alpha_1} \cdots X_n^{\alpha_n}$ and T_w is as in (10).

Suppose a *nontrivial* linear combination maps to 0 under π . We then get

$$\sum c_{w,\alpha} x^\alpha \pi(T_w) = 0$$

in $\text{End}(\mathcal{R})$, where $c_{w,\alpha}$ are scalars in \mathbb{F} , not all zero.

The left side is a rational expression in the square roots of q, t, t_0, t_n, u_0, u_n , and we consider what happens if we specialize the last five indeterminates to 1. By clearing denominators and eliminating common factors, we may ensure that at least *some* of the $c_{w,\alpha}$ have nonzero specializations; by (1), (10), and (13), $\pi(T_w)$ specializes to the action of w as in (9).

Thus we get a nontrivial dependence relation in $\text{End}(\mathcal{R})$ of the form

$$\sum c_{w,\alpha} x^\alpha w = 0, \quad w \in W, \alpha \in \mathbb{Z}^n.$$

Since $W = W_0\tau(\mathbb{Z}^n)$, we can rewrite this as

$$\sum c_{w,\alpha,\beta} x^\alpha w \tau^\beta = 0, \quad w \in W_0, \alpha, \beta \in \mathbb{Z}^n.$$

Collecting the terms for β , we get

$$\sum x^\alpha w p_{w,\alpha}(\tau_1, \dots, \tau_n) = 0, \quad w \in W_0, \alpha \in \mathbb{Z}^n,$$

where $p_{w,\alpha}(x)$ is the Laurent polynomial $\sum_\beta c_{w,\alpha,\beta} x^\beta$. Since $\tau_i(x^\gamma) = (q^{\gamma_i})x^\gamma$, applying the expression to x^γ we obtain

$$\sum x^{\alpha+w\gamma} p_{w,\alpha}(q^{\gamma_1}, \dots, q^{\gamma_n}) = 0.$$

It follows that $p_{w,\alpha}(q^{\gamma_1}, \dots, q^{\gamma_n}) = 0$ for all γ in \mathbb{Z}^n outside the *finite* union of hyperplanes determined by the conditions $\alpha + w\gamma = \alpha' + w'\gamma$ for α, α', w, w' occurring in the last expression. But then $p_{w,\alpha}$ must be identically 0, and we conclude that all $c_{w,\alpha,\beta} = 0$, contrary to the assumption. \square

Let us define $\mathcal{R}_X := \mathbb{F}[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$; then the above proof shows:

3.3. COROLLARY. *The natural maps from H , H_0 , \mathcal{R}_Y and \mathcal{R}_X into \mathcal{H} are injective.*

We shall identify the above algebras with their images in \mathcal{H} . Then we have:

3.4. COROLLARY. *The multiplication maps from $\mathcal{R}_X \otimes H$ and $\mathcal{R}_X \otimes H_0 \otimes \mathcal{R}_Y$ into \mathcal{H} are linear isomorphisms.*

We conclude this section by giving an intrinsic definition of the representation π . First, by the definition of H , it is clear that the map

$$(16) \quad \chi : T_i \mapsto t_i^{1/2}, \quad i = 0, \dots, n$$

extends to a one-dimensional character of H .

3.5. PROPOSITION. *The representation π is isomorphic to $\text{Ind}_H^{\mathcal{H}}(\chi)$.*

Proof. The induced representation is \mathcal{H}/\mathcal{I} where \mathcal{I} is the left ideal of \mathcal{H} generated by $T_i - t_i^{1/2}$, $i = 0, \dots, n$. On the other hand, $\pi \approx \mathcal{H}/\mathcal{J}$ where \mathcal{J} is the annihilator of the cyclic vector $1 \in \mathcal{R}$. It remains to show that $\mathcal{I} = \mathcal{J}$.

First, since $s_i(1) = 1$ it follows from formula (7) that $\pi(T_i - t_i^{1/2})(1) = 0$, and so

$$\mathcal{I} \subseteq \mathcal{J}.$$

Next consider the left ideal I in H generated by $T_i - t_i^{1/2}$. Then we have $H = \mathbb{F} + I$; thus $\mathcal{R}_X \otimes H = \mathcal{R}_X \otimes \mathbb{F} + \mathcal{R}_X \otimes I$. Applying Corollary 3.4 we conclude that

$$\mathcal{H} = \mathcal{R}_X + \mathcal{I}.$$

This mean that the isomorphism $\mathcal{R}_X \rightarrow \mathcal{R}$ given by $X_i \mapsto x_i = \pi(X_i) \cdot 1$ can be factored as the following sequence of *surjective* maps

$$\mathcal{R}_X \rightarrow \mathcal{H}/\mathcal{I} \rightarrow \mathcal{H}/\mathcal{J} \rightarrow \mathcal{R}.$$

In particular, the middle map is an isomorphism, and so $\mathcal{I} = \mathcal{J}$. \square

4. The involution

Let ε denote the involution on \mathbb{F} which sends q, t, t_n, u_0 to their inverses, and which maps

$$t_0 \mapsto u_n^{-1}.$$

We shall show that ε extends to an involution on \mathcal{H} . First we prove

4.1. LEMMA. *Let $U_n \equiv X_1^{-1}T_0Y_1^{-1} = X_1^{-1}T_1^{-1} \cdots T_n^{-1} \cdots T_1^{-1}$, then $U_n \sim u_n$.*

Proof. By (iv) we have $X_i^{-1}T_i^{-1} = T_iX_{i+1}^{-1}$, and applying this repeatedly, we get

$$U_n = (T_1 \cdots T_{n-1})(X_n^{-1}T_n^{-1})(T_{n-1}^{-1} \cdots T_1^{-1}).$$

Thus U_n is conjugate to $X_n^{-1}T_n^{-1}$ and the lemma follows from relation (v). \square

4.2. THEOREM. *The map ε extends to an involution of \mathcal{H} which maps X_i to Y_i , sends T_1, \dots, T_n to their inverses, and maps*

$$T_0 \mapsto U_n^{-1}.$$

Proof. We first verify that the ε -transforms of (i)–(vi) hold in \mathcal{H} . For $i \neq 0$ the relation (i) becomes $T_i^{-1} \sim t_i^{-1}$, which is implied by $T_i \sim t_i$; while for $i = 0$ it becomes $U_n^{-1} \sim u_n^{-1}$, which follows from Lemma 4.1.

All the ε -transforms of the braid relations (ii) are immediate, except for

$$U_n^{-1}T_1^{-1}U_n^{-1}T_1^{-1} \stackrel{?}{=} T_1^{-1}U_n^{-1}T_1^{-1}U_n^{-1}.$$

We shall check this directly. Write $\Phi = T_2 \cdots T_n \cdots T_2$; then $X_1\Phi = \Phi X_1$, and we get

$$\begin{aligned} T_1^{-1}U_n^{-1}T_1^{-1}U_n^{-1} &= \Phi T_1 X_1 \Phi T_1 X_1 = \Phi T_1 \Phi X_1 T_1 X_1 = \Phi T_1 \Phi X_1 X_2 T_1^{-1} \\ U_n^{-1}T_1^{-1}U_n^{-1}T_1^{-1} &= T_1 \Phi T_1 X_1 \Phi T_1 X_1 T_1^{-1} = T_1 \Phi T_1 \Phi X_1 T_1 X_1 T_1^{-1} \\ &= T_1 \Phi T_1 \Phi T_1^{-1} X_1 X_2 T_1^{-1}. \end{aligned}$$

By multiplying both sides by $T_1 X_1^{-1} X_2^{-1} T_1$ on the right, it suffices to show $(T_2 \cdots T_n \cdots T_2)T_1(T_2 \cdots T_n \cdots T_2)T_1 \stackrel{?}{=} T_1(T_2 \cdots T_n \cdots T_2)T_1(T_2 \cdots T_n \cdots T_2)$.

We apply $T_2 T_1 T_2 = T_1 T_2 T_1$ to both sides and commute the resulting T_1 's as far to the extremes as possible. Using $T_1 T_2 T_1 = T_2 T_1 T_2$ once on each side and cancelling, we get

$$(T_3 \cdots T_n \cdots T_3)T_2(T_3 \cdots T_n \cdots T_3)T_2 \stackrel{?}{=} T_2(T_3 \cdots T_n \cdots T_3)T_2(T_3 \cdots T_n \cdots T_3).$$

Iterating, we reach the *true* relation $T_n T_{n-1} T_n T_{n-1} = T_{n-1} T_n T_{n-1} T_n$, which proves (ii).

The ε -transforms of (iii)–(iv) are easily checked. For (v) we get

$$\varepsilon(X_n^{-1}T_n^{-1}) = Y_n^{-1}T_n = (T_{n-1} \cdots T_1)T_0^{-1}(T_1^{-1} \cdots T_{n-1}^{-1})$$

which is conjugate to T_0^{-1} . Hence the desired relation follows from $T_0 \sim t_0$.

Finally, for (vi) we have

$$\varepsilon(U_0) = q^{1/2}U_n Y_1 = q^{1/2}(X_1^{-1}T_0 Y_1^{-1})Y_1 = q^{1/2}X_1^{-1}T_0 = U_0^{-1}.$$

Thus (vi) follows from its original counterpart.

It follows that ε is a homomorphism, and it remains only to prove that $\varepsilon^2 = 1$. Since \mathcal{H} is generated by $\{T_i, X_i, Y_i \mid i = 1, \dots, n\}$, it suffices to show that $\varepsilon(Y_i) = X_i$ for all i . But

$$\varepsilon(Y_1) = T_1^{-1} \cdots T_n^{-1} \cdots T_1^{-1} U_n^{-1} = X_1.$$

Since $T_i X_i T_i = X_{i+1}$ and $T_i^{-1} Y_i T_i^{-1} = Y_{i+1}$, the result follows for all i by induction. \square

5. Intertwiners

In this section we introduce certain commutators in \mathcal{H} , and prove that they enjoy a crucial intertwining property with respect to the commutative family \mathcal{R}_Y .

Definition. We define operators S_i in \mathcal{H} as follows:

$$(17) \quad S_i := [T_i, Y_i], i = 1, \dots, n; \quad S_0 := [Y_1, U_n].$$

We also introduce the following notation, analogous to (7):

$$(18) \quad X^{v+k\delta} := q^{-k} X_1^{v_1} \cdots X_n^{v_n}; \quad Y^{v+k\delta} := q^k Y_1^{v_1} \cdots Y_n^{v_n}, \quad v \in \mathbb{Z}^n, k \in \mathbb{Z}.$$

5.1. THEOREM. *For all \tilde{v} in $\mathbb{Z}^n \times \mathbb{Z}\delta$,*

$$(19) \quad Y^{\tilde{v}} S_i = S_i Y^{s_i(\tilde{v})}; \quad i = 0, \dots, n.$$

Proof. By Theorems 3.2 and 4.2, it is enough to prove the $\pi \circ \varepsilon$ transform of (19). Applying $\pi \circ \varepsilon$ to (18) and (17), we get:

$$\begin{aligned} \pi \circ \varepsilon(Y^{\tilde{v}}) &= x^{\tilde{v}}; \quad \pi \circ \varepsilon(Y^{s_i(\tilde{v})}) = x^{s_i(\tilde{v})} \\ \pi \circ \varepsilon(S_0) &= [x_1, \pi(T_0^{-1})]; \quad \pi \circ \varepsilon(S_i) = [\pi(T_i^{-1}), x_i], \quad i = 1, \dots, n. \end{aligned}$$

Now, an easy calculation in $\text{End}(\mathcal{R})$, using formulas (13) gives

$$(20) \quad \pi \circ \varepsilon(S_i) = \phi_i(x)s_i; \quad \phi_i(x) = \begin{cases} t_0^{-1/2} x_1(1 - cx_1^{-1})(1 - dx_1^{-1}) & i = 0 \\ t_n^{-1/2} x_n^{-1}(1 - ax_n)(1 - bx_n) & i = n \\ t_i^{-1/2} x_{i+1}(1 - t_i x_i x_{i+1}^{-1}) & i \neq 0, n. \end{cases}$$

Thus the $\pi \circ \varepsilon$ transform of (19) becomes the following assertion in $\text{End}(\mathcal{R})$:

$$x^{\tilde{v}} \phi_i(x)s_i \stackrel{?}{=} \phi_i(x)s_i x^{s_i(\tilde{v})}.$$

After cancelling the ϕ_i , this follows from (8). \square

5.2. COROLLARY. *Let a', b', c', d' be the ε transforms of a, b, c, d , then*

$$(21) \quad S_i^2 = \begin{cases} u_n q^{-1} (1 - c' Y_1^{-1})(1 - d' Y_1^{-1})(1 - qc' Y_1)(1 - qd' Y_1) & i = 0 \\ t_n (1 - a' Y_n)(1 - b' Y_n)(1 - a' Y_n^{-1})(1 - b' Y_n^{-1}) & i = n \\ t_i Y_i Y_{i+1} (1 - t_i^{-1} Y_i Y_{i+1}^{-1})(1 - t_i^{-1} Y_i^{-1} Y_{i+1}) & i \neq 0, n. \end{cases}$$

Proof. By (20), we get $\pi \circ \varepsilon(S_i^2) = \phi_i s_i \phi_i s_i = \phi_i s_i (\phi_i) s_i^2 = \phi_i s_i (\phi_i)$, and the result follows from the explicit formula for ϕ_i . \square

In the next section we will use the S_i 's as creation operators for the E_α , starting with the constant function 1, which is an eigenfunction of Y_i , satisfying

$$(22) \quad \pi(Y_i)(1) = q^{\rho_i}(1); \quad i = 1, \dots, n.$$

This is an immediate consequence of the equation $\pi(T_i)(1) = t_i^{1/2}(1)$ and the definitions of Y_i and ρ in (11) and (15). To describe the other eigenvalues, we proceed as follows:

Definition. For α in \mathbb{Z}^n , we define

$w_\alpha :=$ the shortest element in W_0 such that $w_\alpha^{-1} \cdot \alpha$ is a partition;

$\bar{\alpha} := \alpha + w_\alpha \cdot \rho$ where ρ is as in (15);

$\mathcal{R}_\alpha :=$ the space of all $f \in \mathcal{R}$ satisfying $Y_i f = q^{\alpha_i + (w_\alpha \cdot \rho)_i} f$ for all i .

Alternatively, w_α in $W_0 = (\pm 1)^n S_n$ can be described as $w_\alpha := \sigma_\alpha \pi_\alpha$, where $\sigma_\alpha \in (\pm 1)^n$ is simply $(\text{sgn}(\alpha_1), \dots, \text{sgn}(\alpha_n))$ with $\text{sgn}(0)$ defined to be 1; and π_α is the permutation in S_n defined as follows: order the indices *first* by decreasing $|\alpha_i|$, *then* for fixed $|\alpha_i|$ from left to right for $\alpha_i \geq 0$, and *finally* from right to left for $\alpha_i < 0$.

For example if $\alpha = (-2, 2, 1, -1, 0, 1, -1)$, then $\sigma_\alpha = (-1, 1, 1, -1, 1, 1, -1)$ and π_α is the permutation $(2, 1, 3, 6, 7, 4, 5)$.

5.3. THEOREM. *If $s_i \cdot \alpha \neq \alpha$, then $\pi(S_i)$ is a linear isomorphism from \mathcal{R}_α to $\mathcal{R}_{s_i \cdot \alpha}$.*

Proof. Let $f \in \mathcal{R}_\alpha$. Then by (5) and (18) it follows that for all \tilde{v} in $\mathbb{Z}^n \times \mathbb{Z}\delta$,

$$Y^{\tilde{v}}(f) = q^{\langle \tilde{v}, \alpha + w_\alpha \cdot \rho \rangle} f.$$

Let us write $\bar{\alpha} = \alpha + w_\alpha \cdot \rho$. Then from Theorem 5.1 and (3) we get

$$\pi(Y^{\tilde{v}})\pi(S_i)f = \pi(S_i)\pi(Y^{s_i(\tilde{v})})f = q^{\langle s_i(\tilde{v}), \bar{\alpha} \rangle} \pi(S_i)f = q^{\langle \tilde{v}, s_i \cdot \bar{\alpha} \rangle} \pi(S_i)f.$$

Thus to prove that $\pi(S_i)f \in \mathcal{R}_{s_i \cdot \alpha}$, it suffices to show that $s_i \cdot \alpha \neq \alpha$ implies

$$(23) \quad s_i \cdot \bar{\alpha} = \overline{s_i \cdot \alpha}; \quad i = 0, \dots, n.$$

For $i = 0$, write $\beta = s_0 \cdot \alpha = (-\alpha_1 - 1, \alpha_2, \dots, \alpha_n)$. Then we claim that the permutations π_β and π_α are the *same*. Indeed if α_1 is positive then, in the ordering corresponding to π_α , the index 1 is the first among the indices j with $|\alpha_j| = \alpha_1$; while in the ordering for π_β , 1 is the last index in the higher group of indices satisfying $|\beta_j| = \alpha_1 + 1$. Thus the relative position of 1 with respect to other indices stays the same, as do those of other indices with respect to each other. A similar argument works if α_1 is negative, and taking into account the sign change we conclude that:

$$\bar{\beta}_1 = \beta_1 + (w_\beta \cdot \rho)_1 = -\alpha - 1 - (w_\alpha \cdot \rho)_1 = -\bar{\alpha}_1 - 1; \quad \bar{\beta}_i = \bar{\alpha}_i, \quad i \geq 2,$$

which is precisely the content of (23) for $i = 0$.

The argument for $i \geq 1$ is similar and simpler. We observe that if $\beta := s_i \cdot \alpha$ is different from α , then w_β equals $s_i w_\alpha$. Since s_i acts linearly, we get

$$s_i \cdot \bar{\alpha} = s_i \cdot \alpha + s_i \cdot (w_\alpha \cdot \rho) = \beta + w_\beta \cdot \rho = \bar{\beta} = \overline{s_i \cdot \alpha}.$$

Thus we conclude that $\pi(S_i)$ maps \mathcal{R}_α into $\mathcal{R}_{s_i \cdot \alpha}$ for all $i \geq 0$. But, by Corollary 5.2, S_i^2 is in \mathcal{R}_Y ; hence $\pi(S_i^2)$ acts by a scalar c_i on \mathcal{R}_α , which can be readily computed by substituting $Y_i = \bar{\alpha}_i$ in (21). In particular, we see that if $s_i \cdot \alpha \neq \alpha$ then c_i is not zero. Thus $\pi(S_i)$ is a linear isomorphism from \mathcal{R}_α to $\mathcal{R}_{s_i \cdot \alpha}$, with inverse $c_i^{-1}\pi(S_i)$. \square

6. Nonsymmetric Koornwinder polynomials

In this section we will define the nonsymmetric Koornwinder polynomials. The crucial result is:

6.1. THEOREM. *The spaces \mathcal{R}_α are all one-dimensional.*

Proof. We first prove that the spaces \mathcal{R}_α are nonzero. For $\alpha = 0$, w_α is the identity in W_0 and so $\bar{\alpha} = \rho$. Thus by (22), the constant functions belong to \mathcal{R}_α for $\alpha = 0$. For other $\alpha \in \mathbb{Z}^n$ we use Theorem 5.3 together with the fact that the affine action of W on \mathbb{Z}^n is transitive.

Now let $f = \sum_{c_\beta \in \mathbb{F}} c_\beta x^\beta$ be a nonzero function in \mathcal{R}_α . Then f satisfies

$$\pi(Y_i)f := q^{(\alpha+w_\alpha \cdot \rho)_i} f; \quad i = 1, \dots, n.$$

As in the proof of Theorem 3.2, we set t, t_0, t_n, u_0, u_n equal to 1 in the expression. Then ρ specializes to the zero vector, and Y_i specializes to $\pi(\tau_i) = T_{q, x_i}$. Clearing denominators and eliminating common factors, we may also assume that the c_β have finite specializations, not all zero. Letting $g \neq 0$ denote the specialization of f , we get

$$T_{q, x_i} g = q^{\alpha_i} g$$

which means that g is a nonzero multiple of x^α .

In particular, the coefficient c_α has a nonzero specialization and so must be nonzero. The result follows, since if there were two linearly independent functions in \mathcal{R}_α , we could construct a nonzero f with $c_\alpha = 0$. \square

The proof of the theorem shows that a function f in \mathcal{R}_α is uniquely determined by the knowledge of the coefficient of x^α in f .

Definition. The nonsymmetric Koornwinder polynomial E_α is the unique polynomial in \mathcal{R}_α in which the coefficient of x^α is 1.

6.2. THEOREM. *The polynomials E_α form a basis for \mathcal{R} over \mathbb{F} .*

Proof. Let us consider the degree filtration $\mathcal{R}_{(0)} \subseteq \mathcal{R}_{(1)} \subseteq \dots \subseteq \mathcal{R}$, where $\mathcal{R}_{(k)}$ is spanned by all monomials x^α with $|\alpha| := |\alpha_1| + \dots + |\alpha_n| \leq k$. We claim that for $|\alpha| \leq k$,

$$\mathcal{R}_\alpha \subseteq \mathcal{R}_{(k)}.$$

For $\alpha = 0$, we use Theorem 6.1 to see that \mathcal{R}_α consists precisely of the constants, which lie in $\mathcal{R}_{(0)}$. For other α , we observe that by (13) the filtration is invariant under T_i and Y_i , while $U_n = X_1^{-1}T_0Y_1^{-1}$ raises degree by at most one. Thus S_1, \dots, S_n preserve the filtration while S_0 raises degree by at most one. Now any α can be obtained from 0 by applying a sequence of s_i 's in which s_0 occurs exactly $|\alpha|$ times. Applying the corresponding S_i 's to \mathcal{R}_0 , we deduce the claim.

It follows that the set $\{E_\alpha : |\alpha| \leq k\}$ is contained in $\mathcal{R}_{(k)}$ and has the same cardinality as the monomial basis. Therefore it suffices to prove that the E_α are linearly independent. For this we choose a polynomial f in \mathcal{R} which takes distinct values on the finite set $\{q^{\bar{\alpha}} : |\alpha| \leq k\}$. Then the E_α belong to distinct eigenspaces under the operator $\pi(f(Y_1, \dots, Y_n))$ and hence are linearly independent. \square

6.3. COROLLARY. *The representation π is irreducible.*

Proof. Let \mathcal{V} be a $\pi(\mathcal{H})$ -invariant subspace of \mathcal{R} , and suppose $f = \sum c_\alpha E_\alpha$ belongs to \mathcal{V} , with some $c_\beta \neq 0$. Choose a function g in \mathcal{R} such that $g(q^{\bar{\beta}}) = 1/c_\beta$, and $g(q^{\bar{\alpha}}) = 0$ for all other α for which $c_\alpha \neq 0$. Then applying $\pi(g(Y_1, \dots, Y_n))$ to f we conclude that E_β belongs to \mathcal{V} . Now applying the $\pi(S_i)$'s we conclude that every E_α belongs to \mathcal{V} . \square

Next we consider the restriction of π to H .

Definition. For each partition λ we write \mathcal{R}^λ for the subspace of \mathcal{R} spanned by the $\{E_\alpha : \alpha \in W_0 \cdot \lambda\}$.

6.4. COROLLARY. *The \mathcal{R}^λ are irreducible $\pi(H)$ -modules, and \mathcal{R} is their direct sum.*

Proof. For the irreducibility, we repeat the previous argument without involving S_0 . The second assertion follows from Theorem 6.2. \square

Finally we discuss the connection with the symmetric Koornwinder polynomials P_λ .

6.5. COROLLARY. *The symmetric Koornwinder polynomial P_λ can be characterized as the unique W_0 -invariant polynomial in \mathcal{R}^λ which has the coefficient of x^λ equal to 1.*

Proof. We note that if $\alpha \in W_0 \cdot \lambda$, then $\bar{\alpha} = \alpha + w_\alpha \cdot \rho = w_\alpha \cdot (\lambda + \rho)$. In particular, if f is in \mathcal{S} , then $f(\bar{\alpha}) = f(\lambda + \rho)$, and so \mathcal{R}^λ is precisely the $f(\lambda + \rho)$ -eigenspace of $\pi(f(Y_1, \dots, Y_n))$ for $f \in \mathcal{S}$. The result now follows from the characterization (14). \square

Definition. Define $C \in H_0$ by $C := (\sum_{w \in W_0} \chi(T_w)^2)^{-1} \sum_{w \in W_0} \chi(T_w) T_w$.

6.6. COROLLARY. $\pi(C)$ is a projection from \mathcal{R}^λ to $\mathbb{F}P_\lambda$.

Proof. First of all, an easy calculation as in Lemma 2.5 of [S] shows that $T_i C = t_i^{1/2} C$ for $i = 1, \dots, n$; hence $\pi(T_i)\pi(C)f = t_i^{1/2}\pi(C)f$ for all $f \in \mathcal{R}$. By (13), this implies that $\pi(C)f$ is W_0 -invariant, and so must be a multiple of P_λ .

Moreover, for $f \in \mathcal{S}$, $\pi(T_w)f = \chi(T_w)f$; hence $\pi(C)$ acts by the identity on \mathcal{S} . \square

7. Duality

Let \dagger denote the involution on \mathbb{F} which sends q, t, t_0, t_n, u_0, u_n to their inverses.

7.1. PROPOSITION. *The map \dagger extends to an anti-involution on \mathcal{H} satisfying*

$$T_i^\dagger = T_i^{-1}, \quad X_i^\dagger = X_i^{-1}, \quad Y_i^\dagger = Y_i^{-1}.$$

Proof. For the proof we merely observe that each defining relation of \mathcal{H} is \dagger -invariant. \square

Definition. We define the *duality* anti-involution $*$ on \mathcal{H} by

$$h^* = \varepsilon(h^\dagger) = \varepsilon(h)^\dagger, \quad h \in \mathcal{H}.$$

On \mathbb{F} , $*$ simply switches t_0 and u_n ; while on the generators,

$$T_i^* = T_i, \quad X_i^* = Y_i^{-1}, \quad Y_i^* = X_i^{-1}, \quad i = 1 \dots, n; \quad T_0^* = U_n.$$

We also extend $*$ from \mathbb{F} to an involution on \mathcal{R} by defining $x_i^* = x_i^{-1}$ for all i . Observe that if f is in \mathcal{S} , then f is invariant under $x_i \mapsto x_i^{-1}$, and so f^* is obtained just by switching t_0 and u_n in the coefficients of f .

Next, we define ρ^* by the requirement that $q^{\rho^*} = (q^\rho)^*$. Explicitly,

$$q^{\rho_i^*} = (q^{\rho_i})^* = ((t_0 t_n)^{1/2} t^{n-i})^* = (u_n t_n)^{1/2} t^{n-i}.$$

The duality conjecture of Macdonald can be stated as follows:

7.2. CONJECTURE. *For any two partitions λ and μ we see that*

$$\frac{P_\lambda(q^{\mu+\rho^*})}{P_\lambda(q^{\rho^*})} = \frac{P_\mu^*(q^{\lambda+\rho})}{P_\mu^*(q^\rho)}.$$

This is seen to be equivalent to the formulation in (4.4) of [D], after the easy verification that our definition of duality ($t_0 \leftrightarrow u_n$), is the same as that in (4.1) of [D].

To establish Conjecture 7.2 and its analog for E_α , we introduce the following:

Definition. Let S be the map from \mathcal{H} to \mathbb{F} defined by

$$S(h) := F_h(q^{-\rho^*}); \quad \text{where } F_h = \pi(h)(1) \in \mathcal{R}.$$

7.3. THEOREM. *We have $S(h^*) = S(h)^*$ for all $h \in \mathcal{H}$.*

Proof. By linearity and Corollary 3.4 it is enough to prove this for h of the form $X^\alpha T_w Y^\beta$, with $\alpha, \beta \in \mathbb{Z}^n$, $w \in W_0$. Then by (16) and (22),

$$(24) \quad F_h = q^{\langle \beta, \rho \rangle} \chi(T_w) x^\alpha; \quad \text{and } S(h) = q^{\langle \beta, \rho \rangle} \chi(T_w) q^{-\langle \alpha, \rho^* \rangle}.$$

On the other hand $h^* = X^{-\beta} T_w^* Y^{-\alpha}$, and so

$$(25) \quad F_{h^*} = q^{-\langle \alpha, \rho \rangle} \chi(T_w^*) x^{-\beta}; \quad \text{and } S(h^*) = q^{-\langle \alpha, \rho \rangle} \chi(T_w^*) q^{\langle \beta, \rho^* \rangle}.$$

For $w \in W_0$, $\chi(T_w)$ only involves t and t_n ; and T_w^* is simply obtained from T_w by reversing its product expansion (10) in terms of T_1, \dots, T_n . Thus

$$\chi(T_w)^* = \chi(T_w) = \chi(T_w^*); \quad w \in W_0.$$

Since $*$ interchanges ρ and ρ^* , the result now follows by comparing (24) and (25). \square

We now define scalars $\mathcal{E}_{\alpha\beta}, \mathcal{P}_{\lambda\mu}$ in \mathbb{F} by

$$\mathcal{E}_{\alpha\beta} := E_\alpha^*(q^{\bar{\beta}}) E_\beta(q^{-\rho^*}); \quad \mathcal{P}_{\lambda\mu} := P_\lambda^*(q^{\mu+\rho}) P_\mu(q^{-\rho^*}).$$

7.4. THEOREM. *We have $\mathcal{E}_{\alpha\beta}^* = \mathcal{E}_{\beta\alpha}$ and $\mathcal{P}_{\lambda\mu}^* = \mathcal{P}_{\mu\lambda}$.*

Proof. For the first assertion we consider $h := E_\alpha^*(Y) E_\beta(X)$. Then by the definition of $*$, we get $h^* = E_\beta^*(Y) E_\alpha(X)$. Now,

$$F_h := \pi(E_\alpha^*(Y)) E_\beta(x) = E_\alpha^*(q^{\bar{\beta}}) E_\alpha(x),$$

and so $S(h) = \mathcal{E}_{\alpha\beta}$. Similarly $S(h^*) = \mathcal{E}_{\beta\alpha}$, and the result follows from Theorem 7.3.

The second assertion is proved similarly by considering $h := P_\lambda^*(Y) P_\mu(X)$. \square

7.5. COROLLARY. *Conjecture 7.2 is true.*

Proof. Since P_λ is invariant under $x_i \mapsto x_i^{-1}$ we get

$$\mathcal{P}_{\mu\lambda} := P_\mu^*(q^{\lambda+\rho}) P_\lambda(q^{-\rho^*}) = P_\mu^*(q^{\lambda+\rho}) P_\lambda(q^{\rho^*}).$$

On the other hand,

$$\mathcal{P}_{\lambda\mu}^* := \left(P_\lambda^*(q^{\mu+\rho}) P_\mu(q^{-\rho^*}) \right)^* = P_\lambda(q^{-\mu-\rho^*}) P_\mu^*(q^\rho) = P_\lambda(q^{\mu+\rho^*}) P_\mu^*(q^\rho).$$

Thus the result follows from Theorem 7.4. \square

RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ

E-mail address: sahi@math.rutgers.edu

REFERENCES

- [AW] R. ASKEY and J. WILSON, *Some Basic Hypergeometric Orthogonal Polynomials that Generalize Jacobi Polynomials*, Mem. A.M.S. **54** (1985).
- [C1] I. CHEREDNIK, Double affine Hecke algebras and Macdonald's conjectures, Ann. of Math. **141** (1995), 191–216.
- [C2] ———, Macdonald's evaluation conjectures and difference Fourier transform, Invent. Math. **122** (1995), 119–145.
- [C3] ———, Nonsymmetric Macdonald polynomials, IMRN, no. 10 (1995), 483–515.
- [C4] ———, Intertwining operators of double affine Hecke algebras, Selecta Math. **3** (1997), 459–495.
- [D] J. VAN DIEJEN, Self-dual Koornwinder-Macdonald polynomials, Invent. Math. **126** (1996), 319–339.
- [K] F. KNOP, Integrality of two variable Kostka functions, J. Reine Angew. Math. **482** (1997), 177–189.
- [KS] F. KNOP and S. SAHI, A recursion and a combinatorial formula for Jack polynomials, Invent. Math. **128** (1997), 9–22.
- [Ko] T. KOORNWINDER, Askey-Wilson polynomials for root systems of type BC , Contemp. Math. **138** (1992), 189–204.
- [L] G. LUSZTIG, Affine Hecke algebras and their graded version, J. A.M.S. **2** (1989), 599–635.
- [M] I. MACDONALD, Affine Hecke algebras and orthogonal polynomials, Astérisque, No. 237 (1996), 189–207.
- [N] M. NOUMI, Macdonald-Koornwinder polynomials and affine Hecke rings, Sūrikaiseki-kenkyūsho Kōkyūroku **919** (1995), 44–55 (in Japanese).
- [S] S. SAHI, Interpolation, integrarlity, and a generalization of Macdonald's polynomials, IMRN, no 10 (1996), 457–471.

(Received August 13, 1997)