Problems

- 1. How many 0's does 400! end with?
- 2. Find the smallest positive integer having exactly 100 positive divisors.
- 3. Define $a_1 = 1$, and for every n > 1, define $a_{n+1} = a_n + \frac{1}{a_n}$. Prove that $20 < a_{200} < 24$.
- 4. If S is a set of real numbers let S + S be the set of all sums of the form a + b where $a \in S$ and $b \in S$ (where a, b are allowed to be the same number). For each positive integer n, how should you choose S of size n if you want to minimize the size of S + S?
- 5. Among all lists of positive integers that sum to 1000, what is the most their product can be?
- 6. Find all integers a, b such that $a^b = b^a$.
- 7. Determine all polynomials P(x) that satisfy the equations $P(x^2+1) = (P(x))^2 + 1$ and P(0) = 0.
- 8. Let us say that a positive real number x is a *Fermat number* if there are three distinct positive integers a, b, c such that $a^x + b^x = c^x$. Prove that there exist arbitrarily large Fermat numbers (which means that for every real number M there is a fermat number x that is bigger than M).
- 9. Suppose we define the *linear size* of a box B, denote LS(B) to be the sum of its length, width and height. Is it possible to construct two boxes B, C where LS(B) < LS(C) and C fits inside of B?
- 10. Given 6 points in the plane, show that the ratio of the maximum distance between two of them to the minimum distance between two of them is at least $\sqrt{3}$.
- 11. Suppose we color the points of the x-y plane with three colors. Prove that there must be two points at distance one from each other that get the same color.
- 12. Evaluate $\sqrt{2}^{\sqrt{2}^{\sqrt{2}^{n}}}$. (More formally, let $x_1 = \sqrt{2}$, and let $x_{n+1} = \sqrt{2}^{x_n}$ for each $n \ge 1$. Find $\lim_{n\to\infty} x_n$, if it exists.)
- 13. Standard U.S. coins are pennies (1 cent), nickels (5 cents), dimes (10 cents), quarters (25 cents), half dollars (50 cents) and dollars (100 cents). Say that an integer k is *convertible* provided that it is possible to find a collection of k coins that adds up to a dollar. What is the smallest positive integer that is not convertible?
- 14. Evaluate the determinant of the $n \times n$ matrix whose (i, j)th entry is $a^{|i-j|}$.
- 15. For a nonnegative integer n, let f(n) be the number of ways to express n as a sum of powers of 2, where each power of 2 is used at most twice. For example, f(6) = 3 since 6 = 4 + 2 = 4 + 1 + 1 = 2 + 2 + 1 + 1. (We define f(0) = 1). For a positive integer n. let r(n) = f(n)/f(n-1). The function r is a rather amazing function: it is a bijection from the set of positive integers to the set of the positive rational numbers! Prove this.