

```
[> #HW 20 - Alan Ho
[> #OK to post
```

```
> read("DMB.txt")
```

*First Written: Nov. 2021*

*This is DMB.txt, A Maple package to explore Dynamical models in Biology (both discrete and continuous)*

*accompanying the class Dynamical Models in Biology, Rutgers University. Taught by Dr. Z. (Doron Zeilbeger)*

*The most current version is available on WWW at:  
<http://sites.math.rutgers.edu/~zeilberg/tokhniot/DMB.txt> .  
Please report all bugs to: DoronZeil at gmail dot com .*

*For general help, and a list of the MAIN functions,  
type "Help()". For specific help type "Help(procedure\_name);"*

-----  
*For a list of the supporting functions type: Help1();  
For help with any of them type: Help(ProcedureName);*

-----  
*For a list of the functions that give examples of Discrete-time dynamical systems (some famous),  
type: HelpDDM());*

*For help with any of them type: Help(ProcedureName);*

-----  
*For a list of the functions continuous-time dynamical systems (some famous) type: HelpCDM());  
For help with any of them type: Help(ProcedureName);*

(1)

```
> #1)
```

```
> Help(SIRS)
```

*SIRS(s,i,beta,gamma,nu,N): The SIRS dynamical model with parameters beta,gamma, nu,N (see section 6.6 of Edelstein-Keshet), s is the number of Susceptibles, i is the number of infected, (the number of removed is given by N-s-i). N is the total population. Try:*

(2)

*SIRS(s,i,beta,gamma,nu,N);* (2)

> *Help(EquP)*

*EquP(F,x): Given a transformation F in the list of variables finds all the Equilibrium points of the continuous-time dynamical system  $x'(t)=F(x(t))$*

*EquP([5/2\*x\*(1-x)],[x]);*

*EquP([y\*(1-x-y),x\*(3-2\*x-y)],[x,y]);* (3)

> *Help(SEquP)*

*SEquP(F,x): Given a transformation F in the list of variables finds all the Stable Equilibrium points of the continuous-time dynamical system  $x'(t)=F(x(t))$*

*SEquP([5/2\*x\*(1-x)],[x]);*

*SEquP([y\*(1-x-y),x\*(3-2\*x-y)],[x,y]);* (4)

> *Help(TimeSeries)*

*TimeSeries(F,x,pt,h,A,i): Inputs a transformation F in the list of variables x*

*The time-series of  $x[i]$  vs. time of the Dynamical system approximating the the autonomous continuous dynamical process*

*$dx/dt=F(x(t))$  by a discrete time dynamical system with step-size h from  $t=0$  to  $t=A$*

*Try:*

*TimeSeries([x\*(1-y),y\*(1-x)],[x,y],[0.5,0.5], 0.01, 10,1);* (5)

> *Help(PhaseDiag)*

*PhaseDiag(F,x,pt,h,A): Inputs a transformation F in the list of variables x (of length 2), i.e. a mapping from  $R^2$  to  $R^2$  gives the*

*The phase diagram of the solution with initial condition  $x(0)=pt$*

*$dx/dt=F[1](x(t))$  by a discrete time dynamical system with step-size h from  $t=0$  to  $t=A$*

*Try:*

*PhaseDiag([x\*(1-y),y\*(1-x)],[x,y],[0.5,0.5], 0.01, 10);* (6)

> # i)

>  $F := SIRS\left(s, i, \frac{0.3 \cdot 2}{1000}, 5, 2, 1000\right)$

$F := [-0.0006000000000000 \ s \ i + 5000 - 5 \ s - 5 \ i, 0.0006000000000000 \ s \ i - 2 \ i]$  (7)


> *EquP(F, [s, i])*

*{ [1000., 0.], [3333.333333, -1666.666667] }* (8)


> *SEquP(F, [s, i])*

*{ [1000., 0.] }* (9)


> *TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)*



```
> PhaseDiag(F, [s, i], [800, 200], 0.01, 10)
```



```
> TimeSeries([0.00060000000000 s i - 2 i, -0.00060000000000 s i + 5000 - 5 s - 5 i], [i, s], [200, 800], 0.01, 10, 1)
```




>  $F := SIRS\left(s, i, \frac{0.9 \cdot 2}{1000}, 5, 2, 1000\right)$   
 $F := [-0.001800000000 \ s \ i + 5000 - 5 \ s - 5 \ i, 0.001800000000 \ s \ i - 2 \ i]$  (10)


>  $EquP(F, [s, i])$   
 $\{[1000., 0.], [1111.111111, -79.36507937]\}$  (11)

>  $SEquP(F, [s, i])$   
 $\{[1000., 0.]\}$  (12)


>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$



>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$



> `TimeSeries([0.001800000000 s i - 2 i, -0.001800000000 s i + 5000 - 5 s - 5 i], [i, s], [200, 800], 0.01, 10, 1)`




>  $F := \text{SIRS}\left(s, i, \frac{3.9 \cdot 2}{1000}, 5, 2, 1000\right)$   
 $F := [-0.007800000000 s i + 5000 - 5 s - 5 i, 0.007800000000 s i - 2 i]$  (13)


> `EquP(F, [s, i])`  
 $\{[256.4102564, 531.1355311], [1000., 0.]\}$  (14)

> `SEquP(F, [s, i])`  
 $\{[256.4102564, 531.1355311]\}$  (15)


> `TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)`



> *PhaseDiag*(*F*, [*s*, *i*], [800, 200], 0.01, 10)



> *TimeSeries*([0.007800000000 *s i* - 2 *i*, -0.007800000000 *s i* + 5000 - 5 *s* - 5 *i*], [*i*, *s*], [200, 800], 0.01, 10, 1)



> #ii)


>

>  $F := SIRS\left(s, i, \frac{0.3 \cdot 3}{1000}, 6, 3, 1000\right)$   
 $F := [-0.0009000000000000 s i + 6000 - 6 s - 6 i, 0.0009000000000000 s i - 3 i]$  (16)


>  $EquP(F, [s, i])$   
 $\{[1000., 0.], [3333.333333, -1555.555556]\}$  (17)

>  $SEquP(F, [s, i])$   
 $\{[1000., 0.]\}$  (18)


>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$



>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$



> *TimeSeries*( [0.0009000000000  $s i - 3 i$ , -0.0009000000000  $s i + 6000 - 6 s - 6 i$ ], [ $i, s$ ],  
 [200, 800], 0.01, 10, 1)




>  $F := \text{SIRS}\left(s, i, \frac{0.9 \cdot 3}{1000}, 6, 3, 1000\right)$   
 $F := [-0.002700000000 s i + 6000 - 6 s - 6 i, 0.002700000000 s i - 3 i]$  (19)

> *EquP*( $F, [s, i]$ )  
 {[1000., 0.], [1111.111111, -74.07407407]} (20)


> *SEquP*( $F, [s, i]$ )  
 {[1000., 0.]} (21)

> *TimeSeries*( $F, [s, i]$ , [800, 200], 0.01, 10, 1)






> *PhaseDiag*(*F*, [*s*, *i*], [800, 200], 0.01, 10)



> *TimeSeries*([0.002700000000 *s i* - 3 *i*, -0.002700000000 *s i* + 6000 - 6 *s* - 6 *i*], [*i*, *s*], [200, 800], 0.01, 10, 1)




>  $F := SIRS\left(s, i, \frac{3.9 \cdot 3}{1000}, 6, 3, 1000\right)$   
 $F := [-0.01170000000 \ s \ i + 6000 - 6 \ s - 6 \ i, 0.01170000000 \ s \ i - 3 \ i]$  (22)


>  $EquP(F, [s, i])$   
 $\{[256.4102564, 495.7264957], [1000., 0.]\}$  (23)

>  $SEquP(F, [s, i])$   
 $\{[256.4102564, 495.7264957]\}$  (24)


>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$



>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$



```
> TimeSeries([0.01170000000 s i - 3 i, -0.01170000000 s i + 6000 - 6 s - 6 i], [i, s], [200, 800], 0.01, 10, 1)
```




```
> #iii)
```

```
> F := SIRS(s, i, 0.3·4 / 1000, 1, 4, 1000)
      F := [-0.001200000000 s i + 1000 - s - i, 0.001200000000 s i - 4 i] (25)
```


```
> EquP(F, [s, i])
      { [1000., 0.], [3333.333333, -466.6666667] } (26)
```

```
> SEquP(F, [s, i])
      { [1000., 0.] } (27)
```


> *TimeSeries*( $F$ , [ $s$ ,  $i$ ], [800, 200], 0.01, 10, 1)



> *PhaseDiag*( $F$ , [ $s$ ,  $i$ ], [800, 200], 0.01, 10)




> *TimeSeries*( $[0.001200000000 s i - 4 i, -0.001200000000 s i + 1000 - s - i]$ , [ $i$ ,  $s$ ], [200, 800], 0.01, 10, 1)




>  $F := SIRS\left(s, i, \frac{0.9 \cdot 4}{1000}, 1, 4, 1000\right)$   
 $F := [-0.003600000000 \, s i + 1000 - s - i, 0.003600000000 \, s i - 4 \, i]$  (28)

>  $EquP(F, [s, i])$   
 $\{[1000., 0.], [1111.111111, -22.22222222]\}$  (29)


>  $SEquP(F, [s, i])$   
 $\{[1000., 0.]\}$  (30)



>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$



```
> TimeSeries([0.003600000000 s i - 4 i, -0.003600000000 s i + 1000 - s - i], [i, s], [200, 800], 0.01, 10, 1)
```




```
> F := SIRS(s, i, 3.9*4/1000, 1, 4, 1000)
      F := [-0.015600000000 s i + 1000 - s - i, 0.015600000000 s i - 4 i] (31)
```


```
> EquP(F, [s, i])
      {[256.4102564, 148.7179487], [1000., 0.]} (32)
```

```
> SEquP(F, [s, i])
      {[256.4102564, 148.7179487]} (33)
```


```
> TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)
```



> *PhaseDiag*( $F$ , [ $s$ ,  $i$ ], [800, 200], 0.01, 10)



> *TimeSeries*([ $0.01560000000 s i - 4 i$ ,  $-0.01560000000 s i + 1000 - s - i$ ], [ $i$ ,  $s$ ], [200, 800], 0.01, 10, 1)




> #iv)

>  $F := SIRS\left(s, i, \frac{0.3 \cdot 7}{1000}, 10, 7, 1000\right)$   
 $F := [-0.002100000000 \, s \, i + 10000 - 10 \, s - 10 \, i, 0.002100000000 \, s \, i - 7 \, i]$  (34)

>  $EquP(F, [s, i])$   
 $\{[1000., 0.], [3333.333333, -1372.549020]\}$  (35)


>  $SEquP(F, [s, i])$   
 $\{[1000., 0.]\}$  (36)

>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$




>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$





>  $TimeSeries([0.002100000000 s i - 7 i, -0.002100000000 s i + 10000 - 10 s - 10 i], [i, s], [200, 800], 0.01, 10, 1)$




>  $F := SIRS\left(s, i, \frac{0.9 \cdot 7}{1000}, 10, 7, 1000\right)$   
 $F := [-0.006300000000 s i + 10000 - 10 s - 10 i, 0.006300000000 s i - 7 i]$  (37)


>  $EquP(F, [s, i])$   
 $\{[1000., 0.], [1111.111111, -65.35947712]\}$  (38)

>  $SEquP(F, [s, i])$   
 $\{[1000., 0.]\}$  (39)


>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$



> *PhaseDiag*(*F*, [*s*, *i*], [800, 200], 0.01, 10)



> *TimeSeries*([0.006300000000 *s i* - 7 *i*, -0.006300000000 *s i* + 10000 - 10 *s* - 10 *i*], [*i*, *s*], [200, 800], 0.01, 10, 1)




>  $F := SIRS\left(s, i, \frac{3.9 \cdot 7}{1000}, 10, 7, 1000\right)$   
 $F := [-0.02730000000 \, s \, i + 10000 - 10 \, s - 10 \, i, 0.02730000000 \, s \, i - 7 \, i]$  (40)


>  $EquP(F, [s, i])$   
 $\{[256.4102564, 437.4057315], [1000., 0.]\}$  (41)

>  $SEquP(F, [s, i])$   
 $\{[256.4102564, 437.4057315]\}$  (42)


>  $TimeSeries(F, [s, i], [800, 200], 0.01, 10, 1)$



>  $PhaseDiag(F, [s, i], [800, 200], 0.01, 10)$



```
> TimeSeries([0.02730000000 s i - 7 i, -0.02730000000 s i + 10000 - 10 s - 10 i], [i, s], [200, 800], 0.01, 10, 1)
```



```
> #2)
```

```
> F := RandNice([x, y], 3)
```

$$F := [(2 - 2x - 3y)(2 - x - 3y), (1 - x - 2y)(3 - 2x - 2y)] \quad (43)$$

```
> EquP(F, [x, y])
```


$$\left\{ [-1, 1], [1, 0], \left[ \frac{5}{2}, -1 \right], \left[ \frac{5}{4}, \frac{1}{4} \right] \right\} \quad (44)$$

```
> SEquP(F, [x, y])
```


{[1., 0.]}

(45)


> *TimeSeries*(*F*, [*x*, *y*], [1.1, 0.1], 0.01, 10, 1)



> *TimeSeries*([*F*[2], *F*[1]], [*y*, *x*], [0.1, 1.1], 0.01, 10, 1)



> *PhaseDiag*(*F*, [*x*, *y*], [1.1, 0.1], 0.01, 10)



```
> F := RandNice([x, y], 3)
      F := [(1 - 2x - y) (1 - 2x - 2y), (3 - 3x - 2y) (2 - x - y)]
```

(46)


```
> EquP(F, [x, y])
      {[-1, 3], [2, -3/2]}
```

(47)


```
> SEquP(F, [x, y])
      {[2., -1.500000000]}
```

(48)


```
> TimeSeries(F, [x, y], [2.1, -1.4], 0.01, 10, 1)
```



```
> TimeSeries([F[2], F[1]], [y, x], [-1.4, 2.1], 0.01, 10, 1)
```



> *PhaseDiag*(*F*, [*x*, *y*], [2.1, -1.4], 0.01, 10)



> *F* := *RandNice*([*x*, *y*], 3)

$$F := [(3 - x - 3y)(3 - 2x - 3y), (1 - x - 3y)(1 - x - 2y)] \quad (49)$$


> *EquP*(*F*, [*x*, *y*])

$$\left\{ [-3, 2], \left[ 2, -\frac{1}{3} \right], [3, -1] \right\} \quad (50)$$


> *SEquP*(*F*, [*x*, *y*])

$$\{ [3., -1.] \} \quad (51)$$

> *TimeSeries*(*F*, [*x*, *y*], [3.1, -0.9], 0.01, 10, 1)




> *TimeSeries*([F[2], F[1]], [y, x], [-0.9, 3.1], 0.01, 10, 1)



> *PhaseDiag*(F, [x, y], [3.1, -0.9], 0.01, 10)





>

> #3)

> Help(Orbk)

*Orbk(k,z,f,INI,K1,K2): Given a positive integer k, a letter (symbol), z, an expression f of  $z[1], \dots, z[k]$  (representing a multi-variable function of the variables  $z[1], \dots, z[k]$  a vector INI representing the initial values  $[x[1], \dots, x[k]]$ , and (in applications) positive integers K1 and K2, outputs the values of the sequence starting at  $n=K1$  and ending at  $n=K2$ . of the sequence satisfying the difference equation*

$$x[n]=f(x[n-1],x[n-2],\dots, x[n-k+1]):$$

*This is a generalization to higher-order difference equation of procedure Orb(f,x,x0,K1,K2).*

*For example, try:*

*Orbk(1,z,5/2\*z[1]\*(1-z[1]),[0.5],1000,1010);*

*To get the Fibonacci sequence, type:*

*Orbk(2,z,z[1]+z[2],[1,1],1000,1010);*

*To get the part of the orbit between  $n=1000$  and  $n=1010$ , of the 3rd order recurrence given in Eq. (4) of the Ladas-Amleh paper*

*<https://sites.math.rutgers.edu/~zeilberg/Bio21/AmlehLadas.pdf>*

*with initial conditions  $x(0)=1, x(1)=3, x(2)=5$ , Type:*

*Orbk(3,z,z[2]/(z[2]+z[3]),[1.,3.,5.],1000,1010);*

*To get the part of the orbit between  $n=1000$  and  $n=1010$ , of the 3rd order recurrence given in*

Eq. (5) of the Ladas-Amleh paper

with initial conditions  $x(0)=1, x(1)=3, x(2)=5$ , Type:  
 $Orbk(3,z,(z[1]+z[3])/z[2],[1.,3.,5.],1000,1010);$

To get the part of the orbit between  $n=1000$  and  $n=1010$ , of the 3rd order recurrence given in Eq. (6) of the Ladas-Amleh paper

with initial conditions  $x(0)=1, x(1)=3, x(2)=5$ , Type:  
 $Orbk(3,z,(1+z[3])/z[1],[1.,3.,5.],1000,1010);$

To get the part of the orbit between  $n=1000$  and  $n=1010$ , of the 3rd order recurrence given in Eq. (7) of the Ladas-Amleh paper

with initial conditions  $x(0)=1, x(1)=3, x(2)=5$ , Type:  
 $Orbk(3,z,(1+z[1])/(z[2]+z[3]),[1.,3.,5.],1000,1010);$  (52)

>  $Orbk\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, [0.5, 0.5, 0.5, 0.5], 2000, 2010\right)$   
 $[1.342779698, 2.576461980, 1.342779698, 2.576461980, 1.342779698, 2.576461980,$  (53)  
 $1.342779698, 2.576461980, 1.342779698, 2.576461980, 1.342779698]$

>  $Orbk\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, [0.75, 0.75, 0.75, 0.75], 1000, 1010\right)$   
 $[1.464493936, 2.314423019, 1.464493936, 2.314423019, 1.464493936, 2.314423019,$  (54)  
 $1.464493936, 2.314423019, 1.464493936, 2.314423019, 1.464493936]$

>  $Orbk\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, [1.5, 1.5, 1.5, 1.5], 1000, 1010\right)$   
 $[1.734194427, 1.917928943, 1.734194427, 1.917928943, 1.734194427, 1.917928943,$  (55)  
 $1.734194427, 1.917928943, 1.734194427, 1.917928943, 1.734194427]$

>  $Orbk\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, [2.5, 2.5, 2.5, 2.5], 2000, 2010\right)$   
 $[1.978191241, 1.683879292, 1.978191241, 1.683879292, 1.978191241, 1.683879292,$  (56)  
 $1.978191241, 1.683879292, 1.978191241, 1.683879292, 1.978191241]$

>  $Orbk\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, [5.5, 5.5, 5.5, 5.5], 2000, 2010\right)$   
 $[2.456781392, 1.394325757, 2.456781392, 1.394325757, 2.456781392, 1.394325757,$  (57)  
 $2.456781392, 1.394325757, 2.456781392, 1.394325757, 2.456781392]$

>  $ToSys\left(4, z, \frac{3+z[2]+z[3]+z[4]}{1+z[1]+z[3]}, \left[\frac{3+z_2+z_3+z_4}{1+z_1+z_3}, z_1, z_2, z_3\right], [z_1, z_2, z_3, z_4]\right)$  (58)

>  $SFP(\%)$   
 $\{[1.822875656, 1.822875656, 1.822875656, 1.822875656]\}$  (59)

