

Dr. Z's Math151 Handout #2.1 [The Tangent and Velocity Problems]

By Doron Zeilberger

Problem Type 2.1.1 : The point $P(a, f(a))$ lies on the curve $y = f(x)$. (a) If Q is the point $(x, f(x))$, use your calculator to find the slope of the secant line PQ (correct to six decimal places) for the following values of x : (i) x_1 (ii) $a + \text{tiny}$ (iii) $a + \text{'very tiny'}$

(b) Using the results of part (a), guess the value of the slope of the tangent line to the curve at $P(a, f(a))$.

(c) Using the slope from part (b), find the equation of the tangent line to the curve at $P = (a, f(a))$.

Example Problem 2.1.1: The point $P(2, 8)$ lies on the curve $y = x^3$. (a) If Q is the point (x, x^3) , use your calculator to find the slope of the secant line PQ (correct to six decimal places) for the following values of x : (i) 3 (ii) 2.001 (iii) 2.0001.

(b) Using the results of part (a), guess the value of the slope of the tangent line to the curve at $P(2, 8)$.

(c) Using the slope from part (b), find the equation of the tangent line to the curve at $P = (2, 8)$.

Steps

1. (a) The slope of the secant line between $P(a, f(a))$ and $Q(x, f(x))$ is

$$(f(x) - f(a))/(x - a) \quad .$$

2. (b) The answers to (ii) and (iii) should be very close to each other and if they are both close to a 'nice' value, that would be a good guess.

Example

1. (a) (i) $(3^3 - 2^3)/(3 - 2) = (27 - 8)/1 = 19$;

(ii) $((2.001)^3 - 2^3)/(2.001 - 2) = (8.012006 - 8)/.001 = 12.006001$;

(iii) $((2.0001)^3 - 2^3)/(2.0001 - 2) = (8.00120006 - 8)/.0001 = 12.00060$;

2. (b) guessed slope=12.

3. (c) $(y - f(a)) = (\text{slope})(x - a)$

3. (c) $(y - 8) = (12)(x - 2)$, hence,

$$y = 12x - 24 + 8 = 12x - 16.$$

Answer: $y = 12x - 16$.

Problem Type 2.1.2 : (a) In a certain planet the height of a stone thrown vertically upwards with velocity v_0 m/s is given by $h = v_0 t - At^2$. Find the average velocity in the time intervals (i) $[a, b]$ (ii) $[a, a + \text{tiny}]$ (iii) $[a, a + \text{very tiny}]$.

Example Problem 2.1.2: (a) In a certain planet the height of a stone thrown vertically upwards with velocity 100 m/s is given by $h = 100t - t^2$. Find the average velocity in the time intervals

(i) $[1, 2]$ (ii) $[1, 1.01]$ (iii) $[1, 1.001]$.

(b) Estimate the instantaneous velocity after one second.

Steps

1. (a) The average velocity of a particle whose height (or distance) is given by $h(t)$, over a time interval $[a, b]$ is

$$\frac{h(b) - h(a)}{b - a} \quad .$$

2. (b) The answers to (ii) and (iii) should be very close to each other, and if they are both close to a ‘nice’ value, that would be a good estimate.

Example

1. (a) $h(t) = 100t - t^2$ so

$$(i) \quad (h(2) - h(1)) / (2 - 1) = ((100(2) - 2^2) - (100(1) - 1^2)) / 1 = (196 - 99) / 1 = 97.$$

$$(ii) \quad (h(1.01) - h(1)) / (1.01 - 1) = ((100(1.01) - (1.01)^2) - (100(1) - 1^2)) / .01 = 97.99.$$

$$(iii) \quad (h(1.001) - h(1)) / (1.001 - 1) = ((100(1.001) - (1.001)^2) - (100(1) - 1^2)) / .001 = 97.999.$$

2. (b) 98.