

Dr. Z's Math251 Handout #15.4b [Integrations in Cylindrical and Spherical Coordinates]

By Doron Zeilberger

Problem Type 15.4ba: Evaluate

$$\int \int \int_E F(x, y, z) dV \quad ,$$

where E is a solid region described in terms of cylinders and other stuff.

Example Problem 15.4ba: Evaluate

$$\int \int \int_E x^2 dV \quad ,$$

where E is the solid that lies within the cylinder $x^2 + y^2 = 1$, above the plane $z = 0$, and below the cone $z^2 = 4x^2 + 4y^2$.

Steps

1. Chances are that you are supposed to use *cylindrical coordinates*. Express E in the form

$$E = \{ (r, \theta, z) \mid \alpha \leq \theta \leq \beta,$$

$$h_1(\theta) \leq r \leq h_2(\theta), u_1(r, \theta) \leq z \leq u_2(r, \theta) \} .$$

Example

1. Since $x^2 + y^2 = r^2$, the cone $z^2 = 4x^2 + 4y^2$ can be written $z^2 = 4r^2$. The cylinder $x^2 + y^2 = 1$ is really $r = 1$, and this is the “base”. $z^2 = 4r^2$ means that z ranges between $-2r$ and $2r$. **but** we are also told that our solid is **above** the plane $z = 0$, so z ranges between 0 and $2r$. It turns out that

$$E = \{ (r, \theta, z) \mid 0 \leq \theta \leq 2\pi, 0 \leq r \leq 1, 0 \leq z \leq 2r \} .$$

2. Express the integrand $F(x, y, z)$ in cylindrical coordinates, using the “dictionary” **2.**

$$x = r \cos \theta \quad , \quad y = r \sin \theta \quad , \quad x^2 + y^2 = r^2 \quad . \quad \int_0^{2\pi} \int_0^1 \int_0^{2r} (r \cos \theta)^2 r \, dz \, dr \, d\theta \quad .$$

Also $dV = r \, dr \, d\theta \, dz$. Then set up the volume integral as an iterated integral

$$\int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} \int_{u_1(r, \theta)}^{u_2(r, \theta)} F(r \cos \theta, r \sin \theta, z) r \, dz \, dr \, d\theta \quad .$$

3. Evaluate the integral from the inside to the outside.

3. The inside integral is:

$$\int_0^{2r} (r \cos \theta)^2 r \, dz = r^3 \cos^2 \theta \int_0^{2r} dz = 2r^4 \cos^2 \theta \quad .$$

The middle integral is

$$\begin{aligned} & \int_0^1 \left[\int_{-2r}^{2r} (r \cos \theta)^2 r \, dz \right] dr \\ &= \int_0^1 2r^4 \cos^2 \theta \, dr = \cos^2 \theta \int_0^1 2r^4 \, dr = \frac{2}{5} \cos^2 \theta \quad . \end{aligned}$$

The outer integral is

$$\begin{aligned} & \int_0^{2\pi} \left[\int_0^1 \int_{-2r}^{2r} (r \cos \theta)^2 r \, dz \, dr \right] d\theta \\ &= \int_0^{2\pi} \frac{2}{5} \cos^2 \theta \, d\theta \\ &= \frac{2}{5} \int_0^{2\pi} \frac{1 + \cos 2\theta}{2} \, d\theta \\ &= \frac{2}{5} \cdot \left[\frac{\theta + (1/2) \sin 2\theta}{2} \right]_0^{2\pi} = \frac{2\pi}{5} \quad . \end{aligned}$$

Ans.: $\frac{2\pi}{5}$.

Problem Type 15.4bb: Evaluate

$$\int \int \int_E F(x, y, z) \, dV \quad ,$$

where E is bounded by the xz -plane and the hemispheres $y = \sqrt{r_1^2 - x^2 - z^2}$ and $y = \sqrt{r_2^2 - x^2 - z^2}$.

Example Problem 15.4bb: Evaluate

$$\int \int \int_E x^2 dV \quad ,$$

where E is bounded by the xz -plane and the hemispheres $y = \sqrt{1 - x^2 - z^2}$ and $y = \sqrt{4 - x^2 - z^2}$.

Steps

1. This is best handled with spherical coordinates. The hemisphere $y = \sqrt{R^2 - x^2 - z^2}$ is half of the sphere $x^2 + y^2 + z^2 = R^2$ whose equation in spherical coordinates is really simple: $\rho = R$. Since $y > 0$ (the square-root is always positive), the range of θ is between 0 and π . ρ is between r_1 and r_2 and ϕ has its full range 0 to π . So

$$E = \{ (\rho, \theta, \phi) \mid r_1 \leq \rho \leq r_2, 0 \leq \theta \leq \pi, 0 \leq \phi \leq \pi \} \quad .$$

2. Using the ‘dictionary’

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta,$$

$$z = \rho \cos \phi \quad ; \quad dV = \rho^2 \sin \phi d\rho d\theta d\phi \quad .$$

Convert the volume integral into an iterated spherical integral (using the description of E in step 1).

$$\int_0^\pi \int_0^\pi \int_{r_1}^{r_2} F(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi d\rho d\theta d\phi \quad .$$

Example

1. Here the radii are 1 and 2 so

$$E = \{ (\rho, \theta, \phi) \mid 1 \leq \rho \leq 2, 0 \leq \theta \leq \pi, 0 \leq \phi \leq \pi \} \quad .$$

2.

$$\int_0^\pi \int_0^\pi \int_1^2 (\rho \sin \phi \cos \theta)^2 \rho^2 \sin \phi d\rho d\theta d\phi$$

$$= \int_0^\pi \int_0^\pi \int_1^2 \rho^4 \sin^3 \phi \cos^2 \theta d\rho d\theta d\phi \quad .$$

3. If you are lucky and all the limits of integration are numbers (do not involve ρ , ϕ or θ) and the integrand is a product of functions of a single variable, then the iterated integral is simply a product of three simple integrals. Express the big integral like that, and evaluate each single integral separately. Then multiply them together.

Warning: This is only possible if *all* the limits of integration are numbers and the integrand is *completely* separable as a product of functions of a single variable.

3.

$$\begin{aligned} & \int_0^\pi \int_0^\pi \int_1^2 \rho^4 \sin^3 \phi \cos^2 \theta \, d\rho \, d\theta \, d\phi \\ &= \int_0^\pi \sin^3 \phi \, d\phi \int_0^\pi \cos^2 \theta \, d\theta \int_1^2 \rho^4 \, d\rho \quad . \end{aligned}$$

The first integral is ($u = \cos \phi$)

$$\begin{aligned} \int_0^\pi \sin^3 \phi \, d\phi &= \int_0^\pi \sin^2 \phi \, d(-\cos \phi) = \\ \int_0^\pi (1 - \cos^2 \phi) \, d(-\cos \phi) &= - \int_1^{-1} (1 - u^2) \, du = \\ &= -u + \frac{u^3}{3} \Big|_1^{-1} = \frac{4}{3} \quad . \end{aligned}$$

The second integral is

$$\begin{aligned} \int_0^\pi \cos^2 \theta \, d\theta &= \int_0^\pi \frac{1 + \cos 2\theta}{2} \, d\theta = \\ &= \frac{\theta}{2} + \frac{\sin 2\theta}{4} \Big|_0^\pi = \frac{\pi}{2} \quad . \end{aligned}$$

The third integral is

$$\int_1^2 \rho^4 \, d\rho = \frac{\rho^5}{5} \Big|_1^2 = \frac{2^5 - 1^5}{5} = \frac{31}{5} \quad .$$

Multiplying these three single-integrals, we get that the original triple integral equals

$$\frac{4}{3} \cdot \frac{\pi}{2} \cdot \frac{31}{5} = \frac{62\pi}{15} \quad .$$

Ans.: $\frac{62\pi}{15}$.

A Problem from a Previous Final: Evaluate

$$\iiint_B 5(x^2 + y^2 + z^2)^2 \, dV$$

where B is the ball

$$\{(x, y, z) \mid x^2 + y^2 + z^2 \leq 4\} \quad .$$

Ans.: $2560\pi/7$.