

Solutions to the “QUIZ” for Lecture 16

1. Compute the Jacobian of the transformation

$$\Phi(r, s) = (rs, r + s)$$

Sol.: Here $x = rs, y = r + s$ and

$$J = (x_r)(y_s) - (x_s)(y_r) = (s)(1) - (r)(1) = s - r \quad .$$

Ans. $s - r$ (type: function of r and s).

2. Let $\mathcal{D} = \Phi(\mathcal{R})$ where $\Phi(u, v) = (u + v, v^2)$ and $R = [0, 6] \times [1, 2]$. Calculate

$$\int \int_{\mathcal{D}} y \, dA \quad .$$

(Note: it is not necessary to compute D).

Sol. Here the transformation is $x = u + v, y = v^2$. The Jacobian is $J = (x_u)(y_v) - (x_v)(y_u) = (1)(2v) - (1)(0) = 2v$.

By the change of variable formula we have

$$\int \int_{\mathcal{D}} y \, dA = \int \int_{\mathcal{R}} yJ \, dA = \int \int_{\mathcal{R}} (v^2)(2v) \, dA = \int \int_{\mathcal{R}} 2v^3 \, dA$$

R is the rectangle $[0, 6] \times [1, 2]$, which means:

$$\{(u, v) \mid 0 \leq u \leq 6, 1 \leq v \leq 2\}$$

That is both type I and type II. Using the type-I formulation we have

$$\begin{aligned} \int_0^6 \int_1^2 2v^3 \, dv \, du &= \left(\int_0^6 du \right) \left(\int_1^2 2v^3 \, dv \right) = \\ \left(u \Big|_0^6 \right) \left(\frac{v^4}{2} \Big|_1^2 \right) &= (6 - 0) \cdot \left(\frac{2^4 - 1^4}{2} \right) = 6 \cdot \left(\frac{15}{2} \right) = 45 \quad . \end{aligned}$$

Ans. 45 (type number).