



## Automated discovery and proof of congruence theorems for partial sums of combinatorial sequences

William Y. C. Chen, Qing-Hu Hou & Doron Zeilberger

To cite this article: William Y. C. Chen, Qing-Hu Hou & Doron Zeilberger (2016): Automated discovery and proof of congruence theorems for partial sums of combinatorial sequences, Journal of Difference Equations and Applications, DOI: [10.1080/10236198.2016.1142541](https://doi.org/10.1080/10236198.2016.1142541)

To link to this article: <http://dx.doi.org/10.1080/10236198.2016.1142541>



Published online: 03 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 2



View related articles [↗](#)



View Crossmark data [↗](#)

# Automated discovery and proof of congruence theorems for partial sums of combinatorial sequences

William Y. C. Chen<sup>a</sup>, Qing-Hu Hou<sup>a</sup> and Doron Zeilberger<sup>b</sup>

<sup>a</sup>Center for Applied Mathematics, Tianjin University, Tianjin, P.R. China; <sup>b</sup>Department of Mathematics, Rutgers University (New Brunswick), Piscataway, NJ, USA

## ABSTRACT

Many combinatorial sequences (e.g. the Catalan and the Motzkin numbers) may be expressed as the constant term of  $P(x)^k Q(x)$ , for some Laurent polynomials  $P(x)$  and  $Q(x)$  in the variable  $x$  with integer coefficients. Denoting such a sequence by  $a_k$ , we obtain a general formula that determines the congruence class, modulo  $p$ , of the indefinite sum  $\sum_{k=0}^{rp-1} a_k$ , for any prime  $p$ , and any positive integer  $r$ , as a linear combination of sequences that satisfy linear recurrence (alias difference) equations with constant coefficients. This enables us (or rather, our computers) to automatically discover and prove congruence theorems for such partial sums. Moreover, we show that in many cases, the set of the residues is finite, regardless of the prime  $p$ .

## ARTICLE HISTORY

Received 8 January 2016  
Accepted 10 January 2016

## KEYWORDS

Polynomials; combinatorial sequences; Legendre symbol; Laurent series

## AMS SUBJECT CLASSIFICATIONS

12Dxx

## 1. Introduction

Let  $\{a_k\}$  be a sequence of integers, and  $r$  be a positive integer. We focus on the congruences of the partial sum  $\sum_{k=0}^{rp-1} a_k$  modulo a general prime  $p$ . When  $a_k$  is a hypergeometric term and  $r = 1$  we get a truncated hypergeometric series, which is closely related to the Gaussian hypergeometric series introduced by Greene [3]. An interesting example is the congruence for the Apéry numbers [1,2]

$$A\left(\frac{p-1}{2}\right) \equiv \sum_{k=0}^{p-1} \binom{2k}{k}^4 2^{-8k} \equiv \gamma(p) \pmod{p},$$

where

$$A(n) = \sum_{j=0}^n \binom{n+j}{j}^2 \binom{n}{j}^2,$$

and

$$\sum_{n=1}^{\infty} \gamma(n) q^n = q \prod_{n=1}^{\infty} (1 - q^{2n})^4 (1 - q^{4n})^4.$$

These congruences are usually obtained case by case and the proofs are complicated. For example, Pan and Sun [5] used a sophisticated combinatorial identity to deduce that

$$\sum_{k=0}^{p-1} \binom{2k}{k+d} \equiv \left(\frac{p-d}{3}\right) \pmod{p}, \quad d = 0, 1, \dots, p,$$

where  $\left(\cdot\right)$  is the Legendre symbol. We propose an automated method to discover and prove such congruences for a large family of combinatorial sequences  $\{a_k\}$ . More precisely, we assume that  $a_k$  is the constant term of  $P(x)^k Q(x)$  where  $P(x)$  and  $Q(x)$  are two Laurent polynomials in the (single) variable  $x$  with integer coefficients. Rowland and Zeilberger [6] discovered an algorithm to automatically generate *automata* for determining the congruences, modulo a prime  $p$ , of combinatorial sequences (not the partial sums) but for *specific* primes  $p$  (one at a time).

Throughout the paper,  $p$  always denotes a prime number. We write  $a \equiv_p b$  if  $a$  is congruent to  $b$  modulo  $p$ . For a Laurent series  $f(x) = \sum_{k \geq k_0} a_k x^k$ , we use  $CT f(x)$  to denote the coefficient of the free term,  $x^0$ . The set of integers, rational numbers and complex numbers are denoted by  $\mathbb{Z}, \mathbb{Q}$  and  $\mathbb{C}$ , respectively. The finite field with  $p$  elements is denoted by  $\mathbb{F}_p$ .

## 2. Evaluation

In this section, we show that the above-mentioned partial sums are linear combinations of  $C$ -finite sequences, i.e. integer sequences that satisfy a linear recurrence equation with *constant* coefficients (like  $2^n$  and the Fibonacci numbers, to name two examples). This would enable us (and our computers) to discover and prove practically infinitely many theorems about the congruences of such partial sums modulo an *arbitrary* (symbolic!) prime  $p$ .

We have the following formula for the congruences of the partial sums.

**Theorem 2.1:** *Let  $P(x), Q(x)$  be two Laurent polynomials in  $x$  with integer coefficients and*

$$a_k := CT P(x)^k Q(x).$$

*Let  $-m$  and  $-n$  be the lowest degrees of  $P(x)$  and  $Q(x)/(P(x) - 1)$ , respectively. Then for any positive integer  $r$ , and any prime  $p > n$ , we have*

$$\sum_{k=0}^{rp-1} a_k \equiv_p \sum_{j=0}^{rm} c_j S_{(rm-j)p}, \tag{2.1}$$

*where  $c_j$  is the coefficient of  $x^{-rm+j}$  in  $P(x)^r - 1$  and  $S_k$  is the coefficient of  $x^k$  in the Laurent expansion of*

$$\frac{Q(x)}{P(x) - 1}.$$

**Proof:** Noting that CT is a linear operator, we have

$$\sum_{k=0}^{rp-1} a_k = \text{CT} \sum_{k=0}^{rp-1} P(x)^k Q(x) = \text{CT} \frac{(P(x)^{rp} - 1)Q(x)}{P(x) - 1}.$$

Since the coefficients of  $P(x)$  are integers, we have, (by the “Freshman’s Dream Identity” ,  $(a + b)^p \equiv_p a^p + b^p$ ),  $P(x)^p \equiv_p P(x^p)$  and hence

$$\sum_{k=0}^{rp-1} a_k \equiv_p \text{CT} \frac{Q(x)(P(x^p)^r - 1)}{P(x) - 1}.$$

By the definition of  $m$  and  $c_j$ , we see that

$$P(x)^r - 1 = \sum_{j=0}^N c_j x^{-rm+j},$$

for some integer  $N$ . If  $j > rm$ , we have

$$(-rm + j)p - n > n(-rm + j) - n \geq 0,$$

which implies that

$$\text{CT} \frac{Q(x)x^{(-rm+j)p}}{P(x) - 1} = 0.$$

Hence

$$\text{CT} \frac{Q(x)(P(x^p)^r - 1)}{P(x) - 1} = \sum_{j=0}^{rm} c_j \text{CT} \frac{Q(x)x^{(-rm+j)p}}{P(x) - 1} = \sum_{j=0}^{rm} c_j S_{(rm-j)p}.$$

This completes the proof. □

This theorem is implemented in the Maple package `CTcong.txt` available from the webpage <http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/ctcong.html> where the user can also find sample input and output files.

The Maple command-line is

$$\text{TheoG}(P, Q, x, p, C, r),$$

where  $P, Q$  are two Laurent polynomials, with integer coefficients, in the variable  $x$ ,  $p$  is the symbol standing for the prime,  $C$  is the name for the sequence of coefficients of  $Q(x)/(P(x) - 1)$ , while  $r$  is as in Equation (2.1). For example, typing (in a Maple session, after reading our Maple package `CTcong.txt`)

$$\text{TheoG}(1/x+2+x, x^d, x, p, C, 1);$$

immediately outputs

**Corollary 2.2:** Let  $A(i)$  be the constant term of the Laurent polynomial

$$x^d \left( \frac{1}{x} + 2 + x \right)^i,$$

and for any prime  $p$ , let

$$B(p) = \sum_{i=0}^{p-1} A(i).$$

Then

$$B(p) \equiv_p C(p),$$

where  $C(n)$  is the  $C$ -finite sequence defined in terms of the generating function

$$\sum_{i=0}^{\infty} C(i)x^i = \frac{x^{d+1}}{x^2 + x + 1}.$$

Noting that

$$\text{CT } x^d \left( \frac{1}{x} + 2 + x \right)^i = \binom{2i}{i-d} = \binom{2i}{i+d},$$

and

$$\frac{x^{d+1}}{x^2 + x + 1} = x^{d+1} (1 + x^3 + x^6 + \dots - x - x^4 - x^7 - \dots),$$

Corollary 2.2 is equivalent to the congruence relation given by Pan and Sun

$$\sum_{k=0}^{p-1} \binom{2k}{k+d} \equiv_p \binom{p-d}{3}.$$

[Of course this case, and many other ones, for *small*  $r$ , are easily humanly generated.]

Using this approach, we found many congruences, including the congruences for the sums of generalized central trinomial coefficients that were considered by Sun [7].

When  $Q(x)/(P(x) - 1)$  is a rational function such that every root of the denominator is a root of unity, the coefficient of  $x^k$  in  $Q(x)/(P(x) - 1)$  can be expressed as a *quasi-polynomial* in  $k$ . We can search for this quasi-polynomial by the method of undetermined coefficients and thus derive theorems presenting *explicit* forms for the congruences. This is implemented by the procedure `TheoQP` in our Maple package `CTcong.txt`. The command-line is

$$\text{TheoQP}(P, Q, x, p, r, d),$$

where  $P, Q$  are the two Laurent polynomials in  $x$ ,  $p$  is the symbol standing for the prime,  $r$  is as above, and  $d$  is the expected degree of the searched quasi-polynomial. (In practice, you start, optimistically, with  $d = 0$ , and if it fails, you keep increasing  $d$  to 1, then 2, until you either find something, or give up.)

For example, typing

$$\text{TheoQP}(1/x+2+x, 1, x, p, 2, 0);$$

yields

**Corollary 2.3:**

$$\sum_{k=0}^{2p-1} \binom{2k}{k} \equiv_p \begin{cases} 3, & \text{if } p \equiv 1 \pmod{3}, \\ -3, & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

For more examples, we refer to the webpage <http://www.math.rutgers.edu/~zeilberg/mamirim/mamirimhtml/ctcong.html>.

**3. Reduction**

In this section, we consider a further reduction of the coefficients  $S_p, S_{2p}, \dots$  in Equation (2.1). We find that in some cases, the set  $\{S_p \pmod p\}$  of residues is a finite subset of  $\mathbb{Q}$  when  $p$  runs over all primes.

First, let us consider the coefficients  $S_k$  given by

$$\frac{ux + v}{a + bx + cx^2} = \sum_{k=0}^{\infty} S_k x^k,$$

where  $u, v, a, b, c$  are integers,  $a \neq 0$  and  $a + bx + cx^2$  is irreducible over  $\mathbb{Q}$ . Let  $\Delta = b^2 - 4ac$  be the discriminant of  $ax^2 + bx + c$ . Since  $a + bx + cx^2$  is irreducible,  $\Delta \neq 0$ , and hence  $\Delta \not\equiv_p 0$  except for finitely many primes  $p$ .

If  $\Delta$  is a square in the finite field  $\mathbb{F}_p$ , then  $a + bx + cx^2$  is reducible in  $\mathbb{F}_p$  so that

$$\frac{ux + v}{a + bx + cx^2} \equiv_p \frac{A}{1 - \alpha x} + \frac{B}{1 - \beta x},$$

for some  $A, B, \alpha, \beta \in \mathbb{F}_p$ . We thus find that

$$S_{rp} = A\alpha^{rp} + B\beta^{rp} \equiv_p A\alpha^r + B\beta^r = S_r.$$

If  $\Delta$  is not a square in  $\mathbb{F}_p$ , then  $a + bx + cx^2$  is irreducible in  $\mathbb{F}_p$ . Let us consider the extension field  $\mathbb{F}_p(\alpha)$  with  $a\alpha^2 + b\alpha + c = 0$  and  $\alpha \in \mathbb{C}$ . Let  $\beta \in \mathbb{C}$  be another root of the equation  $ax^2 + bx + c = 0$ . By the property of the Frobenius automorphism [8], it follows that in the extension field  $\mathbb{F}_p(\alpha)$ ,

$$\alpha^p = \beta, \quad \beta^p = \alpha.$$

Hence in the field  $\mathbb{F}_p(\alpha)$ , we have

$$S_{rp} = A\alpha^{rp} + B\beta^{rp} = A\beta^r + B\alpha^r = \frac{c^r}{a^r} (A\alpha^{-r} + B\beta^{-r}) = \frac{c^r}{a^r} S_{-r},$$

where  $S_{-r}$  is determined by the initial values  $S_0, S_1$  and the recurrence relation

$$aS_n + bS_{n-1} + cS_{n-2} = 0, \quad n \in \mathbb{Z}.$$

Since  $S_{rp}$  and  $S_{-r}$  are both rational numbers, we obtain that  $S_{rp} \equiv_p \frac{c^r}{a^r} S_{-r}$ .

In general, let  $q(x)$  be an irreducible polynomial in  $\mathbb{Z}[x]$  of degree  $d$  with non-zero constant term and  $\alpha_1, \dots, \alpha_d$  be the  $d$  roots of  $x^d q(1/x)$  in  $\mathbb{C}$ . If the splitting field  $\mathbb{Q}(\alpha_1, \dots, \alpha_d)$  equals  $\mathbb{Q}(\alpha_j)$  for some  $1 \leq j \leq d$ , we say that  $q(x)$  is *simple*. Clearly, every irreducible polynomial of degree 2 is simple.

We have the following finiteness theorem regarding the congruences.

**Theorem 3.1:** *Let  $P(x), Q(x)$  be two Laurent polynomials in  $x$  with integer coefficients and*

$$a_k = \text{CT } P(x)^k Q(x).$$

*Suppose that each irreducible factor  $q(x) \neq x$  of the denominator of  $Q(x)/(P(x) - 1)$  is simple. Then there exists a finite subset  $A$  of  $\mathbb{Q}$  such that for any prime  $p$ ,*

$$\sum_{k=0}^{rp-1} a_k \equiv_p a,$$

for some  $a \in A$ .

**Proof:** By Theorem 2.1, for a sufficiently large prime  $p$ ,  $\sum_{k=0}^{rp-1} a_k$  modulo  $p$  is a linear combination of  $S_0, S_p, S_{2p}, \dots$ , where  $S_k$  is the coefficient of  $x^k$  in the series

$$R(x) = \frac{Q(x)}{P(x) - 1}.$$

To prove the theorem, it suffices to show that for a fixed integer  $n$ , the set

$$\bigcup_p \{S_{np} \pmod p\}$$

of residues is finite when  $p$  runs over all primes.

Consider the partial fraction decomposition of  $R(x)$  over  $\mathbb{Q}$

$$R(x) = g(x) + \sum_{i=1}^m \frac{h_i(x)}{q_i(x)^{\ell_i}},$$

where  $g(x)$  is a Laurent polynomial over  $\mathbb{Q}$  and for each  $i = 1, \dots, m$ ,  $q_i(x) \in \mathbb{Z}[x]$  is irreducible and  $h_i(x) \in \mathbb{Z}[x]$  is a polynomial with  $\deg h_i(x) < \deg q_i(x)$ . In order to prove the finiteness of the set  $\bigcup_p \{S_{np} \pmod p\}$ , it suffices to show that the residues of the coefficients of each summand form a finite set.

Let  $h(x)/q(x)^\ell$  be one summand and

$$\sum_{k=0}^{\infty} s_k x^k = \frac{h(x)}{q(x)^\ell}.$$

Let  $\tilde{q}(x) = x^d q(1/x)$  where  $d = \deg q(x)$  and let  $\alpha_1, \dots, \alpha_d$  be the roots of  $\tilde{q}(x)$ . By assumption, we have  $\mathbb{Q}(\alpha_1, \dots, \alpha_d) = \mathbb{Q}(\alpha)$  with  $\alpha \in \{\alpha_1, \dots, \alpha_d\}$ . Denote the splitting

field  $\mathbb{Q}(\alpha)$  by  $K$ . Since  $q(x)$  is irreducible, we have

$$K = \left\{ \frac{a_0 + a_1\alpha + \cdots + a_{d-1}\alpha^{d-1}}{b} : a_0, \dots, a_{d-1}, b \in \mathbb{Z} \right\}.$$

Let  $p$  be a prime larger than the maximal irreducible factor of the leading coefficient of  $\tilde{q}(x)$ . Then

$$K_p := \left\{ \frac{a_0 + a_1\alpha + \cdots + a_{d-1}\alpha^{d-1}}{b} : p \nmid b \right\} \subset K$$

form a subring of  $K$ . There is a natural ring homomorphism  $\tau : K_p \rightarrow \mathbb{F}_p[x]/\langle \tilde{q}(x) \rangle$  given by

$$\tau \left( \frac{a_0 + a_1\alpha + \cdots + a_{d-1}\alpha^{d-1}}{b} \right) = a_0b^{-1} + a_1b^{-1}x + \cdots + a_{d-1}b^{-1}x^{d-1}.$$

Clearly, the kernel of the map  $\tau$  is  $pK_p$ .

It is well-known that the coefficients  $s_k$  can be expressed as

$$s_k = \sum_{i=1}^d f_i(k)\alpha_i^k,$$

where  $f_i(k)$  is a polynomial over  $\mathbb{Q}(\alpha)$  of degree less than  $\ell$ . It is easy to see that

$$\tau(f_i(np)) = \tau(c_i),$$

where  $c_i$  is the constant term of  $f_i(x)$ . Since  $\tilde{q}(x) \in \mathbb{Z}[x]$ , for each  $i = 1, \dots, d$ , we have

$$\tilde{q}(\tau(\alpha_i^p)) = \tau(\tilde{q}(\alpha_i^p)) = \tau((\tilde{q}(\alpha_i))^p) = 0.$$

Hence  $\tau(\alpha_i^p) = \tau(\alpha_j)$  for some  $1 \leq j \leq d$ . Let  $\sigma$  be the map given by  $\tau(\alpha_i^p) = \tau(\alpha_{\sigma(i)})$ . We thus have

$$\tau(s_{np}) = \sum_{i=1}^d \tau(c_i)\tau(\alpha_{\sigma(i)}^n) = \tau \left( \sum_{i=1}^d c_i\alpha_{\sigma(i)}^n \right).$$

Let

$$\sum_{i=1}^d c_i\alpha_{\sigma(i)}^n = r_0 + r_1\alpha + \cdots + r_{d-1}\alpha^{d-1}.$$

Since  $\tau(s_{np}) \in \mathbb{Q}$ , we have

$$\tau(s_{np}) = \tau(r_0),$$

and hence  $s_{np} \equiv_p r_0$ . Noting that there are only finitely many choices for  $\sigma$ , hence the set  $\cup_p \{s_{np} \pmod p\}$  is finite. □

**Example:** Suppose that

$$P(x) = \frac{x^3 - 2x + 1}{x}, \quad \text{and} \quad Q(x) = 1.$$



We have

$$\frac{Q(x)}{P(x) - 1} = \frac{x}{x^3 - 3x + 1}.$$

Let

$$\alpha = -0.532\dots, \quad \beta = 0.6527\dots, \quad \gamma = 2.879\dots$$

be the three roots of  $x^3 - 3x^2 + 1$ . Using the approximate values of the three roots, we may use the LLL algorithm [4] to find integral relations among  $\beta, \gamma$  and powers of  $\alpha$ . Using Maple, we get two possible relations

$$\beta = 2 + 2\alpha - \alpha^2, \quad \gamma = 1 - 3\alpha + \alpha^2. \quad (3.1)$$

It is easy to verify that

$$(2 + 2\alpha - \alpha^2)^3 - 3(2 + 2\alpha - \alpha^2)^2 + 1 = 0,$$

and

$$(1 - 3\alpha + \alpha^2)^3 - 3(1 - 3\alpha + \alpha^2)^2 + 1 = 0.$$

It follows that  $2 + 2\alpha - \alpha^2$  and  $1 - 3\alpha + \alpha^2$  are roots of  $x^3 - 3x^2 + 1$ . So we obtain the relations in (3.1). Therefore,  $\mathbb{Q}(\alpha, \beta, \gamma) = \mathbb{Q}(\alpha)$  and hence  $x^3 - 3x + 1$  is simple. By Theorem 3.1, the set

$$\left\{ \sum_{k=0}^{2p-1} \text{CT } P(x)^k \pmod{p} \right\}$$

is finite. In fact, when  $p > 3$ , the only possibilities of the residues are  $-1$  and  $2$ .

We conclude with an example where the denominator is not simple.

**Example:** Let

$$P(x) = -2x^2 + 1 + \frac{1}{x}, \quad Q(x) = 1.$$

Then

$$\frac{Q(x)}{P(x) - 1} = \frac{x}{1 - 2x^3} = x + 2x^4 + 2^2x^7 + \dots.$$

Hence for  $p \equiv 1 \pmod{3}$ , we have

$$\sum_{k=0}^{p-1} \text{CT } P(x)^k \equiv_p 2^{\frac{p-1}{3}}.$$

It seems that the set  $\{2^{\frac{p-1}{3}} \pmod{p}\}$  is not finite.

## Acknowledgements

We wish to thank Zhi-Wei Sun for valuable suggestions.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, and the National Science Foundation of China.

## References

- [1] S. Ahlgreen and K. Ono, *A Gaussian hypergeometric series evaluation and Apéry number congruences*, J. Reine. Angew. Math. 518 (2000), pp. 187–212.
- [2] F. Beukers, *Another congruence for the Apéry numbers*, J. Number Theory 25(2) (1987), pp. 201–210.
- [3] J. Greene, *Hypergeometric functions over finite fields*, Trans. Amer. Math. Soc. 301(1) (1987), pp. 77–101.
- [4] A.K. Lenstra, H.W. Lenstra, and L. Lovász, *Factoring polynomials with rational coefficients*, Math. Ann. 261(4) (1982), pp. 515–534.
- [5] H. Pan and Z.-W. Sun, *A combinatorial identity with application to Catalan numbers*, Discrete Math. 306(16) (2006), pp. 1921–1940.
- [6] E. Rowland and D. Zeilberger, *A case study in meta-automation: Automatic generation of congruence automata for combinatorial sequences*, J. Differ. Equ. Appl. 20(7) (2014), pp. 973–988.
- [7] Z.-W. Sun, *Congruences involving generalized central trinomial coefficients*, Sci. China Math. 57(7) (2014), pp. 1375–1400.
- [8] Z.-X. Wan, *Lectures on Finite Fields and Galois Rings*, World Scientific, New Jersey, 2003.