

June 16, 2016

1. SUPER-CONJECTURE 1

Super-Conjecture 1.1. *Let $p \geq 5$ be a prime, and*

$$A = \sum_{n=0}^{p-1} \binom{2n}{n}.$$

Then

$$\begin{aligned} A &\equiv 1 \pmod{p^2} \text{ if } p \equiv 1 \pmod{3}, \\ A &\equiv -1 \pmod{p^2} \text{ if } p \equiv 2 \pmod{3}. \end{aligned}$$

Proof. Already from the paper we know

$$A = CT \frac{(1+x)^{2p}}{x^{p-1}(1+x+x^2)} = CT \frac{(1-x)(1+x)^{2p}}{x^{p-1}(1-x^3)}.$$

So

$$(1) \quad A = \sum_{k \geq 0} \left(\binom{2p}{p-1-3k} - \binom{2p}{p-2-3k} \right)$$

CASE 1 $p \equiv 1 \pmod{3}$. In this case, the $k = (p-1)/3$ term in (1) is 1, and all of the other terms are divisible by p ,

$$A = 1 + pB$$

where

$$B = \frac{1}{p} \sum_{0 \leq k < (p-1)/3} \left(\binom{2p}{p-1-3k} - \binom{2p}{p-2-3k} \right).$$

We'll show that the integer B satisfies $B \equiv 0 \pmod{p}$, so that $A \equiv 1 \pmod{p^2}$.

Since

$$\begin{aligned} \frac{1}{p} \binom{2p}{p-1-3k} &\equiv 2 \frac{(p-1)!}{(p-1-3k)!(3k+1)!} \pmod{p}, \\ \frac{1}{p} \binom{2p}{p-2-3k} &\equiv 2 \frac{(p-1)!}{(p-2-3k)!(3k+2)!} \pmod{p} \end{aligned}$$

we see that

$$B \equiv 2 \sum_{0 \leq k < (p-1)/3} \left(\frac{(p-1)!}{(p-1-3k)!(3k+1)!} - \frac{(p-1)!}{(p-2-3k)!(3k+2)!} \right) \pmod{p}$$

However, by trisecting the binomial theorem this sum of integers is 0

$$\sum_{0 \leq k < (p-1)/3} \left(\frac{(p-1)!}{(p-1-3k)!(3k+1)!} - \frac{(p-1)!}{(p-2-3k)!(3k+2)!} \right) = 0,$$

so

$$B \equiv 0 \pmod{p}.$$

CASE 2 $p \equiv 2 \pmod{3}$. In this case the second term with $k = (p-2)/3$ gives -1 . We proceed as in CASE 1, the trisection identity we need is

$$\sum_{0 \leq k \leq (p-2)/3} \frac{(p-1)!}{(p-1-3k)!(3k+1)!} - \sum_{0 \leq k \leq (p-5)/3} \frac{(p-1)!}{(p-2-3k)!(3k+2)!} = 0,$$

□

2. SUPER-CONJECTURE 1"

Super-Conjecture 2.1. *Let $p \geq 5$ be a prime, r be a positive integer, and*

$$A = \sum_{n=0}^{rp-1} \binom{2n}{n}.$$

Then

$$\begin{aligned} A &\equiv \alpha_r \pmod{p^2} \text{ if } p \equiv 1 \pmod{3}, \\ A &\equiv -\alpha_r \pmod{p^2} \text{ if } p \equiv 2 \pmod{3}. \end{aligned}$$

where

$$\alpha_r = \sum_{n=0}^{r-1} \binom{2n}{n}.$$

Proof. (sketch, assuming a fact (3) you probably know, if true) This time

$$A = CT \frac{(1+x)^{2rp}}{x^{rp-1}(1+x+x^2)} = CT \frac{(1-x)(1+x)^{2rp}}{x^{rp-1}(1-x^3)}.$$

So

$$(2) \quad A = \sum_{k \geq 0} \left(\binom{2rp}{rp-1-3k} - \binom{2rp}{pr-2-3k} \right)$$

As before, we separate the terms which have a p for sure, namely those whose binomial denominator is not a multiple of p from those whose binomial denominator is a multiple of p . The non-multiples of p will sum to 0 as before, and we are left with the extra terms.

CASE 1 $p \equiv 1 \pmod{3}$. Here the terms with denominator parameter a multiple of p are

$$EXTRA = \sum_{j \geq 0} \left(\binom{2rp}{(r-1-3j)p} - \binom{2rp}{(r-2-3j)p} \right).$$

Assuming

$$(3) \quad \binom{Dp}{Ep} \equiv \binom{D}{E} \pmod{p^2}$$

then

$$EXTRA \equiv \sum_{j \geq 0} \left(\binom{2r}{r-1-3j} - \binom{2r}{r-2-3j} \right) = \alpha_r \pmod{p^2}$$

by (1), which works for any positive integer p .

CASE 2 $p \equiv 2 \pmod{3}$. Here the terms with denominator parameter being a multiple of p are

$$EXTRA = \sum_{j \geq 0} \left(\binom{2rp}{(r-2-3j)p} - \binom{2rp}{(r-1-3j)p} \right).$$

then

$$EXTRA \equiv \sum_{j \geq 0} \left(\binom{2r}{r-2-3j} - \binom{2r}{r-1-3j} \right) = -\alpha_r \pmod{p^2}$$

again by (1).

□

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455
E-mail address: `stanton@math.umn.edu`