

A Fibonacci-Counting Proof Begged by Benjamin and Quinn

Doron ZEILBERGER¹

Arthur Benjamin and Jennifer Quinn, in their delightful book[1], asked for *proofs that count* of the identities

$$\sum_i \binom{2n}{i} F_{2i} = 5^n F_{2n} \quad , \quad \sum_i \binom{2n}{i} F_{2i+1} = 5^n F_{2n+1} \quad .$$

Here goes. For any vector of integers u , let $|u|$ denote the sum of its entries. Let, for $\epsilon = 1, 0$,

$$A_\epsilon(n) := \{(w, u) : w \in \{0, 1\}^{2n} \quad , \quad u \in \{1, 2\}^* \quad , \quad 2|w| - |u| = \epsilon \quad \} \quad ,$$

$$B_\epsilon(n) := \{(w, u) : w \in \{1, 2, 3, 4, 5\}^n \quad , \quad u \in \{1, 2\}^* \quad , \quad |u| = 2n - \epsilon \quad \} \quad .$$

The left sides of the identities *count* $A_\epsilon(n)$ and the right sides count $B_\epsilon(n)$, ($\epsilon = 1, 0$). Consider a directed graph with vertices C_0 and C_1 , where there are ten edges joining C_0 to C_0 , five edges joining C_0 to C_1 , five edges joining C_1 to C_0 , and five edges joining C_1 to C_1 . I claim that both $A_\epsilon(n)$ and $B_\epsilon(n)$ are in natural bijection with the set of n -step walks on this directed graph that start at C_ϵ ($\epsilon = 0, 1$).

If $(w, u) \in A_1(n)$, start your journey at vertex C_1 . Let u' be the shortest prefix of u such that $|u'| \geq 2(w_1 + w_2) - 1$, and write $u = u'u''$. Of course $|u'| = 2(w_1 + w_2) - 1$ or $|u'| = 2(w_1 + w_2)$. There are five possibilities for each of these scenarios, and once and for all, name the five edges from C_1 to C_0 and the five edges from C_1 to C_1 by these possibilities, respectively, and follow the appropriate edge, at the same time shrinking (w, u) by replacing it by (w', u'') where w' is w with its first two entries chopped. In the former case (w', u'') belongs to $A_0(n - 1)$, and in the latter case to $A_1(n - 1)$. Now continue recursively.

If $(w, u) \in A_0(n)$, start your journey at vertex C_0 . Let u' be the shortest prefix of u such that $|u'| \geq 2(w_1 + w_2)$, and write $u = u'u''$. Of course $|u'| = 2(w_1 + w_2)$ or $|u'| = 2(w_1 + w_2) + 1$. There are ten possibilities for the former scenario and five possibilities for the latter. Name, once and for all, the ten edges from C_0 to C_0 , and the five edges from C_0 to C_1 by these possibilities, respectively, and follow the appropriate edge, at the same time shrinking (w, u) by replacing it by (w', u'') where w' is w with its first two entries chopped. In the former case (w', u'') belongs to $A_0(n - 1)$, and in the latter case to $A_1(n - 1)$. Now continue recursively.

In the same vein, but much simpler, $B_0(n)$ and $B_1(n)$ correspond to n -step walks starting at C_0 and C_1 on that very same directed graph. The details are left to the reader.

REFERENCE

1. A. Benjamin and J. Quinn, "*Proofs that Really Count: The Art of Combinatorial Proofs*", The Math. Assoc. of Amer. 2003.

¹ Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA. zeilberg@math.rutgers.edu , <http://www.math.rutgers.edu/~zeilberg/>. May 24, 2004. Supported in part by the NSF.