

AUTOMATIC CONJECTURING AND PROVING OF EXACT VALUES OF SOME INFINITE FAMILIES OF INFINITE CONTINUED FRACTIONS

ROBERT DOUGHERTY-BLISS AND DORON ZEILBERGER

ABSTRACT. Inspired by the recent pioneering work, dubbed “The Ramanujan Machine” by Raayoni et al. [6], we (automatically) [rigorously] prove some of their conjectures regarding the exact values of some specific infinite continued fractions, and generalize them to evaluate infinite families (naturally generalizing theirs). Our work complements their beautiful approach, since we use *symbolic* rather, than *numeric* computations, and we instruct the computer to not only discover such evaluations, but at the same time prove them rigorously.

In fond memory of Richard Askey (1933-2019), the great guru of special functions

“Special Functions” is a misnomer, they should be called “Useful functions”-Paul Turan [often quoted by Richard Askey]

1. PREFACE

This tribute to Dick Asky is yet another example how special functions are so useful. Combined with *symbolic computation* (that he was one of the first to realize its great potential) they are even more useful.

2. INTRODUCTION

Continued fractions, possibly finite expressions of the form

$$a_0 + \frac{b_1}{a_1 + \frac{b_2}{a_2 + \frac{b_3}{a_3 + \cdots}}} = a_0 + \frac{b_1}{|a_1|} + \frac{b_2}{|a_2|} + \frac{b_3}{|a_3|} + \cdots ,$$

have been employed since antiquity. Aristarchus of Samos (b. 310 BCE), the great Ancient Greek astronomer and first-rate mathematician, argued that r , the radius

Date: May 4, 2020.

of the Sun divided by the radius of Earth, satisfies

$$\frac{r}{r-1} > \frac{71755875}{61735500}.$$

To provide a simpler bound, he replaced $71755875/61735500$ with $43/37$ and claimed that the same bound would hold. Continued fractions lurk here; the later fraction appears in the former's continued fraction expansion:

$$\begin{aligned} \frac{43}{37} &= 1 + \frac{1}{|6} + \frac{1}{|6} \\ \frac{71755875}{61735500} &= 1 + \frac{1}{|6} + \frac{1}{|6} + \frac{1}{|4} + \frac{1}{|1} + \frac{1}{|2} + \frac{1}{|1} + \frac{1}{|2} + \frac{1}{|1} + \frac{1}{|6}. \end{aligned}$$

Thus Aristarchus and the other Greeks certainly knew *something* about continued fractions, though the full extent of their knowledge is lost to history (see [5, p. 336] and [1, ch. 1]).

The Italian Pietro Cataldi (b. 1548) was the first mathematician to expound a proper theory and notation for continued fractions. He noted that continued fractions, when they converge, alternate between being above and below their limits, and essentially gave us the notation

$$a_0 + \frac{b_1}{|a_1} + \frac{b_2}{|a_2} + \frac{b_3}{|a_3} + \dots$$

which we employ here.

Cataldi may have been the first to present the theory of continued fractions, but the great mathematician Leonhard Euler was responsible for the theory's explosion and widespread application. In *Introductio in Analysin Infinitorum*, his influential analysis textbook, Euler treated continued fractions extensively and showed that they are in a sort of correspondence with infinite series. (Every series can be represented as a continued fraction, and every continued fraction can be represented as a series.) Based on this reasoning, he gave some of the first examples of continued fractions for famous constants:

$$\begin{aligned} e &= \frac{1}{|2} + \frac{1}{|1} + \frac{1}{|2} + \frac{1}{|1} + \frac{1}{|1} + \frac{1}{|4} + \frac{1}{|1} + \frac{1}{|1} + \frac{1}{|6} + \dots \\ \frac{e+1}{e-1} &= \frac{1}{|2} + \frac{1}{|6} + \frac{1}{|10} + \frac{1}{|14} + \frac{1}{|18} + \frac{1}{|22} + \dots \end{aligned}$$

The most well-known general case of Euler’s results is now called *Euler’s continued fraction*:

$$\frac{1}{1 - \frac{r_1}{1 + r_1 - \frac{r_2}{1 + r_2 - \frac{r_3}{1 + r_3 - \dots}}}} = \sum_{k \geq 0} \prod_{j=1}^k r_j.$$

Such identities are intrinsically fascinating, but continued fraction expansions have found wide applications in number-theoretic irrationality proofs. There is always hope that the correct continued fraction will provide a Diophantine approximation sufficiently nice to prove the irrationality of a famous constant, *à la* Roger Apéry’s proof that $\zeta(3)$ is irrational (see [9]).

In this paper, we would like to combine the spirits of Euler and Aristarchus to present an experimental method for automatically discovering (and proving!) continued fraction expansions. We were inspired by “The Ramanujan Machine,” a recent *inverse symbolic calculator* that numerically conjectures continued fraction expansions involving well-known constants [6]. For example, one of their conjectures is

$$(1) \quad \frac{e}{e-2} = 4 - \frac{1}{5} - \frac{2}{6} - \frac{3}{7} - \dots = 4 - \frac{1}{5 - \frac{2}{6 - \frac{3}{7 - \dots}}}.$$

Another is

$$\frac{1}{e-2} = \frac{1}{|1} - \frac{1}{|1} + \frac{2}{|1} - \frac{1}{|1} + \frac{3}{|1} - \frac{1}{|1} + \dots$$

The potential for automatic conjectures is intriguing, but it occurred to us that *symbolic* experiments could yield automatic proofs rather than conjectures. These experiments led to the proof of some of their conjectures and to the discovery and proof of three new infinite families of continued fractions.

Before describing our results, we need some notation which deviates slightly from the norm.

Definition 1. Given two sequences $a(n)$ and $b(n)$ and an integer $m \geq 1$, the m th *convergent* of their general continued fraction is defined by

$$[a(n) : b(n)]_1 = a(1)$$

$$[a(n) : b(n)]_{m+1} = a(1) + \frac{b(1)}{[a(n+1) : b(n+1)]_m}, \quad m \geq 1,$$

whenever these expressions are well-defined. If all convergents are well-defined and $\lim_{m \rightarrow \infty} [a(n) : b(n)]_m$ exists, then the *general continued fraction* $[a(n) : b(n)]$ is defined as

$$[a(n) : b(n)] = \lim_{m \rightarrow \infty} [a(n) : b(n)]_m = a(1) + \frac{b(1)}{a(2) + \frac{b(2)}{a(3) + \dots}}$$

Classical, *simple* continued fractions are those where $b(n) = 1$ for all n . We shall often make reference to a continued fraction $[a(n) : b(n)]$ before we have established its existence; in such cases we usually refer to the convergents or to the sequences $a(n)$ and $b(n)$ themselves.

There is ambiguity in the notation $[a(n) : b(n)]$ —what is the sequence variable?—but we shall always use n as the sequence variable, and other letters as parameters. For example, the sequences in $[n2^m : 1]$ are $a(n) = n2^m$ and $b(n) = 1$, not $a(m) = n2^m$ and $b(m) = 1$.

Our principal discoveries are the following three infinite families:

$$(2) \quad [n + k : an] = \frac{a^D}{(D-1)! \left(e^a - \sum_{s=0}^{D-1} \frac{a^s}{s!} \right)}, \quad D = a + k + 1 \geq 1$$

$$(3) \quad [an^2 + bn + 1 : -an^2 - bn] = \frac{4F(a, b) + 2(2a + b)(a + b + 1)}{4F(a, b) + 2(2a + b)}, \quad a \neq 0, b \geq 0$$

$$(4) \quad [(n-1)^k + n^k, -n^{2k}] = \frac{1}{\zeta(k)}, \quad k \geq 2$$

where

$$F(a, b) = 2 \sum_{k \geq 0} \frac{1}{(k+2)! (3 + b/a)^{\bar{k}} a^k}$$

and

$$\zeta(s) = \sum_{k \geq 1} \frac{1}{k^s}$$

is *Riemann's zeta function*. The symbols $x^{\bar{k}} = x(x+1)\cdots(x+k-1)$ and $x^{\underline{k}} = x(x-1)\cdots(x-k+1)$ are called the *rising and falling factorials*, respectively. Note that the rising factorial, $x^{\bar{k}}$, is often denoted by $(x)_k$, and hence the falling factorial, $x^{\underline{k}}$, is written $(x-k+1)_k$.

We shall get to discovering and proving our results in the following sections, but first one more historical remark.

As in continued fractions as it is elsewhere, so prolific was Euler that new results should be checked against his work for duplicates. For example, what is usually referred to as *Gauss' continued fraction* was discovered by Euler nearly thirty years before Gauss was born! (See [2], or [3] for an English translation.) In Section 4.4 we

shall discuss Euler's continued fraction in relation to ours and argue that, as far as we can tell, our three new infinite families do not follow Euler's.

For a more modern introduction to continued fractions, see [4, ch. X], and [11] for the analytic theory in particular.

3. EXPERIMENTAL CONTINUED FRACTIONS: THE MAPLE PACKAGE GCF.TXT

This article is accompanied by a Maple package, `GCF.txt` available from the webpage of this article

<https://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/gcf.html>,

where one can also find two sample input and output files with computer-generated articles for many special cases of our results. Indeed, while Section 4 contains human-readable proofs, our results were *discovered and proved* through symbolic experimentation with `GCF.txt`.

This is quite different than “The Ramanujan Machine” (TRM) described in [6], both at a high-level and practically. At a high level, TRM takes a constant and tries to fit a family of continued fractions to it. Our experiments work in the opposite direction: we construct a family of continued fractions and try to guess the constants that they generate. While TRM produces only conjectures, our Maple package produces *proofs*. Of course, the dazzling conjectures of TRM are motivation for everything in `GCF.txt`.

Here are short descriptions of the main procedures.

- `GCF(L)`: Inputs a finite list of pairs of **numbers**, $[[a_1, b_1], [a_2, b_2], \dots, [a_k, b_k]]$, outputs the finite (generalized) continued fraction it evaluates to. For example `GCF([[1, 1]$10])` gives $\frac{89}{55}$ (the tenth convergent to the Golden ratio) while `GCF([seq([4*i-2, 1], i=1..20)])`;

gives the 20th convergent to Euler's famous continued fraction for $\frac{e+1}{e-1}$.

$$\frac{376958612213530151806235679061}{174199042280794948413485144460} \quad ,$$

that agrees with it to 60 decimal digits.

- `GCFab(a, b, n, K)`: Now the input parameters, a and b , are *expressions* in the symbol n . K is again a positive integer, and the output is the same as `GCF` applied to

$$[[a(1), b(1)], \dots, [a(K), b(K)]] \quad .$$

- `RDB(a, b, n)`: Inputs expressions a and b in n and outputs (if successful) the explicit expressions for the numerator and denominator of the n -th convergent of the infinite continued fraction, as well as its limit, and the error by using the 50-th convergent. It uses Maple's `rsolve` command that is not guaranteed to work (most linear recurrences are not solvable in closed-form, and even amongst those that are, Maple [probably] does not know how to solve all of them). But whenever it succeeds, it can

be fully trusted, i.e. it gives a **proved** result. Under the hood `rsolve` uses Mark Van Hoeij's algorithm [10], but often the answer can be checked by hand. For instance, in the case of (2), the relevant expressions involve the *incomplete Gamma function*, and can be checked *ab initio* without Maple by using the incomplete Gamma function's well-known (and easily proved) first-order inhomogeneous recurrence.

- `Yaron(k1, a1, n, G)`: is a specialization of `RDB`, namely `RDB(n+k1, a1*n, n)`, but for expository clarity, the incomplete Gamma function `GAMMA(n+2, -a1)` ($\Gamma(n+2, -a1)$) is denoted by `G[n]`. It also proves its results rigorously.

- `YaronV(k1, a1, n, G)`: a verbose form of `Yaron(k1, a1, n, G)`. It outputs a computer-generated article.

Note that for each specific pair of sequences, `RDB` gives the exact evaluation of the infinite continued fraction, but to prove the **general** results of our paper, that were conjectured from the many special cases, we had to 'cheat' and use traditional human mathematics. We believe that much of this human part can also be automated, but leave it to a future paper.

The following section rigorously evaluates an infinite family of general continued fractions, but there is much more to discover. Our infinite family is a quite restricted class of general continued fractions generated by specific linear polynomials. There are other, equally interesting polynomials. For example, `RDB` can discover the identity

$$[3n : -n(2n - 1)] = \frac{4}{3\pi - 8}.$$

We encourage our readers to experiment and discover new, more exotic, families.

4. GENERAL PROOFS

Our main tool for evaluating continued fractions is exploiting their recursive nature.

Definition 2. Given a continued fraction $[a(n) : b(n)]$, define the *numerator* and *denominator* sequences $p(n)$ and $q(n)$ by the recurrences

$$\begin{aligned} p(0) &= a(1) \\ p(1) &= a(1)a(2) + b(1) \\ p(n+2) &= a(n+3)p(n+1) + b(n+2)p(n) \end{aligned}$$

and

$$\begin{aligned} q(0) &= 1 \\ q(1) &= a(2) \\ q(n+2) &= a(n+3)q(n+1) + b(n+2)q(n), \end{aligned}$$

respectively.

Lemma 1. *For all positive integers m ,*

$$\frac{p(m)}{q(m)} = [a(n) : b(n)]_{m+1}.$$

This is a well-known fact from the theory of general continued fractions.

4.1. **First infinite family.** Our general proof of (2) relies on some results in differential equations, so let us first define the necessary objects.

Definition 3. Let

$$M(a, b, z) = \sum_{k \geq 0} \frac{a^{\bar{k}}}{b^{\bar{k}} k!} z^k$$

be the *confluent hypergeometric function of the first-kind*—also known as *Kummer's function*—and

$$U(a, b, z) = \frac{\Gamma(1-b)}{\Gamma(a+1-b)} M(a, b, z) + \frac{\Gamma(b-1)}{\Gamma(a)} z^{1-b} M(a+1-b, 2-b, z)$$

be the *confluent hypergeometric function of the second-kind*—also known as *Tricomi's function*—where

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt$$

is the *gamma function*, defined by the above integral whenever $\Re z > 0$. Also let

$$\Gamma(z, a) = \int_a^\infty e^{-t} t^{z-1} dt$$

be the *incomplete gamma function*.

The key property of $M(a, b, z)$ and $U(a, b, z)$ is that they are linearly independent solutions of *Kummer's differential equation*

$$zw''(z) + (b-z)w'(z) - aw(z) = 0.$$

Kummer's function $M(a, b, z)$ is entire if b is not a nonpositive integer, while $U(a, b, z)$ generally has a pole at the origin. In particular, we have the following well-known result.

Lemma 2. *If $a - b + 1 = -n$ for a nonnegative integer n , then*

$$U(a, b, z) = z^{-a} \sum_{s=0}^n \binom{n}{s} a^{\bar{s}} z^{-s}.$$

See [7, ch. 13] for more details on the confluent hypergeometric functions. Here is a useful lemma from the world of generating functions.

Lemma 3. *Let $f(z)$ be a meromorphic function with a single pole of order $r \geq 2$ at $z_0 \neq 0$. If*

$$f(z) = \sum_{k \geq -r} a_k (z - z_0)^k,$$

then

$$[z^n]f(z) = \frac{(-1)^r a_{-r}}{z_0^{n+r}} \binom{n+r-1}{r-1} (1 + O(1/n)).$$

where $[z^n]f(z)$ is the coefficient on z^n in the expansion of f about the origin.

Proof. Let

$$g(z) = \sum_{-r \leq k < 0} a_k (z - z_0)^k$$

be the principle part of f at z_0 . Expressing $g(z)$ as a power series about the origin in the usual way yields

$$[z^n]g(z) = \sum_{k=1}^r \frac{(-1)^k a_{-k}}{z_0^{k+n}} \binom{n+k-1}{k-1}.$$

In particular, if $r \geq 2$, then this is a polynomial in n of degree $r-1 \geq 1$, and n^{r-1} only appears in the last term. Since $f(z) - g(z)$ is entire, $[z^n](f(z) - g(z)) = O(1)$, and pulling out the leading term after rearranging yields

$$[z^n]f(z) = \frac{(-1)^r a_{-r}}{z_0^{n+r}} \binom{n+r-1}{r-1} (1 + O(1/n))$$

as desired. □

Using the previous lemma, we will now provide asymptotic expansions for $p(n)$ and $q(n)$ for the continued fraction $[n+k : an]$. The key observation is that the exponential generating functions of $p(n)$ and $q(n)$ both satisfy the same second-order differential equation, and that this equation has a nice, meromorphic solution with a single pole at $z = 1$. To avoid repetition, let's first prove a very specific lemma.

Lemma 4. *Let $a(n)$ be a sequence whose exponential generating function $A(z) = \sum_{n \geq 0} \frac{a(n)}{n!} z^n$ satisfies*

$$A(z) = L e^{-\alpha z} M(D, D+2, \alpha(z-1)) + E e^{-\alpha z} U(D, D+2, \alpha(z-1))$$

for some values L and E independent of z , a positive integer D , and a nonzero real α . Then

$$a(n) = E \frac{e^{-\alpha}}{\alpha^{D+1}} D \binom{n+D}{D} (1 + O(1/n)).$$

Proof. Kummer's hypergeometric function is entire, while $U(D, D+2, \alpha(z-1))$ has a single pole at $z=1$. In fact, by Lemma 2, the pole is order $D+1 \geq 2$, and the coefficient on its lowest degree term is D/α^{D+1} . If we write

$$Ee^{-\alpha z}U = e^{-\alpha}Ee^{-\alpha(z-1)}U,$$

then we see that the coefficient on the lowest degree term of $Ee^{-\alpha z}U$ is $Ee^{-\alpha}D/\alpha^{D+1}$, so Lemma 3 implies $a(n) = E\frac{e^{-\alpha}}{\alpha^{D+1}}D\binom{n+D}{D}(1 + O(1/n))$. \square

Theorem 1. *Let a and k be integers. If $D = a + k + 1 \geq 1$, then*

$$[n+k : an] = \frac{a^D}{(D-1)! \left(e^a - \sum_{s=0}^{D-1} \frac{a^s}{s!} \right)},$$

provided that the convergents of the continued fraction are well-defined.

Proof. Let

$$P(z) = \sum_{n \geq 0} \frac{p(n)}{n!} z^n$$

$$Q(z) = \sum_{n \geq 0} \frac{q(n)}{n!} z^n$$

be the exponential generating functions of $p(n)$ and $q(n)$, respectively. By well-known facts about egfs, the recurrences in Definition 2 imply that P and Q both satisfy the second-order differential equation

$$(1-z)f''(z) - (az+k+3)f'(z) - 2af(z) = 0,$$

with initial conditions

$$P(0) = k+1 \quad P'(0) = (k+1)(k+2) + a$$

$$Q(0) = 1 \quad Q'(0) = k+2.$$

This reduces to a special case of *Kummer's equation*. It is easy to check with a computer algebra system that the general solution is

$$f(z) = A(k)e^{-az}M(D, D+2, a(z-1)) + B(k)e^{-az}U(D, D+2, a(z-1))$$

for some sequences $A(k)$ and $B(k)$ which depend on the initial conditions. By Lemma 4, it suffices to compute $B(k)$ for $p(n)$ and $q(n)$ separately.

Let $B_p(a, k)$ be the coefficient on $e^{-az}U$ for $P(z)$, and $B_q(a, k)$ the coefficient on $e^{-az}U$ for $Q(z)$. We may compute these functions by solving the relevant initial condition equations. For instance,

$$B_p(a, k) = (-1)^{a+k} a^{a+k+2} = (-a)^{D+1}.$$

The function $B_q(a, k)$ is significantly more complicated, but still routine to compute. After some coercing, Maple simplifies it as

$$\begin{aligned} B_q(a, k) &= \frac{a(e^a(\Gamma(D+1, a) - \Gamma(D+1))a^{D+1} - a^{2D+1})(-1)^D}{Da^{D+1}} \\ &= (-1)^D \frac{a}{D} (e^a(\Gamma(D+1, a) - \Gamma(D+1)) - a^D). \end{aligned}$$

The incomplete gamma function expression can be written

$$\Gamma(D+1, a) - \Gamma(D+1) = - \int_0^a e^{-t} t^D dt = D! \left(e^{-a} \sum_{s=0}^D \frac{a^s}{s!} - 1 \right),$$

so

$$\begin{aligned} B_q(a, k) &= (-1)^D \frac{a}{D} \left(D! \left(\sum_{s=0}^D \frac{a^s}{s!} - e^a \right) - a^D \right) \\ &= (-1)^D a(D-1)! \left(\sum_{s=0}^{D-1} \frac{a^s}{s!} - e^a \right). \end{aligned}$$

Putting everything together, we have

$$\begin{aligned} [n+k : an] &= \lim_{m \rightarrow \infty} \frac{p(m)}{q(m)} \\ &= \lim_{m \rightarrow \infty} \frac{B_p(a, k)(1 + O(1/m))}{B_q(a, k)(1 + O(1/m))} \\ &= \frac{B_p(a, k)}{B_q(a, k)} \\ &= \frac{a^D}{(D-1)! \left(e^a - \sum_{s=0}^{D-1} \frac{a^s}{s!} \right)}, \end{aligned}$$

as claimed. □

There are some notable special cases. If $a = -k$, then $D = 1$, which yields

$$[n+k : -kn] = \frac{ke^k}{e^k - 1}.$$

If $a = -1$, then $D = k$, and we can write

$$[n+k+1 : -n] = \frac{(-1)^k e}{ek! \sum_{s=0}^k \frac{(-1)^s}{s!} - k!}$$

for nonnegative integers k . Equation 1 is then obtained with $k = 2$:

$$[n + 3 : -n] = \frac{e}{e - 2}.$$

Note that

$$ek! \sum_{s=0}^k \frac{(-1)^s}{s!} = e[k!/e],$$

where $[x]$ denotes the integer nearest to the real x . This is a remarkable coincidence, since $[k!/e]$ is the k th *derangement number*, the number of permutations on k objects with no fixed points. There does not seem to be any immediate combinatorial reason for the derangement numbers to appear.

4.2. Second infinite family. Our second infinite family (3) is much easier to evaluate.

Theorem 2. *Let a and b be nonnegative reals such that $a \neq 0$. Then*

$$[an^2 + bn + 1 : -an^2 - bn] = \frac{4F(a, b) + 2(2a + b)(a + b + 1)}{4F(a, b) + 2(2a + b)},$$

where

$$F(a, b) = 2 \sum_{k \geq 0} \frac{1}{(k + 2)!(3 + b/a)^{\bar{k}} a^k}.$$

Proof. The implied recurrence numerator and denominator recurrences can be solved and easily put into asymptotic form. The solutions are, asymptotically,

$$\begin{aligned} p(n) &= C(a, b, n)((F(a, b)/2 + (2a + b)(a + b + 1)) + O(1/n^2)) \\ q(n) &= \frac{1}{2}C(a, b, n)((F(a, b) + 2(2a + b)) + O(1/n^2)), \end{aligned}$$

where $C(a, b, n)$ is some function. From this, it is easy to see that

$$\lim_{n \rightarrow \infty} \frac{p(n)}{q(n)} = \frac{F(a, b) + 2(2a + b)(a + b + 1)}{F(a, b) + 2(2a + b)}. \quad \square$$

The F function is actually a special case of the general hypergeometric function, and therefore offers numerous opportunities for closed-form evaluation. For example, suppose that $b/a = m - 1/2$ for some nonnegative integer m . Then, from the identity

$$(r - 1/2)^{\bar{k}} = \frac{(2r - 1)^{\overline{2k}}}{4^k r^{\bar{k}}},$$

we have

$$\begin{aligned} (3 + b/a)^{\bar{k}} &= (3 + m - 1/2)^{\bar{k}} \\ &= \frac{(5 + 2m)^{\bar{2k}}}{4^k(3 + m)^{\bar{k}}} \\ &= \frac{(4 + 2m + 2k)!(2 + m)!}{4^k(4 + 2m)!(2 + m + k)!}. \end{aligned}$$

Therefore

$$F(a, b) = \frac{2(4 + 2m)!}{(m + 2)!} \sum_{k \geq 0} \frac{(k + m + 2)!}{(k + 2)!(2k + 2m + 4)!} \left(\frac{4}{a}\right)^k.$$

This remaining sum looks quite burly, but is amenable to routine evaluation after some simplifications. Let us give a brief sketch of how it might work.

In what follows, let us write “ \sim ” to mean “equal up to a multiplicative constant.” First, shifting the summation index by $m + 2$ gives

$$F(a, b) \sim \sum_{k \geq 2} \frac{(k + m)!}{k!(2k + 2m + 1)!} \left(\frac{4}{a}\right)^k.$$

Note that $(k + m)!/k! = (k + m)^{\underline{m}}$, so

$$F(a, b) \sim \sum_{k \geq 2} \frac{(k + m)^{\underline{m}}}{(2k + 2m + 1)!} \left(\frac{4}{a}\right)^k.$$

Now shifting the index by m gives

$$F(a, b) \sim \sum_{k \geq m+2} \frac{k^{\underline{m}}}{(2k + 1)!} \left(\frac{4}{a}\right)^k.$$

At this point we have won, because the series

$$\sum_{k \geq m+2} \frac{1}{(2k + 1)!} z^k$$

is known, and $k^{\underline{m}}$ is a polynomial in k .

More explicitly, set

$$f(z) = \sum_{k \geq m+2} \frac{1}{(2k + 1)!} z^k,$$

and note that $f(z)$ is $z^{-1/2} \sinh \sqrt{z}$ minus a finite number of initial terms. From the elementary theory of generating functions, since k^m is a polynomial in k , we have

$$\sum_{k \geq m+2} \frac{k^m}{(2k+1)!} z^k = (zD)^m f(z),$$

where D is the differentiation operator $Df = f'$. Since we know f , we can carry out the iterated differentiations and then set $z = 4/a$ to obtain an answer. Note that the hyperbolic trigonometric functions are closed under differentiation, so our answer will be in terms of them. The full result is too messy to completely record, but these routine operations can be completed by any computer algebra system.

For example, if $a = 4$ and $b = 6$, then following the above steps will eventually produce

$$\begin{aligned} F(4, 6) &= -308 + 840(\cosh(1) - \sinh(1)) \\ &= -308 + \frac{840}{e}. \end{aligned}$$

In this case we obtain

$$\begin{aligned} [4n^2 + 6n + 1 : -4n^2 - 6n] &= \frac{840/e}{840/e - 280} \\ &= \frac{3}{3 - e}. \end{aligned}$$

Taking $a = 6$ and $b = 9$, we obtain

$$[6n^2 + 9n + 1 : -6n^2 - 9n] = \frac{-9\sqrt{6} \sinh(\sqrt{6}/3) + 18 \cosh(\sqrt{6}/3)}{-9\sqrt{6} \sinh(\sqrt{6}/3) + 18 \cosh(\sqrt{6}/3) - 4}.$$

There are likely many other nice cases.

4.3. Third infinite family. Our third infinite family (4) is also relatively straightforward to establish.

Theorem 3. For integers $k \geq 2$,

$$[(n-1)^k + n^k, -n^{2k}] = \frac{1}{\zeta(k)}.$$

Proof. It is routine to check that $p(n) = (n+1)!^k$ is the numerator sequence for this continued fraction. It is also routine to check (but difficult to discover) that the denominator sequence of the continued fraction is

$$q(n) = (n+1)!^k \left(\frac{\psi(k, n+2)}{(k-1)!} + \zeta(k) \right),$$

where $\psi(k, z)$ is the k th *polygamma function*, which may be defined by

$$\psi(k, z) = (-1)^{k+1} k! \sum_{j \geq 0} \frac{1}{(z+j)^{k+1}}$$

for $z \notin \{-1, -2, \dots\}$. This gives

$$\frac{p(n)}{q(n)} = \frac{1}{\zeta(k)} \frac{1}{\psi(k, n+2)O(1) + 1},$$

where the $O(1)$ is some constant independent of n . It is easy to check with the dominated convergence theorem that $\psi(k, n+2) \rightarrow 0$ as $n \rightarrow \infty$ for all $k \geq 2$, which implies

$$\lim_{n \rightarrow \infty} \frac{p(n)}{q(n)} = \frac{1}{\zeta(k)}$$

as claimed. □

As a demonstration, $k = 3$:

$$\frac{1}{\zeta(3)} = 1 - \frac{1}{9} - \frac{64}{35} - \frac{729}{91} + \dots$$

Just using the terms listed, we have:

$$1 - \frac{1}{9} - \frac{64}{35} - \frac{729}{91} = \frac{1728}{2035} \approx 0.84914004914004914$$

$$\frac{1}{\zeta(3)} \approx 0.83190737258070746.$$

Not a great approximation, but it is something.

4.4. The method of Euler. Our method of proving continued fractions is relatively direct and simple: write down the “obvious” recurrences, solve them, then take a limit. Another approach is to combine some specializations of well-known infinite families with manipulation techniques. This situation is analogous to the problems of evaluating sums or integrals—try some standard techniques, or attempt to contort your problem until it matches some well-known, general result.

One such general result for continued fractions is *Euler’s continued fraction* we mentioned in the introduction:

$$\frac{1}{1 - \frac{r_1}{1 + r_1 - \frac{r_2}{1 + r_2 - \frac{r_3}{1 + r_3 - \dots}}}} = \sum_{k \geq 0} \prod_{j=1}^k r_j.$$

Slight manipulation yields

$$\frac{-r_1}{1 + r_1 - \frac{r_2}{1 + r_2 - \frac{r_3}{1 + r_3 - \dots}}} = \frac{1}{\sum_{k \geq 0} \prod_{j=1}^k r_j} - 1,$$

or, in our notation,

$$[r(n-1) + 1 : -r(n)] = r(0) + \frac{1}{\sum_{k \geq 0} \prod_{j=1}^k r(j)}.$$

Euler's continued fraction directly applies to situations where "numerator + denominator = 1." (In our notation this is "numerator + *previous* denominator = 1.") None of our families are of this form, but Euler's reach can be expanded with an *equivalence transformation*. Namely, it is easy to prove by induction that

$$[a(n) : b(n)] = [c(n-1)a(n) : c(n-1)c(n)b(n)]$$

for any strictly nonzero sequence $\{c(n)\}$ with $c(0) = 1$, provided that both continued fractions exist.

In light of this, one way to evaluate $[a(n) : b(n)]$ is to find sequences $r(n)$ and $c(n)$ such that

$$\begin{aligned} a(n) &= c(n-1)(r(n-1) + 1) \\ b(n) &= -c(n-1)c(n)r(n), \end{aligned}$$

for $n \geq 1$. Once these sequences are in hand, we can write

$$\begin{aligned} [a(n) : b(n)] &= [c(n-1)(r(n-1) + 1) : -c(n-1)c(n)r(n)] \\ &= [r(n-1) + 1 : -r(n)] \\ &= r(0) + \left(\sum_{k \geq 0} \prod_{j=1}^k \frac{b(j)}{a(j+1) + b(j)} \right)^{-1}. \end{aligned}$$

Our problem reduces to computing the remaining sum.

It is not obvious how to find such sequences. For instance, take $a = -1$ and $k = 1$ in (2). Then we must solve the equations

$$\begin{aligned} n + 1 &= c(n-1)(r(n-1) + 1) \\ -n &= -c(n-1)c(n)r(n) \end{aligned}$$

for $c(n)$ and $r(n)$. It is easy to find that

$$r(n) = \frac{n}{c(n-1)n - n + 2c(n-1)},$$

but solving for $c(n)$ is not so easy. It is possible that some suitable *ansatz* could let us *guess* $c(n)$, but this is another question entirely.

Thus, we believe that our results are not specializations of Euler's most well-known continued fraction. It is still possible that there is a simpler derivation of our results from his that we missed, but the main purpose of this paper is to facilitate the *discovery* of new results. Simpler proofs can always be found after the fact.

5. CONCLUSION

We have given a Maple package and three new infinite families of general continued fractions, but there is surely more to do. For instance, it seems like one very general family is $[P(n) + 1 : -P(n)]$ where P is any polynomial. These questions could be investigated with our `GCF.txt`. It also seems likely that our "human proofs" involving asymptotic analysis could be automated, or simplified using one of Euler's many results on continued fractions. We leave these avenues for later papers to explore.

The authors would like to thank the OEIS [8] for its help in identifying promising sequences of integers that arose in the evaluation of these continued fractions, as well as Alex Kontorovich and Danny Calegari for helpful feedback on an earlier version of this paper.

REFERENCES

- [1] Brezinski, Claude. *History of Continued Fractions and Padé Approximants*. Vol. 12. Springer Science & Business Media, 2012.
- [2] Euler, Leonhard. De fractionibus continuis observationes. *Commentarii academiae scientiarum Petropolitanae* (1750): 32-81.
- [3] Euler, L. and Aycok, A., 2018. Observations on continued fractions. arXiv preprint arXiv:1808.07006.
- [4] G.H. Hardy and E.M. Wright. *An Introduction to the Theory of Numbers* (Oxford University Press, 2008).
- [5] Heath, Thomas. *Aristarchus of Samos, the Ancient Copernicus*. Dover Publications, 2004.
- [6] Raayoni, G., Pisha, G., Manor, Y., Mendlovic, U., Haviv, D., Hadad, Y. and Kaminer, I., 2019. The Ramanujan Machine: Automatically Generated Conjectures on Fundamental Constants. arXiv preprint arXiv:1907.00205.
- [7] National Institute of Standards and Technology. (2020), Digital Library of Mathematical Functions <https://dlmf.nist.gov/>.
- [8] OEIS Foundation Inc. (2020), The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>.
- [9] Van der Poorten, Alfred. A Proof that Euler Missed... Apéry's Proof of the Irrationality of $\zeta(3)$. *Math. Intelligencer* 1.4 (1979): 195-203.
- [10] Van Hoeij, M., 1999. Finite Singularities and Hypergeometric Solutions of Linear Recurrence Equations. *Journal of Pure and Applied Algebra*, 139(1-3), pp.109-131.
- [11] Wall, Hubert. *Analytic Theory of Continued Fractions*. Courier Dover Publications, 2018.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY (NEW BRUNSWICK), HILL CENTER-BUSCH CAMPUS, 110 FRELINGHUYSEN RD., PISCATAWAY, NJ 08854-8019, USA

E-mail address: robert.w.bliss@gmail.com

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY (NEW BRUNSWICK), HILL CENTER-BUSCH CAMPUS, 110 FRELINGHUYSEN RD., PISCATAWAY, NJ 08854-8019, USA

E-mail address: doronzeil@gmail.com