

The Gift Exchange Problem, Part II

By Moa Apagodu, David Applegate, Neil Sloane, and Doron Zeilberger

So far we had a k -free (partial) recurrence for $E_\sigma(n, k)$ (for $\sigma = 2, 3$) (that implied recurrences for G_σ) and the trivial n -free (partial) recurrence, (5), for the general $E_\sigma(n, k)$, It turns out that $E_\sigma(n, k)$ satisfies a pure (ordinary) recurrence in k , and that $G_\sigma(n)$ satisfies a linear recurrence with polynomial coefficients, in n , for *every* σ .

Indeed: we have

$$\sum_k E_\sigma(n, k) \frac{y^k}{k!} = \frac{1}{n!} \left(\sum_{i=0}^{\sigma+1} \frac{y^i}{i!} \right)^n$$

Using the J.C.P.Miller ([Z1]) method we get the *pure recurrence* that enables us, for a given n , to compute fast the numbers $E_\sigma(n, k)$ for a *fixed* n , without having to use the $E_\sigma(n-1, k)$'s

$$\sum_{i=0}^{\sigma} \left(\frac{n}{i!} - \frac{k}{(i+1)!} + \frac{1}{(i-1)! + i!} \right) E_\sigma(n, k-i) = 0 \quad .$$

We also have, using Euler's integral representation for the factorial

$$k! = \int_0^\infty e^{-y} y^k dy \quad ,$$

that we have the integral representation

$$G_\sigma(n) = \frac{1}{n!} \int_0^\infty e^{-y} \left(\sum_{i=0}^{\sigma+1} \frac{y^i}{i!} \right)^n dy.$$

It follows from the (discrete) Almkvist-Zeilberger algorithm [AlZ] that $G_\sigma(n)$ satisfies *some* homogeneous linear recurrence with polynomial (in n) coefficients , and it follows from Theorem AZ of [ApZ] that its order is $\sigma + 1$. The Maple package `DavidNeil` accompanying this article can easily compute these recurrences, and the the recurrences for $1 \leq \sigma \leq 6$, together with the implied asymptotics (using the Poincare-Birkhoff-Trijinski method as exposted in [WiZ] and implemented in [Z2]) to the sixth-order, can be found in:

<http://www.math.rutgers.edu/~zeilberg/tokhniot/oDavidNeil1> .

Regarding the *generating function*

$$g_\sigma(x) := \sum_{n=0}^{\infty} G_\sigma(n) x^n,$$

it follows from the (discrete) Almkvist-Zeilberger algorithm [AlZ] that $g_\sigma(x)$ satisfies *some* homogeneous linear differential equation with polynomial (in x) coefficients , and it follows from the obvious analog of Theorem AZ of [ApZ] that its order is $\sigma + 1$.

See

<http://www.math.rutgers.edu/~zeilberg/tokhniot/oDavidNeil2>

for the differential equations for $1 \leq \sigma \leq 6$.

These differential equations transcribe to linear recurrences in $G_\sigma(n)$, that are *not* minimal-order, and for small σ coincide with the ones obtained above using Sister Celine's technique.

Additional References

[AlZ] G. Almkvist and D. Zeilberger, The method of differentiating under the integral sign, J. Symbolic Computation 10 (1990), 571-591.

[ApZ] Moa Apagodu and Doron Zeilberger, *Multi-Variable Zeilberger and Almkvist-Zeilberger Algorithms and the Sharpening of Wilf-Zeilberger Theory*, Adv. Appl. Math. **37**(2006), 139-152.

[WiZ] Jet Wimp and Doron Zeilberger, *Resurrecting the asymptotics of linear recurrences*, J. Math. Anal. Appl. **111**(1985), 162-177.

[Z1] Doron Zeilberger *The J.C.P. Miller Recurrence for exponentiating a polynomial, and its q-Analog*, J. Difference Eqs. and Appls. **1**(1995), 57-60.

[Z2] Doron Zeilberger, *AsyRec: A Maple package for Computing the Asymptotics of Solutions of Linear Recurrence Equations with Polynomial Coefficients*, The Personal Journal of Ekhad and Zeilberger, <http://www.math.rutgers.edu/~zeilberg/pj.html>