

**Proof of a Conjecture of Neil Sloane Concerning Claude Lenormand’s  
“Raboter” Operation (OEIS sequence A318921)**

*Doron ZEILBERGER*

*Dedicated to Neil James Alexander Sloane (born Oct. X, MCMXXXIX) on his turning  
1001111 years young (100101 days late)*

Yesterday, Neil Sloan gave a great talk([S1]), where among many other fascinating results and open problems, he mentioned a conjecture that he made less than two months ago, concerning the “planing” operation (*raboter*) introduced in 2004 by sequence enthusiast Claude Lenormand, and described in [S2]. He also mentioned that he has no idea how hard it is to prove, and made a ‘meta-conjecture’ that it may be a ‘low-hanging fruit’. I will now show that both conjecture and meta-conjecture are true.

By examining the definition of <http://oeis.org/A318921> in terms of the binary representation of  $n$ , it is readily seen that  $r(n)$  may be defined recursively as follows

$$r(n) := \begin{cases} 2r(n/2), & \text{if } n \equiv 0 \pmod{4}; \\ r(\frac{n-1}{2}), & \text{if } n \equiv 1 \pmod{4}; \\ r(n/2), & \text{if } n \equiv 2 \pmod{4}; \\ 2r(\frac{n-1}{2}) + 1, & \text{if } n \equiv 3 \pmod{4}, \end{cases}$$

subject to the initial conditions  $r(0) = 0$  and  $r(1) = 0$ .

Sloane defined, for, integers  $k, k \geq 1$ ,

$$L(k) := \sum_{n=2^k}^{2^{k+1}-1} r(n) \quad ,$$

and conjectured the following

**Fact:**  $L(k) = 2 \cdot 3^{k-1} - 2^{k-1}$  .

Indeed, breaking up the sum into the four congruence classes modulo four, we have

$$\begin{aligned} L(k) &= \sum_{n=2^k}^{2^{k+1}-1} r(n) \\ &= \sum_{m=2^{k-2}}^{2^{k-1}-1} r(4m) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(4m+1) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(4m+2) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(4m+3) \end{aligned}$$

$$\begin{aligned}
&= \sum_{m=2^{k-2}}^{2^{k-1}-1} 2r(2m) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m+1) + \sum_{m=2^{k-2}}^{2^{k-1}-1} (2r(2m+1) + 1) \\
&= 3 \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m+1) + 2 \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m+1) + \sum_{m=2^{k-2}}^{2^{k-1}-1} 1 \\
&= 3 \left( \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m) + \sum_{m=2^{k-2}}^{2^{k-1}-1} r(2m+1) \right) + 2^{k-2} \\
&= 3 \left( \sum_{m=2^{k-1}}^{2^k-1} r(m) \right) + 2^{k-2} = 3L(k-1) + 2^{k-2} .
\end{aligned}$$

Hence the sequence  $L(k)$  satisfies the first-order inhomogeneous recurrence with constant coefficients

$$L(k) - 3L(k-1) = 2^{k-2} .$$

But  $R(k) := 2 \cdot 3^{k-1} - 2^{k-1}$  also satisfies the same recurrence, i.e.

$$R(k) - 3R(k-1) = 2^{k-2} ,$$

(check!), and the fact follows by induction on  $k$ , since it is true for  $k = 1$ .  $\square$

## References

- [S1] N.J.A. Sloane, *Coordination Sequences, Planing Numbers, and Other Recent Sequences*, talk given in Rutgers University Experimental Mathematics Seminar, Nov. 15, 2018.  
part 1: <https://vimeo.com/301216222> ; part 2: <https://vimeo.com/301219515> .  
slides: <http://sites.math.rutgers.edu/~my237/expmath/EMNov2018.pdf> .
- [S2] N.J.A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, **Sequence A318921**,  
<http://oeis.org/A318921> .

Doron Zeilberger, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.  
Email: DoronZeil at gmail dot com .

Exclusively published in the Personal Journal of Shalosh B. Ekhad and Doron Zeilberger.

Nov. 16, 2018.