

Proof of a Conjecture of Amitai Regev About Hook Multi-Sets

Doron ZEILBERGER¹

Given an integer-partition $\lambda \vdash n$ and a box (a cell) $v = [i, j] \in \lambda$ it determines the arm length $a_v (= \lambda_i - j)$, the leg length $l_v (= \lambda'_j - i)$, and the left length $f_v (= j - 1)$. Fill v with a pair of numbers in two different ways:

First Filling: Fill v with (a_v, l_v) .

Second Filling: Fill v with (a_v, f_v) .

This yields the following two multi-sets of pairs:

$$A_1(n) = \bigcup_{\lambda \vdash n} \{(a_v, l_v) | v \in \lambda\} \quad ,$$

$$A_2(n) = \bigcup_{\lambda \vdash n} \{(a_v, f_v) | v \in \lambda\} \quad .$$

The purpose of this short note is to prove Amitai Regev[R]' intriguing

Conjecture: For all non-negative integers n , $A_1(n) = A_2(n)$.

As usual, $(z)_a := (1 - z)(1 - qz) \cdots (1 - q^{a-1}z)$.

The conjecture would follow from the following two lemmas.

Lemma 1: Let $M_1(c, d)(n)$ be the number of times the pair (c, d) shows up in $A_1(n)$, then

$$\sum_{n=0}^{\infty} M_1(c, d)(n)q^n = \frac{q^{c+d+1}}{1 - q^{c+d+1}} \cdot \frac{1}{(q)_{\infty}} \quad . \tag{1}$$

Lemma 2: Let $M_2(c, d)(n)$ be the number of times the pair (c, d) shows up in $A_2(n)$, then

$$\sum_{n=0}^{\infty} M_2(c, d)(n)q^n = \frac{q^{c+d+1}}{1 - q^{c+d+1}} \cdot \frac{1}{(q)_{\infty}} \quad . \tag{2}$$

Proof of Lemma 2: $M_2(c, d)(n)$ counts the number of Ferrers diagrams of n where one of the cells that has (right) arm c and left-arm d is **marked**. Obviously it belongs to a row of length $c + d + 1$,

¹ Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA. `zeilberg at math dot rutgers dot edu` , <http://www.math.rutgers.edu/~zeilberg> . Sept. 13, 2009. Exclusive to the Personal Journal of Ekhad and Zeilberger and arxiv.org . Supported in part by the NSF.

and each such row has exactly one such cell. Hence this is the same as counting the number of Ferrers diagrams of n where one of the rows of length $c + d + 1$ is marked. We can construct such a Ferrers diagram (with any number of cells) by first drawing that row of length $c + d + 1$ (weight q^{c+d+1}) then putting **below** it an arbitrary Ferrers diagram with largest part $\leq c + d + 1$, whose generating function is $1/((1 - q)(1 - q^2) \cdots (1 - q^{c+d+1}))$, and then placing **above** the above-mentioned fixed row any Ferrers diagram whose *smallest* part is $\geq c + d + 1$, whose generating function is $1/((1 - q^{c+d+1})(1 - q^{c+d+2}) \cdots)$. Combining, we get that the generating function of such marked creatures, which is the left side of (2), is the right side of (2) \square .

Before proving Lemma 1 we have to recall certain basic facts from q -land.

Fact 1 (The q -Binomial Theorem)

$$\frac{1}{(z)_{a+1}} = \sum_{j=0}^{\infty} \frac{(q)_{a+j}}{(q)_a (q)_j} z^j \quad .$$

(This is easily proved by induction on a).

When $a = \infty$ this simplifies to

Fact 2

$$\frac{1}{(z)_{\infty}} = \sum_{j=0}^{\infty} \frac{z^j}{(q)_j} \quad .$$

Fact 3: The generating function for Ferrers diagrams bounded in an m by n rectangle is $\frac{(q)_{m+n}}{(q)_m (q)_n}$.

(Let's recall the proof of this elementary fact. Let the generating function be $F(m, n; q)$. Consider the last cell of the top row. If it is occupied, the generating function of these diagrams is $q^n F(m - 1, n)$ (remove the fully-occupied top row), if it is not, it is $F(m, n - 1)$ (delete the empty rightmost column), getting the recurrence $F(m, n; q) = q^n F(m - 1, n; q) + F(m, n - 1; q)$. Then verify that the same recurrence is satisfied by $\frac{(q)_{m+n}}{(q)_m (q)_n}$, and check the trivial initial conditions $m = 0$ and $n = 0$.)

Proof of Lemma 1: The left-side of (1) is the generating function for Ferrers diagrams where one **hook** with arm-length c and leg-length d is **marked**. Let's figure out the generating function (weight-enumerator) for all such (c, d) -hook-marked Ferrers diagrams.

Suppose the corner of that hook is at cell $(i + 1, j + 1)$ (i.e. the $(i + 1)$ -row and the $(j + 1)$ -column). Here $0 \leq i < \infty$ and $0 \leq j < \infty$. Let's look at its *anatomy*. It consists of **seven parts**.

1. Strictly **left of and above** cell $(i + 1, j + 1)$. This is a fully occupied i by j rectangle with weight q^{ij} .

2. Above the arm (of length $c + 1$). This is a fully occupied i by $c + 1$ rectangle with weight $q^{(c+1)i}$.

3. To the left of the leg (of length $d + 1$). This is a fully occupied $d + 1$ by j rectangle with weight $q^{(d+1)j}$.

4. The Ferrers diagram with $\leq i$ rows lying **above and to the right** of the arm. The generating function of this is $1/(q)_i$.

5. The Ferrers diagram with $\leq j$ columns lying **below and to the left** of the leg. The generating function of this is $1/(q)_j$.

6. The hook itself. This gives generating function q^{c+d+1} .

7. The Ferrers diagram formed **inside** the hook, i.e. lying below the arm and to the right of the leg. By Fact 3 its generating function is $\frac{(q)_{c+d}}{(q)_c(q)_d}$.

Combining, we see that the generating function for these (c, d) -hook-marked Ferrers diagrams is

$$\frac{(q)_{c+d}}{(q)_c(q)_d} \cdot q^{c+d+1} \cdot q^{ij+i(c+1)+j(d+1)} \cdot \frac{1}{(q)_i} \cdot \frac{1}{(q)_j} .$$

Summing over **all** $0 \leq i, j < \infty$, we get that the generating function on the left of (1) equals

$$\begin{aligned} & \frac{(q)_{c+d}}{(q)_c(q)_d} q^{c+d+1} \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} q^{ij+i(c+1)+j(d+1)} \frac{1}{(q)_i} \frac{1}{(q)_j} \\ &= \frac{(q)_{c+d}}{(q)_c(q)_d} q^{c+d+1} \sum_{i=0}^{\infty} \frac{1}{(q)_i} q^{i(c+1)} \sum_{j=0}^{\infty} q^{j(i+d+1)} \frac{1}{(q)_j} \\ &= \frac{(q)_{c+d}}{(q)_c(q)_d} q^{c+d+1} \sum_{i=0}^{\infty} \frac{1}{(q)_i} q^{i(c+1)} \frac{1}{(q^{d+i+1})_{\infty}} , \end{aligned}$$

by Fact 2 with $z = q^{d+i+1}$). This equals

$$\begin{aligned} & \frac{(q)_{c+d}}{(q)_c} q^{c+d+1} \sum_{i=0}^{\infty} q^{i(c+1)} \frac{1}{(q)_{\infty}} \frac{(q^{i+1})_d}{(q)_d} \\ &= \frac{q^{c+d+1}}{(q)_{\infty}} \frac{(q)_{c+d}}{(q)_c} \sum_{i=0}^{\infty} q^{i(c+1)} \frac{(q)_{i+d}}{(q)_d(q)_i} \\ &= \frac{q^{c+d+1}}{(q)_{\infty}} \frac{(q)_{c+d}}{(q)_c} \frac{1}{(q^{c+1})_{d+1}} , \end{aligned}$$

by Fact 1 with $z = q^{c+1}$. Finally, this equals

$$= \frac{q^{c+d+1}}{(q)_{\infty}} \cdot \frac{(1-q)(1-q^2) \cdots (1-q^{c+d})}{(1-q)(1-q^2) \cdots (1-q^{c+d+1})} = \frac{q^{c+d+1}}{(q)_{\infty}} \cdot \frac{1}{(1-q^{c+d+1})} \quad \square.$$

Reference

[R] A. Regev, *private communication*.