THE 1958 PEKERIS-ACCAD-WEIZAC GROUND-BREAKING COLLABORATION THAT COMPUTED GROUND STATES OF TWO-ELECTRON ATOMS (AND ITS 2010 REDUX)

CHRISTOPH KOUTSCHAN * AND DORON ZEILBERGER **

We have come a long way

In order to *appreciate* how good we as mathematicians and scientists have it today, with extremely fast hardware and lots and lots of memory, as well as with readily available high-level software, both for *numeric* and *symbolic* computation, it may be a good idea to go back to the early days of electronic computers and carefully examine, as a case study, a problem that was considered a huge challenge back then, and compare notes. We chose C.L. Pekeris' [9] 1958 seminal work[1] on the ground state energies of two-electron atoms. In particular, we will do all computations *ab initio* with today's software and hardware.

Schrödinger

Let's recall the (time-independent) Schrödinger equation for the *state function* (alias *wave function*) $\psi(x, y, z)$ of a one-electron atom with a stationary nucleus (see, for example, [8] Eq. (30-1) with $N = 1$), in *atomic units*:

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} + 2\left(E + \frac{Z}{r}\right)\right)\psi(x, y, z) \ = \ 0,$$

where Z denotes the nuclear charge, E the energy of the system, and $r = \sqrt{x^2 + y^2 + z^2}$ is the distance of the electron to the nucleus.

Schrödinger's solution of this eigenvalue problem is one of the greatest classics of modern physics, familiar to all physics (and chemistry, but unfortunately not math) students, using *separation of (dependent) variables*, and getting explicit and exact results for the *eigenvalues* (the possible energy levels E) and even for the corresponding *eigenfunctions* ψ. Since the latter (or more precisely: their squares) are interpreted as probability distributions, certain restrictions have to be imposed on ψ; in particular, the integral of ψ^2 over the whole domain must be finite. The eigenvalues then are exactly those values of E for which the Schrödinger equation admits such a solution. It turns out that these eigenfunctions are expressible in terms of the venerable special functions of mathematical physics, namely (associated) Legendre and (associated) Laguerre polynomials.

Since exactly the same predictions (about the energy levels) were already made by the "old", *ad hoc*, Bohr-Sommerfeld, quantum mechanics, the "new" wave- and matrix- quantum theories needed to predict facts that were beyond the scope of the old theory, thereby offering a crucial *confirmation*. That's why Schrödinger himself, Hylleraas, and many other physicists tried to derive the energy levels (alias eigenvalues) for two-electron atoms, whose Schrödinger equation, for the wave function $\psi = \psi(x_1, y_1, z_1, x_2, y_2, z_2)$ is

$$\left(\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial y_1^2} + \frac{\partial^2}{\partial z_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial y_2^2} + \frac{\partial^2}{\partial z_2^2} + 2\left(E + \frac{Z}{r_1} + \frac{Z}{r_2} - \frac{1}{r_{12}}\right)\right)\psi \ = \ 0,$$

Date: May 26, 2010.

* Supported by NFS-DMS 0070567 as a postdoctoral fellow, and by grant P20162 of the Austrian FWF.

** Supported in part by the United States of America National Science Foundation.

[1] available on-line from http://www.math.rutgers.edu/~zeilberg/akherim/pekeris1958.pdf (viewed June 12, 2016)