

- (25) 1. Calculate the following limits. Give a brief justification of your answers without reference to calculator computations or graphing.

(a)  $\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3}$

(b)  $\lim_{x \rightarrow 0} \frac{\sin 3x}{\tan 2x}$

(c)  $\lim_{x \rightarrow \infty} \frac{5x^4 + 7x^3 + 2x^2 + 10}{7x^4 + 5x^3 + 2x + 1}$

(d)  $\lim_{x \rightarrow \infty} \frac{e^{x/2}}{x^2}$

(e)  $\lim_{x \rightarrow 3} \frac{x^2}{\ln x}$

- (10) 2. Compute the derivative of  $\sqrt{x+3}$  **directly from the definition**.

- (25) 3. Compute the derivatives with respect to  $x$  of the following functions. Algebraic simplification of the answers need not be performed.

(a)  $\ln(x) \cos(2x)$

(b)  $\frac{e^x}{2x^3 + x}$

(c)  $\int_0^x \sec t \, dt$

(d)  $\int_0^{x^3} e^{t^2} \, dt$

(e)  $\sqrt{x^4 + 3}$

- (10) 4. Suppose that  $f$  is a function with first and second derivatives. Suppose in addition that the following values are known:  $f'(0) = 2$ ,  $f'(1) = 3$ ,  $f''(0) = 4$ , and  $f''(1) = 5$ . If  $g(x) = f(\ln(x))$ , what are  $g'(1)$  and  $g''(1)$ ?

- (15) 5. Find the following indefinite integrals:

(a)  $\int (x^3 + \frac{3}{x} + \cos x) \, dx$

(b)  $\int (2x + 1) \sec^2(x^2 + x) \, dx$

(c)  $\int \frac{6x^2 - 4}{(x^3 - 2x + 1)^3} \, dx$

(18) 6. Compute the following:

(a)  $\int_1^2 \frac{\sqrt{x} + 8}{x} dx$

(b) The area under the graph of  $y = 2 + x^2 + \sin x$  on the interval  $[0, \pi]$ .

(c)  $\int_0^\pi x^2 \sin(x^3) dx$

(10) 7. In the following,  $A$  and  $B$  are constants. Let  $f$  be the function defined by

$$f(x) = \begin{cases} x^3 + Ax & \text{if } x \leq 1 \\ Bx^2 + 2 & \text{if } x > 1 \end{cases}$$

(a) What is  $\lim_{x \rightarrow 1^-} f(x)$ ?

(b) What is  $\lim_{x \rightarrow 1^+} f(x)$ ?

(c) How must  $A$  and  $B$  be related if  $f(x)$  is continuous at  $x = 1$ ?

(d) What must the values of  $A$  and  $B$  be if  $f(x)$  is differentiable at  $x = 1$ ?

(9) 8. Use the linearization of  $\tan x$  at  $x = \pi/4$  to estimate the value of  $\tan(\pi/4 + 0.13)$ .

(10) 9. Find an equation for the tangent line to the graph of  $2x^3y^2 + x^2y^3 = 16$  at the point  $(1, 2)$ .

(10) 10. In this problem, assume that coordinates are given in feet. A point is moving along the  $x$ -axis in such a way that its acceleration at time  $t$  is  $t + \cos 2t$  ft/sec<sup>2</sup>.

(a) Suppose the velocity of the point at  $t = 0$  is 3 ft/sec. Describe the velocity of the point as a function of  $t$ .


(b) Suppose the coordinate of the point at  $t = 0$  is 10. Describe the position of the point at time  $t$ .

(8) 11. Compute the value of the Riemann sum for the function  $2^x$  on the interval  $[-1, 2]$  using the partition  $-1, 0, 1, 2$  and taking as the representative points the right endpoint of each subinterval.


(10) 12. What point on the graph of  $y = \sqrt{x}$  is closest to the point  $(3, 0)$ ?

(10) 13. Find equations for all horizontal and vertical asymptotes of the function  $\frac{4e^{-x} + 3}{7e^{-x} - 2}$ .

(10) 14. Here is the graph of a function  $f$ .



On the axes below, sketch the graph of the derivative of  $f$ .



- (10) 15. You may find it hard, but imagine you are watching a balloon in the shape of a cube being inflated. At a certain moment the volume of the balloon is 8 cubic feet and the volume is increasing at the rate of 0.3 cubic feet per minute. How fast is the surface area of the balloon increasing at that moment?
- (10) 16. In the space below, sketch the graph of a function  $f$  with the following properties:  $f(x)$  is defined and differentiable for all real numbers  $x$  except  $x = -3$  and  $x = 2$ . The graph of  $f$  has vertical asymptotes at  $x = -3$  and  $x = 2$ .

$$\lim_{x \rightarrow \infty} f(x) = -1 \quad \text{and} \quad \lim_{x \rightarrow -\infty} f(x) = 2.$$

The graph of  $f$  is concave down on the intervals  $(-\infty, -3)$  and  $(2, \infty)$  and the graph is concave up on the interval  $(-3, 2)$ .