EXPLICIT EQUATIONS OF THE FAKE PROJECTIVE PLANE $(a = 7, p = 2, \emptyset, D_3X_7)$

LEV BORISOV, MATTIE JI, AND YANXIN LI

ABSTRACT. We find explicit equations of the fake projective plane $(a = 7, p = 2, \emptyset, D_3X_7)$, which lies in the same class as the fake projective plane $(a = 7, p = 2, \emptyset, D_32_7)$ with 21 automorphisms whose equations were previously found by Borisov and Keum. The method involves finding a birational model of a common Galois cover of these two surfaces.

1. INTRODUCTION

Fake projective planes (FPPs for short) are defined as complex algebraic surfaces of general type with Hodge numbers equal to that of \mathbb{CP}^2 . The first example of an FPP was constructed by Mumford in [Mu79] using the method of 2-adic uniformization. Efforts of multiple researchers over the ensuing decades resulted in a complete classification of these surfaces as free quotients of the complex 2-ball

$$\mathcal{B}^2 = \{|z_1|^2 + |z_2|^2 < 1\} \subset \mathbb{C}^2$$

by explicit discrete subgroups of PU(2, 1). We refer the reader to [BK20] for the history of the field.

However, if one is given a fake projective plane \mathcal{B}^2/Γ with explicit generators and relations of Γ , it is still a highly nontrivial problem to write explicit equations of \mathcal{B}^2/Γ inside some projective space. The issue here is that there is no known algorithm for calculating modular forms in a way that would allow one to find polynomial equations among them. An extensive program of finding equations of fake projective planes, initiated in [BK20] and continued in [BF20, BBF20, Bo23, BL23, BJLM23], has up until now produced explicit equations for 10 out of 50 conjugate pairs of FPPs. This paper describes a method of obtaining one more pair, bringing the count up to 11. As is in all other efforts of finding FPPs, heavy computer calculations are necessary. We provide the code and the results in the supplementary materials [BJL23+]. Our primary tool is Mathematica [Math], with some computations performed in Magma [Mag] and Macaulay2 [Mac]. The fake projective plane $X = (a = 7, p = 2, \emptyset, D_3X_7)$ lies in the same class as the fake projective plane $Y = (a = 7, p = 2, \emptyset, D_32_7)$ which implies that there is a surface Z which is an étale Galois cover of both X and Y. Our construction hinges on the observation that in this particular case the Galois groups in question are the dihedral group D_{14} of 14 elements and the cyclic group C_{14} . More precisely, there exists the following diagram of maps of surfaces.

(1)
$$Y \xleftarrow{\mu} W \xleftarrow{\pi} Z \xrightarrow{\rho} X$$

where

- μ is an unramified cyclic C_2 quotient map;
- π is an unramified cyclic C_7 quotient map;
- $\mu \circ \pi$ is an unramified quotient map by the group isomorphic to D_{14} ;
- ρ is an unramified cyclic C_{14} quotient map.

The above is, of course, a direct consequence of the relations among the fundamental groups of these surfaces, which we denote by $\Gamma_X, \ldots, \Gamma_W$. Note that all of them are finite index subgroups in the maximal arithmetic subgroup $\overline{\Gamma}$ computed in [CS11+].

The paper is organized as follows. In Section 2 we study the relationship between the subgroups $\Gamma_X, \Gamma_Y, \Gamma_Z, \Gamma_W$ of $\overline{\Gamma}$ which is key to our approach. In Section 3 we describe how we obtain an embedding of W in \mathbb{CP}^{19} and a birational model of Z in \mathbb{CP}^{12} . In Section 4 we describe the method which allowed us to pass from Z to an embedding of X into \mathbb{CP}^9 . Finally, in Section 5 we make a couple of comments regarding possible further directions of this project.

Acknowledgements. This project sprang from the REU experience organized as part of the larger DIMACS REU at Rutgers University. This research was conducted while the second author was participating in the 2023 DIMACS REU program at the Rutgers University Department of Mathematics, mentored by the first author. We are grateful to Lazaros Gallos, Kristen Hendricks, and Rutgers University DIMACS for organizing the REU program and making this project possible. We also thank Sargam Mondal for useful questions and comments.

 $\mathbf{2}$

2. Galois covers

As was pointed out in the Introduction, the fake projective plane $X = (a = 7, p = 2, \emptyset, D_3X_7)$ is in the same class (in the sense of [PY07, PY10, CS10]) as the well-understood fake projective plane $Y = (a = 7, p = 2, \emptyset, D_32_7)$ whose equations were constructed in [BK20]. This means that the corresponding subgroups Γ_X and Γ_Y of PU(2, 1) are finite index subgroups in some arithmetic group $\overline{\Gamma}$. In our case, Γ_Y is normal while Γ_X is not, and both have index 21, see [CS11+].

While any two finite index subgroups contain a common normal subgroup of finite index, this index can in general be quite large. Fortunately, in our case, we found a subgroup $\Gamma_Z \subset \overline{\Gamma}$ which is a normal subgroup of index 14 in both Γ_X and Γ_Y , so that $Z = \mathcal{B}^2/\Gamma_Z$ is an unramified Galois cover of both X and Y.

Proposition 2.1. The commutator subgroup $\Gamma_Z = (\Gamma_X)'$ is a normal subgroup of $\overline{\Gamma}$. Moreover, Γ_Z is contained in both Γ_X and Γ_Y . The quotient group $\overline{\Gamma}/\Gamma_Z$ has order 294 and is isomorphic to a certain semidirect product $(D_{14} \times C_7) \rtimes C_3$ with generators (t_1, t_2, t_3, t_4) and relations

$$1 = t_1^2 = t_2^7 = (t_1 t_2)^2 = t_3^7, \ t_1 t_3 = t_3 t_1, t_2 t_3 = t_3 t_2,$$

$$1 = t_4^3, \ t_4 t_1 t_4^{-1} = t_1, \ t_4 t_2 t_4^{-1} = t_2^4, \ t_4 t_3 t_4^{-1} = t_3^2.$$

Here D_{14} denotes the dihedral group of 14 elements generated by t_1 and t_2 and the action of C_3 is seen in the above relations.

Proof. The group $\overline{\Gamma}$ is computed in [CS11+, bargammapresentations.txt] as the finitely presented group with generators z and b and relations

$$\begin{array}{c} z^7, (b^{-2}z)^3, (b^2z^{-2}b^2z^2)^3, (b^2z^{-2}b^2z^4)^3, b^3z^{-2}b^{-1}z^2b^{-2}z, \\ b^3zb^3z^3bz^2b^{-1}z^{-1}, b^3z^2b^2z^{-2}b^{-1}z^{-1}b^{-3}zb^{-1}z^{-1}. \end{array}$$

The subgroups Γ_X and Γ_Y are generated by

$$b^3, zb^3z, bz^2b^{-1}z$$

and

$$b^3, (zbz^{-1})^3, bzb^2z^{-2}, zbz^3b^{-1}$$

respectively.

Magma [Mag] readily computes [BJL23+, Section2.txt] that $\Gamma_Z = (\Gamma_X)'$ is a normal subgroup of Γ_Y and $\overline{\Gamma}$. After that, with some trial

and error, we found generators of $\overline{\Gamma}/\Gamma_Z$ (images of)

$$t_1 = b^3, t_2 = bzb^2 z^{-2}b^3, t_3 = bz^2 b^{-1}z, t_4 = b^4$$

which satisfy the relations of the proposition. See [BJL23+, Section2.txt] for the Magma verification.

From now on we will tacitly identify the quotient $\overline{\Gamma}/\Gamma_Z$ with the above semidirect product.

Proposition 2.2. The quotients Γ_X/Γ_Z and Γ_Y/Γ_Z are given by $(C_2 \times C_7) \rtimes \{1\}$ and $(D_{14} \times \{1\}) \rtimes \{1\}$ respectively, for an appropriate choice of $C_2 \subset D_{14}$.

Proof. We verify in [BJL23+, Section2.txt] that the quotients Γ_X/Γ_Z and Γ_Y/Γ_Z are generated by $\{t_1, t_3\}$ and $\{t_1, t_2\}$ respectively.

Corollary 2.3. The surfaces X and Y are given by $X = Z/C_{14}$ and $Y = Z/D_{14}$.

Remark 2.4. The elements t_3 and t_4 generate the quotient group $\overline{\Gamma}/\Gamma_Y$, isomorphic to $C_7 \rtimes C_3$. It is the automorphism group of Y and it acts on the homogeneous coordinates of the embedding of Y into \mathbb{CP}^9 by scaling and/or permuting the coordinates, see [BK20].

We will also be interested in the intermediate surface $W = Z/C_7$ with the action of C_7 coming from the normal subgroup of D_{14} in $\overline{\Gamma}/\Gamma_Z$. The corresponding group Γ_W is an index two subgroup of Γ_Y generated by Γ_Z and t_2 . The surface W is the unramified double cover of Y which corresponds to the unique nontrivial automorphism-invariant 2-torsion line bundle on Y, see [BK20, BJLM23]. Indeed, both Γ_W and Γ_Y are normal in $\overline{\Gamma}$ (see [BJL23+, Section2.txt]) and thus the index two subgroup $\Gamma_W \subset \Gamma_Y$ is preserved by the conjugation action of $\overline{\Gamma}/\Gamma_Y$.

In what follows we will need the following.

Proposition 2.5. The Hodge numbers of Z are $h^{1,0}(Z) = 0$, $h^{1,1}(Z) = 14$ and $h^{2,0}(Z) = 13$.

Proof. The Euler characteristics of \mathcal{O}_Z is 14, since $Z \to Y$ is unramified. We used the Magma file [BJL23+, Section2.txt] to check that $h^{1,0}(Z) = 0$ since the abelianization of Z is finite. This implies that $h^{2,0}(Z) = 13$, and $h^{1,1}(Z) = 14$ follows from $\chi_{top}(Z) = 14\chi_{top}(Y) = 42$. Moreover, we can identify the representation of $\overline{\Gamma}/\Gamma_Z$ on the 13dimensional space $H^0(Z, K_Z)$. Let us first discuss the action of $D_{14} \times C_7$. Recall that D_{14} has two one-dimensional representations, which we will call V_+ and V_- coming from $C_2 = D_{14}/C_7$ and three twodimensional representations which we will call V_1, V_2 and V_4 . Specifically, the generator of $C_7 \subset D_{14}$ has eigenvalues $\zeta_7^{\pm i}$ on V_i , and there is a basis of C_7 eigenvectors of V_i on which $C_2 \subset D_{14}$ acts as a permutation. Irreducible representations of $D_{14} \times C_7$ are then tensor products of the above representations of D_{14} and a one-dimensional representation of C_7 , and we denote them accordingly. For example, $V_{2,4}$ stands for the tensor product of V_2 with the representation of C_7 that sends its generator to ζ_7^4 . It has a basis $(r_{2,4}, r_{-2,4})$ where subscripts denote the weights modulo 7 with respect to the action of $C_7 \times C_7$.

Proposition 2.6. As a representation of $D_{14} \times C_7$, after a choice of a generator of $\{1\} \times C_7$, the space $H^0(Z, K_Z)$ is isomorphic to a direct sum of representations

 $V_{-,0} \oplus V_{1,1} \oplus V_{4,2} \oplus V_{2,4} \oplus V_{1,a} \oplus V_{4,2a} \oplus V_{2,4a}$

for some $a \in \{3, 5, 6\} \mod 7$.

Proof. Since the actions of D_{14} and C_{14} have no fixed points and $H^1(Z, K_Z) = 0$, the holomorphic Lefschetz fixed point formula implies that $H^0(Z, K_Z) \oplus H^2(Z, K_Z)$ is a regular representation for both of these groups. Since $H^2(Z, K_Z)$ is a trivial one-dimensional representation, we see that $H^0(Z, K_Z)$ is isomorphic to

 $V_{-,0} \oplus V_{1,a_1} \oplus V_{1,a_2} \oplus V_{2,a_3} \oplus V_{2,a_4} \oplus V_{4,a_5} \oplus V_{4,a_6}$

with (a_1, \ldots, a_6) a permutation of $(1, \ldots 6)$.

By picking a generator of $\{1\} \times C_7$, we can assume without loss of generality that the first two V-s are $V_{1,1}$ and $V_{1,a}$ for some $a \notin \{0,1\} \mod 7$. Furthermore, we have the action of the group C_3 which conjugates $V_{a,b}$ into $V_{4a,2b}$ because of Proposition 2.1. This uniquely determines the rest of the V-s. It remains to observe that a can not be 2 or 4 either because it would contradict the above permutation property.

Remark 2.7. Since we can choose a generator of $\{1\} \times C_7$ so that $V_{1,a}$ becomes $V_{1,1}$ and $V_{1,1}$ becomes $V_{1,a^{-1}}$, we can further reduce to a = 3 or a = 6 cases. We will later see that a = 3 case works. In principle, we could have used more extensive Magma calculations and the holomorphic Lefschetz formula to fully specify the representation of $\operatorname{Aut}(Z)$ on

 $H^0(Z, K_Z)$, but we find it easier to use this approach, which will be enough for our purposes.

Corollary 2.8. There is a basis

 $(r_{0,0}, r_{1,1}, r_{-1,1}, r_{4,2}, r_{-4,2}, r_{2,4}, r_{-2,4}, r_{1,a}, r_{-1,a}, r_{4,2a}, r_{-4,2a}, r_{2,4a}, r_{-2,4a})$ of $H^0(Z, K_Z)$, indexed by subscripts in $(C_7)^2$, with the following action of the automorphism group of Z:

- the generator of $C_7 \times \{1\}$ sends $r_{i,j}$ to $\zeta_7^i r_{i,j}$;
- the generator of $\{1\} \times C_7$ sends $r_{i,j}$ to $\zeta_7^j r_{i,j}$;
- the generator of $C_2 \subset D_{14}$ sends $r_{0,0}$ to $(-r_{0,0})$ and interchanges $r_{i,j}$ and $r_{-i,j}$;
- the generator of C_3 sends $r_{i,j}$ to $r_{4i,2j}$.

Proof. Follows immediately from Proposition 2.6.

3. Finding explicit equations of Z

In this section, we describe the rather lengthy process that allowed us to construct a birational model of Z as a repeated cyclic cover of Y.

In the first step, we constructed the double cover W of Y as a surface in its bicanonical embedding into \mathbb{CP}^{19} , cut out by 100 quadratic equations. While it was already accessible via methods of [BK20], we used a different approach. We note that $W \to Y$ corresponds to the 2-torsion divisor D on Y and $H^0(W, 2K_W)$ can be naturally identified with $H^0(Y, 2K_Y) \oplus H^0(Y, 2K_Y + D)$. The action of Aut(Y) extends to $H^0(Y, 2K_Y + D)$ and there is exactly one invariant section s on it, which we find as follows.

Recall that there is a unique ample divisor class H on Y such that $3H = K_Y$. In [BJLM23], we studied linear systems |nH+T| for small n and torsion divisors T. Specifically, |4H + D| has three C_7 -equivariant sections that are permuted by C_3 , and we know explicit equations of these curves on Y. This allows us to find the curve $\{s = 0\}$ on Y by looking for an automorphism-invariant section of $6K_Y = 18H$ that vanishes on the three aforementioned curves in |4H + D| linear system and using

$$6H + D = 18H - 3(4H + D).$$

Once we have found this curve $\{s = 0\}$, we are able to find an Aut(Y)invariant section of 12H = 2(6H + D) which vanishes twice on it, i.e.

6

it is equal to $s^{\otimes 2}$, up to scaling, which we denote by U_{10} . We can then use s to embed sections of $2K_Y + D$ into $H^0(Y, 4K_Y)$. Note that the map of 55-dimensional spaces $\operatorname{Sym}^2(H^0(Y, 2K_Y)) \to H^0(Y, 4K_Y)$ is an isomorphism, and we can view sections of $4K_Y$ as quadratic polynomials in the defining variables U_0, \ldots, U_9 of Y in [BK20]. The image of $\otimes s : H^0(2K_Y + D) \to H^0(4K_Y)$ is the ten-dimensional space of quadratic polynomials in U_0, \ldots, U_9 which are zero on $\{s = 0\}$. We pick a basis of it with a nice action of $\operatorname{Aut}(Y)$ (the C_7 acts by scaling the basis elements and C_3 acts by permuting them) and denote it by U_{10}, \ldots, U_{19} .

We constructed the basis (P_0, \ldots, P_{19}) of $H^0(W, 2K_W)$ by looking at $P_i = U_i$ for $i = 0, \ldots, 9$ and $P_i = \frac{U_i}{\sqrt{U_{10}}}$ for $i = 10, \ldots, 19$. We then picked random points on Y to find quadratic relations on P_i . Note that the covering involution of $W \to Y$ acts by (+1) on P_0, \ldots, P_9 and by (-1) on P_{10}, \ldots, P_{19} . We also had to be careful with the scaling of U_{10} to ensure that our model of W is defined over the field $\mathbb{Q}(\sqrt{-7})$. More precisely, we used

$$U_{10} = \frac{(49 + 13i\sqrt{7})}{56} \Big(\frac{1}{64} (-17 - 7i\sqrt{7})U_0^2 + U_1U_4 + U_2U_5 + U_3U_6 + \frac{1}{8} (-1 + i\sqrt{7})U_1U_7 + \frac{1}{8} (-1 + i\sqrt{7})U_2U_8 + \frac{1}{8} (-1 + i\sqrt{7})U_3U_9 \Big),$$

see [BJL23+, Section3.nb].

Claim 3.1. The double cover W of Y is cut out by 100 quadratic equations from [BJL23+, W_equations.txt].

Remark 3.2. We did not try to do all of the formal checks on W, as the number of variables is likely too high, although we did check its Hilbert polynomial in [BJL23+, W_check_Hilbert.txt].

Remark 3.3. We choose to make a distinction between statements that are verified by hand or by computer using symbolic manipulation, as opposed to the statements that are derived by picking some random points on the varieties in question, as is often the case in our Mathematica calculations. We call the former propositions, lemmas, et cetera, and use the term "claim" to indicate the latter.

The next step (passing from W to Z) is arguably the most technical in the entire process. It is similar to the method of [BF20]. Let $r_{1,1}, r_{-1,1}, r_{1,a}, r_{-1,a}$ be sections of $H^0(Z, K_Z)$ as in Corollary 2.8. Consider

$$s_1 = r_{1,1}r_{-1,a}, s_2 = r_{-1,1}r_{1,a}, s_3 = r_{1,1}r_{-1,1}, s_4 = r_{1,a}r_{-1,a}.$$

These s_i are invariant with respect to $C_7 \times \{1\}$ and therefore are pullbacks of sections of $H^0(W, 2K_W)$. They must satisfy a number of restrictions.

- There holds $s_1s_2 = s_3s_4$.
- Sections s_1 and s_2 have weight (a+1) with respect to the $\{1\} \times C_7$ action on W.
- Sections s_3 and s_4 have weight 2 and 2a respectively under the above action.
- The covering involution of $W \to Y$ switches s_1 and s_2 and preserves s_3 and s_4 .

This information turns out to be sufficient to reconstruct s_i , up to scaling.

Claim 3.4. For the choice of a = 3 we can pick

$$s_{1} = -P_{6} - \frac{1}{4}P_{9} - \frac{1}{4}(1 + i\sqrt{7})P_{16} + P_{19}, \ s_{2} = -P_{6} - \frac{1}{4}P_{9} + \frac{1}{4}(1 + i\sqrt{7})P_{16} - P_{19},$$

$$s_{3} = P_{5}, \ s_{4} = \frac{1}{8}(5 + i\sqrt{7})P_{1}.$$

Here P_i are the coordinates on \mathbb{CP}^{19} in which W is embedded. There are no solutions for a = 6.

Remark 3.5. If $\pi: Z \to W$ is a quotient map, we have

$$\pi_*\mathcal{O}_Z \cong \bigoplus_{i=0}^0 T^{\otimes i}$$

where T is a 7-torsion line bundle on W. The above is also precisely the splitting of $\pi_* \mathcal{O}_Z$ into eigenbundles of the C_7 -action. Since $r_{i,j}$ are eigenfunctions of the $C_7 \times \{1\}$ action on K_Z , they can be viewed as sections of $\mathcal{O}_W(3H) \otimes T^{\otimes i}$ and this defines curves $\{r_{i,j} = 0\}$ in W. Note that while we have not yet really constructed $r_{i,j}$, we can indeed construct these curves. For instance, $r_{1,1} = 0$ is the set of common zeros of s_1 and s_3 on W.

Once all the s_i are found, there are multiple approaches to finding a birational model of Z. We chose an approach that manifestly preserved the C_3 action, in addition to the $D_{14} \times C_7$ action. Specifically, the field

of rational functions of Z is a degree 7 cyclic extension of the field of rational functions of W, and we can pick the basis of it to be

(2)
$$(1, \frac{r_{1,1}}{r_{0,0}}, \frac{r_{4,2}}{r_{0,0}}, \frac{r_{2,4}}{r_{0,0}}, \frac{r_{-1,1}}{r_{0,0}}, \frac{r_{-4,2}}{r_{0,0}}, \frac{r_{-2,4}}{r_{0,0}})$$

since these cover all possible first subscripts. We can then compute the multiplication table and use it to find the polynomial relations among $r_{i,j}$. We take care to keep C_3 action throughout.

For example, suppose we want to multiply $\left(\frac{r_{4,2}}{r_{0,0}}\right)\left(\frac{r_{2,4}}{r_{0,0}}\right)$. We know that the product will be in the ζ_7^6 eigenspace of $C_7 \subset D_{14}$, so we have

$$\left(\frac{r_{4,2}}{r_{0,0}}\right)\left(\frac{r_{2,4}}{r_{0,0}}\right) = F\left(\frac{r_{-1,1}}{r_{0,0}}\right)$$

for some rational function F on W. Then F has simple zeros at $\{r_{2,4} = 0\}$ and $\{r_{4,2} = 0\}$ and poles at $\{r_{0,0} = 0\}$ and $\{r_{-1,1} = 0\}$. This determines F uniquely up to scaling, and it can be found explicitly as follows. We know that the divisor of s_3 is $\{r_{-1,1} = 0\} + \{r_{1,1} = 0\}$ and the divisor of P_{10} is $2\{r_{0,0} = 0\}$. Therefore, the divisor of Fs_3P_{10} is

$$\{r_{2,4} = 0\} + \{r_{4,2} = 0\} + \{r_{1,1} = 0\} + \{r_{0,0} = 0\}$$

which means that Fs_3P_{10} is a global section of $2K_W$. Such sections are given by quadratic polynomials in P_0, \ldots, P_{19} (we make sure to look at a C_3 -invariant subset of degree two monomials that span the complement to the space of 100 relations of W) and it is a simple matter to put conditions of vanishing on random points of the above curves to determine F. We then use C_3 action to populate the multiplication table appropriately.

Similarly, if we want to find the square of, for example, $r_{2,4}$, we see that

$$\left(\frac{r_{2,4}}{r_{0,0}}\right)^2 = F\left(\frac{r_{4,2}}{r_{0,0}}\right).$$

For the $s'_3 = r_{4,2}r_{-4,2}$ (a C_3 -translate of s_3) we see that the divisor of Fs'_3P_{10} is

$$2\{r_{2,4} = 0\} + \{r_{-4,2} = 0\} + \{r_{0,0} = 0\}.$$

We observe that F = 0 is singular at $x \in \{r_{2,4} = 0\}$ if and only if the gradient of F vanishes on the tangent space of x in W, which again allows us to find F up to scaling.

Once the multiplication table is populated, up to scaling, the associativity property is used to fix scalings (up to some natural ambiguity coming from C_3 -invariant scaling of $r_{i,j}$). We can then compute (recall that a = 3)

 $\frac{r_{1,3}}{r_{0,0}}, \frac{r_{4,6}}{r_{0,0}}, \frac{r_{2,5}}{r_{0,0}}, \frac{r_{-1,3}}{r_{0,0}}, \frac{r_{-4,6}}{r_{0,0}}, \frac{r_{-2,5}}{r_{0,0}}$

in terms of the basis (2) and use multiplication table to compute the relations among the 13 basis elements $r_{i,j}$ of $H^0(Z, K_Z)$. While we do not get an embedding, we still get a birational model of Z in \mathbb{CP}^{12} . The details can be found in [BJL23+, Section3.nb].

Remark 3.6. We used the notation

$$(Z_0, \dots, Z_{12}) = (r_{0,0}, r_{1,1}, r_{4,2}, r_{2,4}, r_{-1,1}, r_{-4,2}, r_{-2,4}, r_{1,3}, r_{4,6}, r_{2,5}, r_{-1,3}, r_{-4,6}, r_{-2,5})$$

for the homogeneous coordinates on the birational model of Z which hopefully does not cause confusion with Z itself. It is important that C_3 symmetry is maintained throughout the computation. We also have control over the action of $C_2 \subset D_{14}$, which negates $r_{0,0}$ and permutes the rest of $r_{i,j}$. Specifically, the action of generators g_3 and g_2 of the respective cyclic groups is given by

$$g_3(Z_0, \dots, Z_{12}) = (Z_0, Z_2, Z_3, Z_1, Z_5, Z_6, Z_4, Z_8, Z_9, Z_7, Z_{11}, Z_{12}, Z_{10}),$$

$$g_2(Z_0, \dots, Z_{12}) = (-Z_0, Z_4, Z_5, Z_6, Z_1, Z_2, Z_3, Z_{10}, Z_{11}, Z_{12}, Z_7, Z_8, Z_9).$$

4. Finding explicit equations of X.

In this section, we explain how we obtained equations of X in its bicanonical embedding.

Since X is obtained from Z by taking C_{14} quotient (by the $C_2 \subset D_{14}$ and $\{1\} \times C_7$) in principle one can take the C_{14} -invariant global sections of $2K_Z$ and look for polynomial equations among them. The only difficulty that we had to overcome is the fact that $\text{Sym}^2(H^0(Z, K_Z)) \rightarrow$ $H^0(Z, 2K_Z)$ is not surjective, and we can only get a dimension 7 subspace of the dimension 10 space $H^0(W, 2K_W) \cong H^0(Z, 2K_Z)^{C_{14}}$ this way. Specifically, these are given by linear combinations of

$$Z_0^2, Z_{12}Z_5 + Z_2Z_9, Z_{10}Z_6 + Z_3Z_7, Z_{11}Z_4 + Z_1Z_8, Z_{10}Z_3 + Z_6Z_7, Z_1Z_{11} + Z_4Z_8, Z_{12}Z_2 + Z_5Z_9.$$

We resolved this issue in a fairly straightforward way. Multiplication by Z_0 sends $H^0(Z, 2K_Z)^{C_{14}}$ into the space of C_7 -invariant, C_2 -antiinvariant sections of $H^0(Z, 3Z)$ which are zero on $\{r_{0,0} = 0\} \subset Z$. While some of these are simply products of Z_0 with a C_{14} -invariant quadratic polynomials above, we got three additional basis elements, specifically the C_3 orbit of

$$Z_{1}Z_{10}^{2} - \frac{1}{4}(-7 + 3i\sqrt{7})Z_{10}Z_{2}Z_{5} - 2Z_{1}Z_{3}Z_{5} + \frac{1}{2}(1 - i\sqrt{7})Z_{12}^{2}Z_{6}$$

+ $2Z_{2}Z_{4}Z_{6} - Z_{11}Z_{6}^{2} - Z_{1}Z_{10}Z_{7} + Z_{10}Z_{4}Z_{7} + \frac{1}{4}(-7 + 3i\sqrt{7})Z_{2}Z_{5}Z_{7}$
- $Z_{4}Z_{7}^{2} + Z_{3}^{2}Z_{8} - \frac{1}{2}(1 - i\sqrt{7})Z_{3}Z_{9}^{2},$

and then computed the 84 cubic equations on them.

The final verifications of the smoothness of the resulting surface and of it being the fake projective plane in bicanonical embedding were performed in [BJL23+, FPP_M2.txt, FPP_smoothness_Hodge.txt] using Magma [Mag] and Macaulay2 [Mac]. To lower the runtime of our verification codes, we reduced the equations of X modulo p = 37 which contains a square root of -7 and a non-trivial cubic root of 1. This allowed us to apply a change of coordinates on the equations of X so that the C_3 action is diagonalized. This process eliminates roughly 2/3 of the monomials used in the original equations.

5. Further directions

Techniques of this paper might allow for the computation of explicit equations of some other fake projective planes which are commensurable to the ones with known equations. We leave this for further REU projects, as this is a suitable training ground for young algebraic geometers. The sticky point is whether we can find a common normal subgroup Γ_Z in the two groups Γ_X and Γ_Y in question that has a solvable quotient Γ_Y/Γ_Z . It is considerably more difficult to construct Galois covers which are not repeated cyclic covers.

Finally, just as a teaser to the reader, we would like to point out that we only know how to match *complex conjugate pairs* of fake projective planes with their explicit embeddings, up to complex conjugation. For example, we are not aware of any method that would allow one to precisely say which choice of $\sqrt{-7}$ in our equations of X corresponds to the specific choice of the generators of Γ_X in [CS11+].

References

[Bo23] L. Borisov, On equations of fake projective planes with automorphism group of order 21, Épijournal Géom. Algébrique 7 (2023), Art. 17, https://arxiv.org/pdf/2109.02070.pdf.

- [BBF20] L. Borisov, Anders Buch and E. Fatighenti, A journey from the octonionic ℙ² to a fake ℙ², Proc. Amer. Math. Soc.150 (2022), no.4, 1467– 1475.
- [BF20] L. Borisov and E. Fatighenti, New explicit constructions of surfaces of general type, preprint arXiv:2004.02637.
- [BJL23+]Borisov, L. М. Ji and Υ. Li, Supplematerials for this article. mentary link athttps://github.com/mattie-ji/FPP-a7-p2-emptyset-D3-X7 and http://www.math.rutgers.edu/~borisov/FPP-a=7-p=2-emptyset-D3-X7.
- [BJLM23] L. Borisov, M. Ji, Y. Li and S. Mondal, On the Geometry of a Fake Projective Plane with 21 Automorphisms, preprint arXiv:2308.10429.
- [BK20] L. Borisov and J. Keum, *Explicit equations of a fake projective plane*, Duke Math. J. 169 (2020), no. 6, 1135–1162.
- [BL23] L. Borisov and Z. Lihn, Realizing a Fake Projective Plane as a Degree 25 Surface in \mathbb{P}^5 , preprint arXiv:2301.09155.
- [CS10] Donald I. Cartwright and Tim Steger, *Enumeration of the 50 fake projective planes* Comptes Rendus Mathematique, 348(1):11–13, 2010.
- [CS11+] Donald I. Cartwright and Tim Steger, http://www.maths.usyd.edu.au/u/donaldc/fakeprojectiveplanes/
- [Mac] D. Grayson and M. Stillman, Macaulay2, a software system for research in algebraic geometry, https://www.macaulay2.com.
- [Mag] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, J. Symbolic Comput., 24 (1997), 235–265; v2.23-2.
- [Math] Mathematica 12.0.0, Wolfram Research, Inc., 2019.
- [Mu79] D. Mumford, An algebraic surface with K ample, $(K^2) = 9$, $p_g = q = 0$, American J. of Math., 101 (1): 233–244.
- [PY07] G. Prasad and S.-K. Yeung, Fake projective planes, Inventiones Math., 168 (2): 321–370.
- [PY10] G. Prasad and S.-K. Yeung, Addendum to "Fake projective planes", Inventiones Math., 182 (1): 213–227.

HILL CENTER DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NJ 08854

Email address: borisov@math.rutgers.edu

URL: https://sites.math.rutgers.edu/~borisov

BROWN UNIVERSITY, DEPARTMENT OF MATHEMATICS, BOX 1917, 151 THAYER STREET, PROVIDENCE, RI 02912, USA

Email address: mattie_ji@brown.edu

HILL CENTER DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NJ 08854

Email address: yl1203@scarletmail.rutgers.edu