

Symmetry in Nonlinear PDE's

Haïm Brezis

The question of symmetry in nonlinear partial differential equations has been the subject of intensive investigations over the past 25 years. The general theme is the following. Suppose the domain Ω , as well as the boundary condition on $\partial\Omega$, has some symmetry, for example radial symmetry, axial symmetry or symmetry with respect to some hyperplane. Do solutions of nonlinear partial differential equations in Ω inherit these symmetries? In a related direction, one may consider overdetermined problems on a general domain Ω , for example the solution of a second-order PDE satisfying both a constant Dirichlet and a constant Neumann condition on $\partial\Omega$. Does this imply that the domain Ω is a ball or the complement of a ball?

Remarkable progress has been achieved through the work of Louis Nirenberg and his collaborators, especially on the first question. I will review some of their basic results. They are concerned with *positive* solutions of a *single* PDE. Related questions may be asked for *systems*. Some suggestive partial results have been obtained but the general situation is still far from satisfactory. I will describe some outstanding open problems.

1. Symmetry via Moving Planes

The main result in the celebrated paper by B. Gidas, W. M. Ni, and L. Nirenberg [GNN1] from 1979 is the following:

THEOREM 1. *Let $\Omega = B$ be the open unit ball in \mathbb{R}^n . Assume $u \in C^2(\bar{\Omega})$ satisfies*

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

where f is C^1 . Then u is radially symmetric and the radial derivative $u'(r)$ is negative for $0 < r < 1$.

As we will see the method of proof relies on the maximum principle used in conjunction with the method of moving planes due to A. D. Alexandroff [Al]. This type of argument had been initiated by J. Serrin [Se] in 1972 in proving the radial symmetry of the *domain* for overdetermined problems. The paper of B. Gidas,

W. M. Ni, and L. Nirenberg [GNN1] has become enormously popular for several reasons:

- (a) It established radial symmetry of the solution for a large class of problems. It became an incentive for investigating symmetry in numerous other situations.
- (b) The method is very flexible.

In fact, it has been adapted with success to a large variety of questions arising in concrete problems. For example, C. J. Amick and L. E. Fraenkel [AF] have used it in connection with vortex rings. W. Craig and P. Sternberg [CS] have used it to settle an open problem on water waves. H. Berestycki and L. Nirenberg [BN2] have used it in connection with problems arising in combustion.

A beautiful generalization of Theorem 1 is the following

THEOREM 2 ([BN1]). *Let Ω be a general bounded convex set in \mathbb{R}^n which is symmetric about some hyperplane, say $x_1 = 0$. Assume $u \in C^2(\Omega) \cap C(\bar{\Omega})$ satisfies (1) with f locally Lipschitz. Then u is symmetric with respect to x_1 and $\frac{\partial u}{\partial x_1} < 0$ for $0 < x_1$ in Ω .*

Theorem 2 was originally proved in [GNN1] under additional assumptions, for example $\partial\Omega$ had to be of class C^2 ; in particular, the simple case of a cube could not be handled. These restrictions were lifted by H. Berestycki and L. Nirenberg [BN1] who also gave a very elegant proof. I cannot resist the pleasure of describing their argument which deserves to become part of the classical literature.

The proof uses Stampacchia's version of the maximum principle. The standard form of the maximum principle asserts that if a function w satisfies

$$(2) \quad -\Delta w + c(x)w \leq 0 \quad \text{in } \omega,$$

$$(3) \quad w \leq 0 \quad \text{on } \partial\omega,$$

with

$$(4) \quad c(x) \geq 0 \quad \text{in } \omega,$$

then

$$w \leq 0 \quad \text{in } \omega.$$

In Stampacchia's form assumption (4) is weakened. One merely assumes that $c^- = \max(-c, 0)$ is small in some appropriate L^p norm. Suppose, for simplicity, that $n \geq 3$ and let S_n be the best Sobolev constant in \mathbb{R}^n , i.e.,

$$S_n = \inf_{\varphi \in H_0^1(\omega)} \frac{\|\nabla\varphi\|_2^2}{\|\varphi\|_{2n/(n-2)}^2}.$$

This number, which is independent of ω (and depends only on n), can be computed explicitly (see, e.g., [Au]).

LEMMA 1. *Assume w satisfies (2)–(3) with*

$$(5) \quad \|c^-\|_{n/2} < S_n.$$

Then $w \leq 0$ in ω .

In particular (5) always holds in small domains provided the sup-norm of c^- is bounded. For example the maximum principle is valid whenever

$$(6) \quad \|c^-\|_\infty |\omega|^{2/n} < S_n.$$

Another formulation suggested by S. R. S. Varadhan may be found in [BN1].

PROOF OF LEMMA 1. Multiplying (2) by $w^+ = \max(w, 0)$ and integrating by parts yields

$$\int |\nabla w^+|^2 + \int c^+(w^+)^2 - \int c^-(w^+)^2 \leq 0.$$

Thus

$$S_n \|w^+\|_{2n/(n-2)}^2 \leq \int c^-(w^+)^2 \leq \|c^-\|_{n/2} \|w^+\|_{2n/(n-2)}^2.$$

Applying (5) we find that $w^+ = 0$, i.e., $w \leq 0$ in ω . □

PROOF OF THEOREM 2. Write $x = (x_1, y)$ with $y = (x_2, x_3, \dots, x_n)$ and set

$$a = \max\{x_1; (x_1, y) \in \Omega\}.$$

We will prove that

$$(7) \quad u(x_1, y) < u(x'_1, y) \quad \forall x = (x_1, y) \in \Omega \text{ with } x_1 > 0 \text{ and } \forall x'_1 \text{ with } |x'_1| < x_1.$$

Inequality (7) yields

$$(8) \quad u(x_1, y) \leq u(-x_1, y),$$

and applying (8) to $\tilde{u}(x_1, y) = u(-x_1, y)$, which is also a solution of (1), one finds that $u(-x_1, y) = u(x_1, y)$, i.e., u is symmetric with respect to x_1 . The fact that $\frac{\partial u}{\partial x_1} < 0$ for $0 < x_1$ in Ω is an easy consequence of (7). □

For $0 < \lambda < a$, set

$$\Sigma(\lambda) = \{x = (x_1, y) \in \Omega; x_1 > \lambda\}$$

and

$$w^\lambda(x) = u(2\lambda - x_1, y) - u(x_1, y) \quad \text{for } x \in \Sigma(\lambda).$$

Note that w^λ is well defined on $\Sigma(\lambda)$ since Ω is convex and symmetric about the hyperplane $x_1 = 0$.

Inequality (7), to be proved, is equivalent to

$$(9) \quad w^\lambda(x) > 0 \quad \forall x \in \Sigma(\lambda), \quad \forall \lambda \in (0, a).$$

The function w^λ satisfies

$$-\Delta w^\lambda + c^\lambda(x)w^\lambda = 0 \quad \text{in } \Sigma(\lambda)$$

where

$$c^\lambda(x) = \begin{cases} \frac{f(u(x_1, y)) - f(u(2\lambda - x_1, y))}{w^\lambda(x)} & \text{if } w^\lambda(x) \neq 0 \\ 0 & \text{if } w^\lambda(x) = 0. \end{cases}$$

Clearly $\|c^\lambda\|_\infty \leq L$ where L is the Lipschitz constant of f on the interval $[-\|u\|_\infty, \|u\|_\infty]$. Moreover

$$w^\lambda \geq 0 \quad \text{on } \partial\Sigma(\lambda),$$

$$w^\lambda \not\equiv 0 \quad \text{on } \partial\Sigma(\lambda).$$

For λ near a , $\Sigma(\lambda)$ has small measure and we deduce from Lemma 1 that $w^\lambda \geq 0$ in $\Sigma(\lambda)$.

Let

$$\Lambda = \{\lambda \in (0, a); w^\lambda \geq 0 \text{ in } \Sigma(\lambda)\},$$

so that Λ is not empty. Clearly Λ is closed in $(0, a)$. We claim that Λ is open.

Fix some $\lambda \in \Lambda$. By the strong maximum principle applied to w^λ in $\Sigma(\lambda)$ we see that

$$w^\lambda > 0 \quad \text{in } \Sigma(\Lambda).$$

Let K be any (smooth) compact set in $\Sigma(\lambda)$ such that $|\Sigma(\mu) \setminus K|$ is sufficiently small for all μ near λ . Sufficiently small refers to Lemma 1 applied in $\Sigma(\mu) \setminus K$ with $\|c\|_\infty \leq L$ (the Lipschitz constant of f).

Since

$$w^\lambda(x) \geq \delta > 0 \quad \text{in } K$$

we have, by continuity,

$$w^\mu(x) \geq 0 \quad \text{in } K$$

for all μ near λ . In particular

$$w^\mu(x) \geq 0 \quad \text{on the boundary of } \Sigma(\mu) \setminus K.$$

Applying Lemma 1 to w^μ in $\Sigma(\mu) \setminus K$ we see that $w^\mu \geq 0$ in $\Sigma(\mu) \setminus K$ and thus $w^\mu \geq 0$ in $\Sigma(\mu)$. Hence $\mu \in \Lambda$ for all μ near λ , i.e., Λ is open.

REMARK 1. The assumption that Ω is convex is essential. For example if Ω is an annulus, radial symmetry may fail. (In fact, nonradial solutions can have lower energy than the radial ones.) We have constructed in [BrN1] (see also [D]) *nonradial* positive solutions of

$$\begin{cases} -\Delta u = u^p + \lambda u & \text{in } \Omega = \text{annulus} \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Radial symmetry of solutions of (1) when Ω is all of \mathbb{R}^n (with $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$) has been originally studied in [GNN2]; important extensions may be found in [CGS], [Li], [LN], [CL], and [Z]. In particular, the radial symmetry is useful in order to give a complete description of all positive solutions of $-\Delta u = u^{(n+2)/(n-2)}$ in \mathbb{R}^n . These functions are the extremals for the Sobolev inequality $\int |\nabla \varphi|^2 \geq S \|\varphi\|_{2n/(n-2)}^2$. This classification plays an important role—after blow-up—in the study of the Yamabe problem, in particular in the work of R. Schoen [Sc].

In the case where Ω is a half-space $\Omega = \{x \in \mathbb{R}^n; x_n > 0\}$, with zero Dirichlet condition, H. Berestycki, L. A. Caffarelli, and L. Nirenberg [BCN] have established symmetry (i.e., $u = u(x_n)$) and monotonicity provided u is bounded and $f(\sup u) \leq 0$. The case of a half-space Ω with a nonlinear Neumann condition has been investigated in [CFS], [E], and [LZ] and [T]. Such results have applications to prescribed curvature problems on manifolds with boundary.

Symmetry and monotonicity in infinite cylindrical domains have been studied by H. Berestycki and L. Nirenberg [BN2] in connection with travelling front solutions arising in combustion. The case where Ω is the exterior of a ball has been considered in [AB]. The interested reader will find further variations on this theme in the expository paper [Be].

The moving plane method has also been applied to establish symmetry of solutions for some classes of *systems* of PDE's; see [Ba1], [DF], and [Tr]. In particular, for the Liouville system,

$$-\Delta u_i = \exp(\sum_{j=1}^n a_{ij} u_j) \quad \text{in } \mathbb{R}^2, \quad 1 \leq i \leq n,$$

with $u_i > 0$. M. Chipot, I. Shafrir, and G. Wolansky [CSW] have proved under mild assumptions, that each u_i is radially symmetric and decreasing about some

point x_i in \mathbb{R}^2 . (An interesting earlier approach by S. Chanillo and M. Kiessling [CK1], based solely on an isoperimetric inequality and the Pohozaev identity, led to similar conclusions under stronger assumptions.) However, the application of the moving plane method to systems has been, so far, very limited. I would like to describe next two types of systems where other techniques have been successful.

2. Questions of Symmetry for the Ginzburg-Landau System

The Ginzburg-Landau system consists of a coupled system of two equations in \mathbb{R}^2 ,

$$(10) \quad -\Delta u = u(1 - |u|^2) \quad \text{in } \mathbb{R}^2.$$

Here u takes its values in \mathbb{R}^2 and it is also convenient to view u as a complex number. Despite its simple appearance, problem (10) has a rich structure, which is not yet fully understood. It is an interesting laboratory for testing new methods.

One is concerned with solutions of (10) satisfying

$$(11) \quad |u(x)| \rightarrow 1 \quad \text{as } |x| \rightarrow \infty.$$

It is easy to construct solutions of (10)–(11) in polar coordinates, using separation of variables. Given any integer $q \in \mathbb{Z}$ the function

$$(12) \quad u = u(r, \theta) = e^{iq\theta} f(r)$$

is a solution of (10)–(11) provided the real-valued function f satisfies the ordinary differential equation

$$(13) \quad \begin{cases} -f'' - \frac{1}{r}f' + \frac{q^2}{r^2}f = f(1 - f^2) & \text{on } (0, \infty) \\ f(0) = 0 \quad \text{and} \quad f(\infty) = 1. \end{cases}$$

It is not difficult to see that for every integer q problem (13) has a unique solution f_q (see, e.g., Appendices II, III in [BBH] and also [HH]). Hence, we obtain a family of special solutions $u_q = e^{iq\theta} f_q(r)$ for $q \in \mathbb{Z}$. An outstanding open problem is whether these are the only solutions of (10)–(11):

Open Problem 1. Let u be any solution of (10)–(11). Is $u = u_q$ for some $q \in \mathbb{Z}$, modulo translation and rotation?

An unusual quantization phenomenon takes place for solutions of (10) having the property that $|u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$, fast enough so that

$$(14) \quad \int_{\mathbb{R}^2} (|u|^2 - 1)^2 < \infty.$$

REMARK 2. It is easy to show that if u is any solution of (10) satisfying (14) then $|u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$. The converse is not known:

Open Problem 2. Suppose u is a solution of (10) such that $|u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$. Does (14) hold?

THEOREM 3. ([BMR]) *Let u be a solution of (10) satisfying (14). Let $q = \text{deg}(u, \infty)$ be the degree of u at infinity, i.e., the winding number of the map*

$$x \in S^1 \mapsto \frac{u(Rx)}{|u(Rx)|} \in S^1 \quad \text{for large } R.$$

Then

$$(15) \quad \frac{1}{2\pi} \int_{\mathbb{R}^2} (|u|^2 - 1)^2 = q^2.$$

The “radial” solution $u_q = e^{iq\theta} f_q(r)$ described above satisfies (15). It is not known, for general q , whether the only solution of (10) satisfying (15) is u_q , modulo translation, rotation and complex conjugation. The answer is positive for $q = 0$ and $q = 1$. The case $q = 0$ is an easy consequence of Liouville theorem. The case $q = 1$ is a remarkable result of P. Mironescu ([M2]) described in Theorem 4 below.

SKETCH OF THE PROOF OF THEOREM 3. The main ingredient is the Pohozaev identity applied to (10). It asserts that

$$(16) \quad \int_{B_R} (|u|^2 - 1)^2 = \frac{R}{2} \int_{S_R} (|u|^2 - 1)^2 + R \int_{S_R} (|u_t|^2 - |u_n|^2).$$

Here $B_R = \{x \in \mathbb{R}^2; |x| < R\}$, $S_R = \{x \in \mathbb{R}^2; |x| = R\}$, u_t and u_n denote respectively the tangential and normal derivatives of u along S_R . Identity (16) is obtained, as usual, through the multiplication of (10) by $xu_x + yu_y = ru_r$ and integration on B_R . Using (14) one shows (see, e.g., [Sh1] and [Br]) that, as $R \rightarrow \infty$,

$$R \int_{S_R} (|u|^2 - 1)^2 \rightarrow 0 \quad \text{and} \quad R \int_{S_R} |u_n|^2 \rightarrow 0.$$

The important term in (16)—the one which “carries” the degree—is u_t . More precisely one shows that, as $R \rightarrow \infty$,

$$R \int_{S_R} |u_t|^2 \rightarrow 2\pi q^2.$$

□

Now, to the result of Mironescu:

THEOREM 4. *Let u be a solution of (10) satisfying (15) with $q = 1$. Then u has radial symmetry, i.e., $u = e^{i\theta} f_1(r)$ modulo rotation, translation, and complex conjugation.*

SKETCH OF THE PROOF OF THEOREM 4. Let $f(r) = f_1(r)$ be the unique solution of (13) corresponding to $q = 1$. Since $\deg(u, \infty) \neq 0$ the function u must have at least one zero. After translation we may assume that $u(0) = 0$. Set $v = u/f$. Using (10) and (13) it is easy to derive a PDE satisfied by v :

$$(17) \quad -\Delta v - \frac{2f'}{f} v_r - \frac{v}{r^2} = f^2 v (1 - |v|^2)$$

where v_r is the radial derivative of v , i.e., $v_r = \frac{1}{|x|} (x \cdot \nabla v)$.

Applying the Pohozaev identity to (17) yields

$$(18) \quad \int_{B_R} \left[\frac{2rf'}{f} |v_r|^2 + \frac{1}{2} (f^2 + rff') (|v|^2 - 1)^2 \right] = \int_{S_R} [\dots]$$

where $[\dots]$ is a lengthy expression involving v, v_r, f , and f' . The solution f of (13) is known to be monotone increasing (see, e.g., [HH]), so that the integrand on the left-hand side of (18) is nonnegative. A careful asymptotic analysis of $u(x)$ as $|x| \rightarrow \infty$ (see [Sh1] and [Br]) combined with Theorem 3 shows that the right-hand side in (18) tends to 0 as $r \rightarrow \infty$. Thus $v_r \equiv 0$ and $|v| \equiv 1$. Going back to (17)

we obtain $v_{\theta\theta} + v = 0$, i.e., $v = e^{i(\theta+\theta_0)}$ or $v = e^{-i(\theta+\theta_0)}$. Returning to u we find $u = e^{i\theta} f(r)$ or $u = e^{-i\theta} f(r)$, modulo a rotation. \square

REMARK 3. The Pohozaev identity seems to play a distinguished role in proving symmetry for two-dimensional problems. P. L. Lions [Lio] has given a proof of Theorem 1 in 2-d which does not make use of the moving plane method. It relies on a clever combination of the Pohozaev identity with an isoperimetric inequality. Related ideas may be found in [Ba2], [CK1], and [CK2]. Unfortunately the method seems to be restricted to two-dimensional problems. One may consider the analogue of (10) in higher dimension and, there, no symmetry result is known:

Open Problem 3. Let $u : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a solution of

$$-\Delta u = u(1 - |u|^2) \quad \text{on } \mathbb{R}^n, \quad n \geq 3$$

with $|u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$ (possibly with a “good” rate of convergence). Assume $\deg(u, \infty) = \pm 1$. Does u have the form

$$u(x) = \frac{x}{|x|} f(r)$$

(modulo translation and isometry), where $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a smooth function, such that $f(0) = 0$ and $f(\infty) = 1$?

REMARK 4. The proof of Theorem 4 provides some information for general values of q . Let u be a solution of (10) satisfying (15) with $q \geq 2$. Assume that u has only one zero (of degree q). Then $u = e^{iq\theta} f_q(r)$. This result raises an interesting variant of Problem 1:

Open Problem 4. Let u be a solution of (10) with $|u(x)| \rightarrow 1$ as $|x| \rightarrow \infty$. Can u have more than one zero?

Theorem 4 has important implications, for example:

THEOREM 5. ([M2]) *Let u be a solution of (10) which is a local minimizer of the energy*

$$E(v, \Omega) = \frac{1}{2} \int_{\Omega} |\nabla v|^2 + \frac{1}{4} \int_{\Omega} (|v|^2 - 1)^2$$

in the sense that for every bounded domain $\Omega \subset \mathbb{R}^2$,

$$E(u, \Omega) \leq E(v, \Omega), \quad \forall v \text{ such that } v = u \text{ on } \partial\Omega.$$

Then either u is a constant or $u = e^{i\theta} f_1(r)$ (modulo translation, rotation, and complex conjugation).

The proof of Theorem 5 uses Theorem 4 in conjunction with a result of E. Sandier [Sa2] and I. Shafrir [Sh1] (u a local minimizer $\Rightarrow \frac{1}{2\pi} \int_{\mathbb{R}^2} (|u|^2 - 1)^2 = 1$).

Theorem 5 is very useful in analyzing the structure of the Ginzburg-Landau vortices near the vortex core. Let Ω be a bounded domain in \mathbb{R}^2 and let $g : \partial\Omega \rightarrow S^1$ be a smooth boundary condition of degree $d > 0$. Let u_ε be a minimizer of the Ginzburg-Landau energy

$$E_\varepsilon(v) = \frac{1}{2} \int_{\Omega} |\nabla v|^2 + \frac{1}{4\varepsilon^2} \int_{\Omega} (|v|^2 - 1)^2$$

with boundary condition $v = g$ on $\partial\Omega$. One of the main results in [BBH] asserts that for ε small, u_ε has exactly d zeroes $a_\varepsilon^1, a_\varepsilon^2, \dots, a_\varepsilon^d$ and that (along a subsequence)

$$(19) \quad u_\varepsilon(z) \rightarrow u_*(z) = e^{i\psi(z)} \prod_{i=1}^d \frac{z - a^i}{|z - a^i|}$$

where $a^i = \lim_{\varepsilon \rightarrow 0} a_\varepsilon^i$ and ψ is a real-valued harmonic function in Ω . The convergence in (19) holds in $C_{\text{loc}}^k(\Omega \setminus \{a_1, a_2, \dots, a_d\})$, for every k . However, there was no information in [BBH] about the mode of convergence of u_ε to u_* near its singularities. As a consequence of Theorem 5 we now have:

THEOREM 6. ([Sh2], [M2]) *Let $U(z) = \frac{z}{|z|} f_1(z)$ where f_1 is the solution of (13) with $q = 1$. Then*

$$(20) \quad \lim_{\varepsilon \rightarrow 0} \left\| u_\varepsilon(z) - e^{i\psi(z)} \prod_{i=1}^d U\left(\frac{z - a_\varepsilon^i}{\varepsilon}\right) \right\|_{L^\infty(\Omega)} = 0.$$

The product $\prod_{i=1}^d$ in (20) denotes the product of complex numbers. Theorem 6 is derived from Theorem 5 via a blow-up argument. One shows that, as $\varepsilon \rightarrow 0$, $u_\varepsilon(\varepsilon z + a_\varepsilon^i)$ converges to a solution of (10) which is a local minimizer of the energy in the sense of Theorem 5 and we may now identify this blow-up limit as $U(z)$ (modulo a rotation).

One may ask similar questions on the disc but the situation is widely open. Consider, for example, the equation

$$(21) \quad \begin{cases} -\Delta u = au(1 - |u|^2) & \text{in } B = \text{the unit disc in } \mathbb{R}^2 \\ u(x) = x & \text{on } \partial B \end{cases}$$

where $a > 0$ is a constant and $u : B \rightarrow \mathbb{C} = \mathbb{R}^2$. It is easy to construct a “radial” solution

$$u = u(r, \theta) = e^{i\theta} f(r)$$

where f satisfies the ordinary differential equation

$$\begin{cases} -f'' - \frac{1}{r}f' + \frac{1}{r^2}f = af(1 - f^2) & \text{in } (0, 1) \\ f(0) = 0 \text{ and } f(1) = 1. \end{cases}$$

This f is uniquely determined (see [BBH] and [HH]).

Open Problem 5. Is the radial solution $e^{i\theta} f(r)$ the only solution of (21)?

If $a \leq \lambda_1$ (the first eigenvalue of $-\Delta$ on B with zero Dirichlet condition) the answer is positive since the energy functional

$$E(v) = \frac{1}{2} \int_B |\nabla v|^2 + \frac{a}{4} \int_B (|v|^2 - 1)^2$$

is strictly convex and thus (21) has a solution.

REMARK 5. The argument described in the proof of Theorem 4 is still valid provided u/f makes sense at 0, i.e., $u(0) = 0$. Thus, another formulation of Open Problem 5 is:

Open Problem 5'. Does any solution u of (21) vanish at 0?

A weaker form of Open Problem 5, which I find quite intriguing is:

Open Problem 6. Is the radial solution $u = e^{i\theta} f(r)$ a minimizer of the energy E ? More generally, if B is the unit ball in $\mathbb{R}^n, n \geq 2$ and $u : B \rightarrow \mathbb{R}^n$, is there a minimizer of E of the form $\frac{x}{|x|} f(|x|)$?

P. Mironescu [M1] (see also [LL]) has given a partial answer. He proves that the radial solution u is a local minimizer in the sense that $E(u) \leq E(v)$ for all $v \in H^1$ such that $v(x) = x$ on ∂B and $\|v - u\|_{H^1}$ is small. In the scalar case rearrangement techniques (see, e.g., [Ba1]) are often used to prove that minimizers have radial symmetry. But in the vector-valued case no such method is available (see, however, the discussion after Theorem 9 below). Therefore it would be very interesting to settle Open Problem 6.

3. Questions of Symmetry for Minimizing Harmonic Maps

Another simple nonlinear PDE system which has received much attention in recent years is the system of harmonic maps. Since we are interested in questions of symmetry we will assume that the target space is a sphere, say $S^{k-1}, k \geq 2$. The unknown u is a map from a domain $\Omega \subset \mathbb{R}^n$ with values into \mathbb{R}^k satisfying

$$(22) \quad \begin{cases} -\Delta u = u|\nabla u|^2 & \text{in } \Omega \\ |u| = 1 & \text{in } \Omega \\ u = g & \text{on } \partial\Omega \end{cases}$$

where $g : \partial\Omega \rightarrow S^{k-1}$ is a given (smooth) boundary condition. The solutions of (22) arise as critical points of the Dirichlet integral

$$E(v) = \int_{\Omega} |\nabla v|^2$$

subject to the constraint

$$v \in H_g^1(\Omega; S^{k-1}) = \{v : \Omega \rightarrow \mathbb{R}^k; \int_{\Omega} |\nabla v|^2 < \infty, |v| = 1 \text{ in } \Omega \text{ and } v = g \text{ on } \partial\Omega\}.$$

Of particular interest are minimizing harmonic maps, i.e., minimizers of E in $H_g^1(\Omega; S^{k-1})$. *Minimizing* harmonic maps seem to inherit some symmetry properties of the data. However, general harmonic maps, i.e., arbitrary (weak) solutions of (22) usually break symmetry. Here are some results.

THEOREM 7. *Let $\Omega = B$ be the unit ball in $\mathbb{R}^n, n \geq 3$. Assume $k = n$ and $g(x) = x$. Then $u(x) = x/|x|$ is a minimizing harmonic map; in fact it is the unique minimizer of E in $H_g^1(\Omega; S^{n-1})$.*

This result was originally proved by W. Jäger and H. Kaul [JK] when $n \geq 7$ (they even show that $x/|x|$ is a minimizer in $H_g^1(\Omega; S^n)$, where S^{n-1} is identified with an equator of S^n). Theorem 7 is due to H. Brezis, J. M. Coron, and E. Lieb [BCL] when $n = 3$ and to F. H. Lin [Lin] for general $n \geq 3$. The proof of F. H. Lin is especially ingenious and elegant. The restriction $n \geq 3$ is needed. When $n = 2$ the class of testing functions $H_g^1(\Omega; S^1)$ is empty; this is a consequence of the fact that there is a degree theory for maps in $H^{1/2}(S^1; S^1)$ (see [BBH] and [BrN2]).

REMARK 6. There is no hope to prove that general (i.e., nonminimizing) harmonic maps inherit the radial symmetry of the boundary condition. In fact, T.

Rivière [R] has constructed an abundance of weird solutions of (22) when Ω is the unit ball in \mathbb{R}^3 , $k = 3$ and g is any nonconstant boundary condition (in particular $g(x) = x$).

REMARK 7. F. Almgren and E. Lieb [AL] have pointed out that natural notions of symmetry may be broken, even for minimizing harmonic maps. Consider, for example in 3-d the notion of mirror symmetry through the xy plane, i.e.,

$$\begin{aligned}u_1(x, y, -z) &= u_1(x, y, z) \\u_2(x, y, -z) &= u_2(x, y, z) \\u_3(x, y, -z) &= -u_3(x, y, z).\end{aligned}$$

They have constructed an example where Ω is the unit ball in \mathbb{R}^3 , $k = 3$, the boundary condition g has mirror symmetry, but no minimizer has mirror symmetry.

When Ω is a two-dimensional domain and $k \geq 3$ there seems to be a better chance for symmetry. Here are some situations where symmetry holds.

Let Ω be the unit disc in \mathbb{R}^2 and consider maps $u : \Omega \rightarrow S^2$. We say that u has radial symmetry if it can be written in the form

$$u(x, y) = (a(r)x, a(r)y, b(r))$$

where $r = (x^2 + y^2)^{1/2}$, $a(r)$ and $b(r)$ are real-valued functions such that $a^2(r) + b^2(r) = 1$.

THEOREM 8. ([BC]) *Let Ω be the unit disc in \mathbb{R}^2 and let*

$$g(x, y) = (Rx, Ry, \sqrt{1 - R^2}) \quad \text{for } (x, y) \in \partial\Omega \text{ with } 0 < R \leq 1.$$

Then any minimizer of E in $H_g^1(\Omega; B^2)$ has radial symmetry. (In fact, there are precisely two minimizers.)

Open Problem 7. Is the same conclusion true for general (nonminimizing) solutions of (22)?

THEOREM 9. *Let $0 < \rho < 1$ and consider the annulus*

$$\Omega_\rho = \{(x, y) \in \mathbb{R}^2; \rho^2 < x^2 + y^2 < 1\}.$$

Consider the boundary condition

$$g(x, y) = \frac{1}{2}(x, y, 0) \quad \text{on } \partial\Omega_\rho.$$

Then any minimizer of E in $H_g^1(\Omega_\rho; S^2)$ has radial symmetry.

Theorem 9 was originally proved by E. Sandier [Sa1] in connection with results of F. Bethuel, H. Brezis, B. Coleman, and F. Hélein [BBCH]. A new proof was given by S. Kaniel and I. Shafrir [KS]. It relies on a very interesting symmetrization device, which could possibly be useful for other vector-valued problems. It is also quite unexpected to have radial symmetry in the annulus since there are examples of broken symmetry for the annulus even in the scalar case (see Remark 1).

References

- [AB] A. Aftalion and J. Busca, *Symétrie radiale pour des problèmes elliptiques surdéterminés posés dans des domaines extérieurs*, C. R. Acad. Sci. Paris **324** (1997), pp. 633–638.
- [Al] A. D. Alexandroff, *A characteristic property of the spheres*, Ann. Mat. Pura Appl. **58** (1962), pp. 303–354.

- [AL] F. Almgren and E. Lieb, *Singularities of energy minimizing maps from the ball to the sphere: Examples, counterexamples and bounds*, Ann. of Math. **128** (1988), pp. 483–530.
- [AF] C. J. Amick and L. E. Fraenkel, *Uniqueness of Hill's spherical vortex*, Arch. Rational Mech. Anal. **92** (1986), pp. 91–119.
- [Au] T. Aubin, *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, Springer-Verlag, New York, 1982.
- [Ba1] M. Badiale, *Monotonicity of solutions for elliptic systems on unbounded domains*, Bull. U.M.I. A **6** (1992), pp. 59–64.
- [Ba2] C. Bandle, *Isoperimetric Inequalities and Applications*, Pitman, Boston, 1980.
- [Be] H. Berestycki, *Qualitative properties of semilinear elliptic equations in unbounded domains*, to appear.
- [BCN] H. Berestycki, L. A. Caffarelli, and L. Nirenberg, *Symmetry for elliptic equations in a half space*, pp. 27–42 in *Boundary Value Problems for Partial Differential Equations and Applications*, C. Baiocchi et al., eds., Masson, Paris, 1993.
- [BN1] H. Berestycki and L. Nirenberg, *On the method of moving planes and the sliding method*, Bol. Soc. Brasil. Mat. **22** (1991), pp. 1–37.
- [BN2] H. Berestycki and L. Nirenberg, *Travelling waves in cylinders*, Ann. Inst. H. Poincaré, Anal. Non Linéaire **9** (1992), pp. 497–572.
- [BBCH] F. Bethuel, H. Brezis, B. Coleman, and F. Hélein, *Bifurcation analysis for minimizing harmonic maps describing the equilibrium of nematic phases between cylinders*, Arch. Rational Mech. Anal. **118** (1992), pp. 149–168.
- [BBH] F. Bethuel, H. Brezis, and F. Hélein, *Ginzburg-Landau Vortices*, Birkhäuser, Boston, 1994.
- [Br] H. Brezis, *Lectures on the Ginzburg-Landau Vortices*, Scuola Normale Superiore, Pisa, 1998.
- [BC] H. Brezis and J. M. Coron, *Large solutions for harmonic maps in two dimensions*, Comm. Math. Phys. **92** (1983), pp. 203–215.
- [BCL] H. Brezis, J. M. Coron, and E. Lieb, *Harmonic maps with defects*, Comm. Math. Phys. **107** (1986), pp. 649–705.
- [BMR] H. Brezis, F. Merle, and T. Rivière, *Quantization effects for $-\Delta u = u(1 - |u|^2)$ in \mathbb{R}^2* , Arch. Rational Mech. Anal. **126** (1994), pp. 123–148.
- [BrN1] H. Brezis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), pp. 437–477.
- [BrN2] H. Brezis and L. Nirenberg, *Degree theory and BMO*, Selecta Math. **1** (1995), pp. 197–263, and **2** (1996), pp. 1–60.
- [CGS] L. Caffarelli, B. Gidas, and J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), pp. 271–297.
- [CK1] S. Chanillo and M. Kiessling, *Conformally invariant systems of nonlinear PDE's of Liouville type*, Geom. Funct. Anal. **5** (1995), pp. 924–947.
- [CK2] S. Chanillo and M. Kiessling, *Curl-free Ginzburg-Landau vortices*, Nonlinear Anal., to appear.
- [CL] W. Chen and C. Li, *Classification of solutions of some nonlinear elliptic equations*, Duke Math. J. **63** (1991), pp. 615–623.
- [CFS] M. Chipot, M. Fila, and I. Shafrir, *On the solutions to some elliptic equations with nonlinear Neumann boundary conditions*, Adv. Differential Equations **1** (1996), pp. 91–110.
- [CSW] M. Chipot, I. Shafrir, and G. Wolansky, *On the solutions of Liouville systems*, to appear.
- [CS] W. Craig and P. Sternberg, *Symmetry of solitary waves*, Comm. Partial Differential Equations **13** (1988), pp. 603–633.
- [D] E. N. Dancer, *Global breaking of symmetry of positive solutions on two dimensional annuli*, Differential Integral Equations **5** (1992), pp. 903–913.
- [DF] D. G. DeFigueiredo and P. L. Felmer, *A Liouville-type theorem for elliptic systems*, Ann. Sci. Norm. Sup. Pisa **21** (1994), pp. 387–397.
- [E] J. F. Escobar, *Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate*, Comm. Pure Appl. Math. **43** (1990), pp. 857–883.
- [GNN1] B. Gidas, W. M. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), pp. 209–243.

- [GNN2] B. Gidas, W. M. Ni, and L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n* , pp. 369–402 in: Math. Anal. Appl.; Part A, Advances in Math. Suppl. Studies 7A, L. Nachbin, ed., Academic Press, New York, 1981.
- [HH] M. Hervé and R. M. Hervé, *Etude qualitative des solutions réelles d'une équation différentielle liée à l'équation de Ginzburg-Landau*, Ann. Inst. H. Poincaré, Anal. Non Linéaire **11** (1994), pp. 427–440.
- [JK] W. Jäger and H. Kaul, *Rotational symmetric harmonic maps from a ball into a sphere and the regularity problem for weak solutions of elliptic systems*, J. Reine Angew. Math. **343** (1983), pp. 146–161.
- [KS] S. Kaniel and I. Shafrir, *A new symmetrization method for vector valued maps*, C. R. Acad. Sci. Paris **315** (1992), pp. 413–416.
- [Li] C. Li, *Monotonicity and symmetry of solutions of fully nonlinear elliptic equations on unbounded domains*, Comm. Partial Differential Equations **16** (1991), pp. 585–615.
- [LN] Y. Li and W. M. Ni, *On the asymptotic behavior and radial symmetry of positive solutions of semilinear equations in \mathbb{R}^n* , Arch. Rational Mech. Anal. **118** (1992), pp. 195–222.
- [LZ] Y. Li and M. Zhu, *Uniqueness theorems through the method of moving spheres*, Duke Math. J. **80** (1995), pp. 383–417.
- [LL] E. Lieb and M. Loss, *Symmetry of the Ginzburg-Landau minimizer in a disc*, Math. Res. Letters **1** (1994), pp. 701–715.
- [Lin] F. H. Lin, *Une remarque sur l'application $x/|x|$* , C. R. Acad. Sci. Paris **305** (1987), pp. 529–531.
- [Lio] P. L. Lions, *Two geometrical properties of solutions of semilinear problems*, Appl. Anal. **12** (1981), pp. 267–272.
- [M1] P. Mironescu, *On the stability of radial solutions of the Ginzburg-Landau equation*, J. Funct. Anal. **130** (1995), pp. 334–344.
- [M2] P. Mironescu, *Les minimiseurs locaux pour l'équation de Ginzburg-Landau sont à symétrie radiale*, C. R. Acad. Sci. Paris **323** (1996), pp. 593–598.
- [R] T. Rivière, *Applications harmoniques de B^3 dans S^2 partout discontinues*, C. R. Acad. Sci. Paris **314** (1992), pp. 719–723; see also *Everywhere discontinuous harmonic maps into spheres*, Acta Math. **175** (1995), pp. 197–226.
- [Sa1] E. Sandier, *The symmetry of minimizing harmonic maps from a two-dimensional domain to the sphere*, Ann. Inst. H. Poincaré, Anal. Non Linéaire **10** (1993), pp. 549–559.
- [Sa2] E. Sandier, *Locally minimizing solutions of $-\Delta u = u(1 - |u|^2)$ in \mathbb{R}^2* , Proc. Roy. Soc. Edinburgh, to appear.
- [Sc] R. Schoen, *On the number of constant scalar curvature metrics in a conformal class*, pp. 311–320 in Differential Geometry, H. B. Lawson et al., eds., Pitman Monographs, Wiley, New York, 1991.
- [Se] J. Serrin, *A symmetry theorem in potential theory*, Arch. Rational Mech. Anal. **43** (1971), pp. 304–318.
- [Sh1] I. Shafrir, *Remarks on solutions of $-\Delta u = u(1 - |u|^2)$ on \mathbb{R}^2* , C. R. Acad. Sci. Paris **318** (1994), pp. 327–331.
- [Sh2] I. Shafrir, *L^∞ approximation for minimizers of the Ginzburg-Landau functional*, C. R. Acad. Sci. Paris **321** (1995), pp. 705–710.
- [T] S. Terracini, *Symmetry properties of positive solutions to some elliptic equations with nonlinear boundary conditions*, Differential Integral Equations **8** (1995), pp. 1911–1922.
- [Tr] W. C. Troy, *Symmetry properties in systems of semilinear elliptic equations*, J. Differential Equations **42** (1981), pp. 400–413.
- [Z] H. Zou, *Symmetry of positive solutions of $\Delta u + u^p = 0$ in \mathbb{R}^N* , J. Differential Equations **120** (1995), pp. 46–88.

ANALYSE NUMERIQUE, UNIVERSITE P. ET M. CURIE, F-75252 PARIS CEDEX 05 FRANCE
E-mail address: brezis@ann.jussieu.fr

DEPARTMENT OF MATHEMATICS, RUTGERS, THE STATE UNIVERSITY, HILL CENTER, BUSCH CAMPUS, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NEW JERSEY 08854-8019
E-mail address: brezis@math.rutgers.edu