

WEYL GROUP ORBITS ON KAC-MOODY ROOT SYSTEMS

LISA CARBONE, WALTER FREYN, AND DIEGO PENTA

ABSTRACT. Let $\mathcal{D}$ be a Dynkin diagram, let $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ be the simple roots of the corresponding Kac-Moody root system and let W denote the Weyl group. We show that for $i \neq j$, the simple roots α_i and α_j are in the same W -orbit if and only if vertices i and j in the Dynkin diagram corresponding to α_i and α_j are connected by a path consisting only of single edges. It follows that the disjoint orbits of the Weyl group on real roots are in one-to-one correspondence with connected components of a graph obtained from the Dynkin diagram by removing all multi-edges, arrows and edge-labels. We introduce the notion of ‘the Cayley graph $\mathcal{P}$ of the Weyl group action on real roots’. The connected components of $\mathcal{P}$ are in one-to-one correspondence with the disjoint orbits of W . The vertices of $\mathcal{P}$ in a given connected component are in one-to-one correspondence with the roots in the corresponding Weyl group orbit and W acts transitively on each connected component of $\mathcal{P}$. For a symmetric hyperbolic generalized Cartan matrix A of rank ≥ 4 we prove that any 2 real roots of the same length lie in the same W -orbit. We show that if the generalized Cartan matrix A contains zeros then there are simple roots that are stabilized by simple reflections in W , that is, W does not act simply transitively on real roots. We give sufficient conditions in terms of the generalized Cartan matrix A (equivalently $\mathcal{D}$) for W to stabilize a real root. Using symmetry properties of the imaginary light cone in the hyperbolic case, we deduce that the number of W -orbits on imaginary roots on a hyperboloid of fixed radius is bounded above by the number of root lattice points on the hyperboloid that intersect the closure of the fundamental region for W .

1. INTRODUCTION

Kac-Moody algebras are the most natural generalization to infinite dimensions of the notion of a finite dimensional semisimple Lie algebra. The data for building a Kac-Moody algebra contains an integer matrix called a generalized Cartan matrix. Let A be an $n \times n$ generalized Cartan matrix with Kac-Moody algebra $\mathfrak{g} = \mathfrak{g}(A)$, root system $\Delta = \Delta(A)$, real roots $\Phi \subseteq \Delta$, and Weyl group $W = W(A)$. Let $\Pi = \{\alpha_1, \dots, \alpha_n\}$ denote a fixed base of Δ .

The group W acts on Δ , but a certain subset of Δ can be described purely in terms of the action of W on Π . This is the subset of ‘real roots’ $\Phi = W\Pi$, characterized by the property that their ‘squared lengths’ with respect to a symmetric invariant bilinear form is positive. This is in contrast to the so-called ‘imaginary roots’ in Δ whose squared length is zero or negative. If A is a Cartan matrix of finite type, then all roots are real and so $\Delta = W\Pi$ for a root basis Π , but if A is not of finite type, then Δ has a non-trivial subset of imaginary roots.

Our object of study is the orbit structure of W on the subset of real roots of an arbitrary Kac-Moody root system with Dynkin diagram $\mathcal{D}$ and simple roots $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$. We show that for $i \neq j$, the simple roots α_i and α_j are in the same W -orbit if and only if vertices i and j

The first author was supported in part by NSF grant number DMS-1101282.

in the Dynkin diagram corresponding to α_i and α_j are connected by a path consisting only of single edges. It follows that the disjoint orbits of the Weyl group on real roots are in one-to-one correspondence with connected components of a graph obtained from the Dynkin diagram by removing all multi-edges, arrows and edge-labels. Applying the classification of hyperbolic Dynkin diagrams ([CCCMNNP]), it follows that the maximal number of disjoint Weyl group orbits on real roots of a hyperbolic root system is 4.

We introduce the notion of ‘the Cayley graph $\mathcal{P}$ of the Weyl group action on real roots’ which we associate to the Dynkin diagram, or equivalently to the generalized Cartan matrix. The connected components of $\mathcal{P}$ are in one-to-one correspondence with the disjoint orbits of W . The vertices of $\mathcal{P}$ in a given connected component are in one-to-one correspondence with the roots in the corresponding Weyl group orbit and W acts transitively on each connected component of $\mathcal{P}$.

It should be noted that for a finite root system associated to a simple Lie algebra, all roots of the same length lie in the same W -orbit. This result does not extend to infinite dimensional root systems. In particular, given a Kac–Moody root system, real roots of the same length may lie in different W -orbits. Given a symmetric generalized Cartan matrix A of noncompact hyperbolic type such that real roots of the same length lie in distinct orbits, we show that there is a nonsymmetric generalized Cartan matrix A' with the same Weyl group as A and the same Coxeter matrix as A , such that the real roots of different lengths lie in distinct W -orbits. In other words, for any Kac–Moody Weyl group W there is a symmetrizable Kac–Moody algebra such that the lengths of real roots in different orbits are different.

For a symmetric hyperbolic generalized Cartan matrix A of rank ≥ 4 or symmetrizable hyperbolic generalized Cartan matrix A of rank ≥ 7 , we prove that any 2 real roots of the same length lie in the same W -orbit.

We show that if the generalized Cartan matrix A contains zeros then there are simple roots that are stabilized by simple reflections in W , that is, W does not act simply transitively on real roots. We give sufficient conditions in terms of the generalized Cartan matrix A (equivalently $\mathcal{D}$) for W to stabilize a real root.

We examine some elementary properties of the Weyl group orbits on imaginary roots using symmetry properties of the imaginary light cone in the hyperbolic case. We deduce that the number of W -orbits on imaginary roots on a hyperboloid of fixed radius is bounded above by the number of root lattice points on the hyperboloid that intersect the closure of the fundamental region for W .

The real and imaginary roots of hyperbolic root systems and their images under the Weyl group are known to have a physical interpretation. For example, in cosmological billiards, the walls of the billiard table are related to physical fluxes that, in turn, are related to real roots ([DHN]). In [BGH] Brown, Ganor and Helfgott show that real roots correspond to fluxes or instantons, and imaginary roots correspond to particles and branes. In [EHTW] Englert, Houart, Taormina, and West give a physical interpretation of Weyl group reflections in terms of M-theory. Many physical applications require a full understanding of the structure of the W -orbits on imaginary roots.

After preparation of this manuscript, we learned that in the finite dimensional case, Carter has given a characterization of the conjugacy classes of W in the case that W is finite ([Ca]). By identifying roots with $\mathbb{Z}/2\mathbb{Z}$ -subgroups Carter's work may be used to deduce some results on the structure of Weyl group orbits when W is finite.

The authors wish to thank Daniel Allcock and Siddhartha Sahi for helpful conversations. Alexander Conway contributed a great deal to the early part of this project and many of his ideas are contained within. We take pleasure in thanking him for his input. We are grateful to Yusra Naqvi for helpful discussions regarding Weyl group orbits on imaginary roots.

2. KAC-MOODY ALGEBRAS

We may construct a Kac-Moody algebra from certain data which includes a *generalized Cartan matrix*. This is a square matrix $A = (a_{ij})_{i,j \in I}$ whose entries satisfy the conditions

$$a_{ij} \in \mathbb{Z}, i, j \in I,$$

$$a_{ii} = 2, i \in I,$$

$$a_{ij} \leq 0 \text{ if } i \neq j, \text{ and}$$

$$a_{ij} = 0 \iff a_{ji} = 0.$$

We may also consider the case that A is *symmetrizable*: there exist positive rational numbers $q_1, \dots, q_\ell$, such that the matrix DA is symmetric, where $D = \text{diag}(q_1, \dots, q_\ell)$.

By a *proper submatrix* of A , we mean a matrix of the form $A_\theta = (a_{ij})_{i,j \in \theta}$, where θ is a proper subset of $\{1, \dots, \ell\}$. We say that A is *indecomposable* if there is no partition of the set $\{1, \dots, \ell\}$ into two non-empty subsets so that $a_{ij} = 0$ whenever i belongs to the first subset, while j belongs to the second.

Possible types

Finite type A is positive-definite, $\det(A) > 0$. In this case A is the Cartan matrix of a finite dimensional semisimple Lie algebra.

Affine type A is positive-semidefinite, but not positive-definite, $\det(A) = 0$.

Indefinite type A is neither of finite nor affine type, $\det(A) < 0$.

Hyperbolic type A is neither of finite nor affine type, but every proper, indecomposable submatrix is either of finite or of affine type, $\det(A) < 0$.

The Dynkin diagram of a generalized Cartan matrix

The *Dynkin diagram* $\mathcal{D}$ of a generalized Cartan matrix $A = (a_{ij})$ is the graph with one node for each row (or column) of A . If $i \neq j$ and $a_{ij}a_{ji} \leq 4$ then we connect i and j with $\max(|a_{ij}|, |a_{ji}|)$

edges together with an arrow towards i if $|a_{ij}| > 1$. If $a_{ij}a_{ji} > 4$ we draw a bold face line labeled with the ordered pair $(|a_{ij}|, |a_{ji}|)$.

2.1. The Kac-Moody algebra of a generalized Cartan matrix. Let $\langle \cdot, \cdot \rangle : \mathfrak{h}^* \longrightarrow \mathfrak{h}$ denote the natural nondegenerate bilinear pairing between a vector space $\mathfrak{h}$ and its dual.

Given:

- a generalized Cartan matrix $A = (a_{ij})_{i,j \in I}$, $I = \{1, \dots, \ell\}$, and
- a finite dimensional vector space $\mathfrak{h}$ (Cartan subalgebra) with $\dim(\mathfrak{h}) = 2\ell - \text{rank}(A)$, and
- a choice of *simple roots* $\Pi = \{\alpha_1, \dots, \alpha_\ell\} \subseteq \mathfrak{h}^*$ and *simple coroots* $\Pi^\vee = \{\alpha_1^\vee, \dots, \alpha_\ell^\vee\} \subseteq \mathfrak{h}$ such that Π and $\Pi^\vee$ are linearly independent and such that $\langle \alpha_j, \alpha_i^\vee \rangle = \alpha_j(\alpha_i^\vee) = a_{ij}$, $i = 1, \dots, \ell$,

we may associate a Lie algebra $\mathfrak{g} = \mathfrak{g}(A)$ over K , a field, generated by $\mathfrak{h}$ and elements $(e_i)_{i \in I}$, $(f_i)_{i \in I}$ subject to relations ([M1], [M2], and [K], Theorem 9.11):

$$[\mathfrak{h}, \mathfrak{h}] = 0,$$

$$[h, e_i] = \alpha_i(h)e_i, h \in \mathfrak{h},$$

$$[h, f_i] = -\alpha_i(h)f_i, h \in \mathfrak{h},$$

$$[e_i, f_i] = \alpha_i^\vee,$$

$$[e_i, f_j] = 0, i \neq j,$$

$$(\text{ad } e_i)^{-a_{ij}+1}(e_j) = 0, i \neq j,$$

$$(\text{ad } f_i)^{-a_{ij}+1}(f_j) = 0, i \neq j,$$

where $(\text{ad}(x))(y) = [x, y]$.

2.2. Weyl group. For each simple root α_i , $i = 1, \dots, \ell$, we define the simple root reflection

$$w_i(\alpha_j) = \alpha_j - \alpha_j(\alpha_i^\vee)\alpha_i.$$

It follows from the formula that $w_i(\alpha_i) = -\alpha_i$. The w_i generate a subgroup

$$W = W(A) \subseteq \text{Aut}(\mathfrak{h}^*),$$

called the *Weyl group* of A . The group W acts on the set of all roots. For $i, j \in I$, and for $i \neq j$, we set

$$c_{ii} = 1, \quad c_{ij} = 2, 3, 4, 6, \text{ or } \infty$$

according as

$$a_{ij}a_{ji} = 0, 1, 2, 3, \text{ or } \geq 4$$

respectively. Then $W = W(A)$ is the Coxeter group with presentation:

$$W = \langle w_i \mid i \in I, (w_i w_j)^{c_{ij}} = 1, \text{ if } c_{ij} \neq \infty \rangle.$$

If $\mathfrak{g}$ is of finite type, W is a finite group. If $\mathfrak{g}$ is infinite dimensional, W is infinite. The group W acts on the set of all roots.

2.3. Invariant form. If A is a symmetrizable generalized Cartan matrix, the algebra $\mathfrak{g} = \mathfrak{g}(A)$ admits a well-defined non-degenerate symmetric bilinear form $(\cdot | \cdot)$ which plays the role of ‘squared length’ of a root ([K], Theorem 2.2). If $\mathfrak{g}$ is of finite type, then $(\cdot | \cdot)$ is the usual Killing form. The form $(\cdot | \cdot)$ is an analog of the Killing form if $\mathfrak{g} = \mathfrak{g}(A)$ is infinite dimensional and $(\cdot | \cdot)|_{\mathfrak{h}^*}$ is W -invariant.

2.4. Real and imaginary roots. If $\mathfrak{g} = \mathfrak{g}(A)$ is finite dimensional, all roots are Weyl group translates of simple roots. That is,

$$\Delta = W\Pi.$$

For infinite dimensional Kac-Moody root systems, there are additional mysterious roots of negative norm (‘squared length’) called imaginary roots.

A root $\alpha \in \Delta$ is called a *real root* if there exists $w \in W$ such that $w\alpha$ is a simple root. A root α which is not real is called *imaginary*. We denote by Δ^{re} the real roots, Δ^{im} the imaginary roots, $\Delta_{\pm}^{re}$ the positive and negative Weyl roots, and $\Delta_{\pm}^{im}$ the positive and negative imaginary roots. Then

$$\begin{aligned}\Delta &= \Delta^{re} \sqcup \Delta^{im} \\ \Delta^{re} &= \Delta_+^{re} \sqcup \Delta_-^{re}, \\ \Delta^{im} &= \Delta_+^{im} \sqcup \Delta_-^{im},\end{aligned}$$

and

$$\Delta^{re} = W\Pi.$$

Real and imaginary roots may be characterized by their ‘squared length’. If A is symmetrizable, we have ([K], Prop 5.1 and 5.2):

$$\begin{aligned}\alpha \in \Delta^{re} &\iff (\alpha | \alpha) > 0 \\ \alpha \in \Delta^{im} &\iff (\alpha | \alpha) \leq 0.\end{aligned}$$

3. WEYL GROUP ORBITS ON REAL ROOTS

Let A be a generalized Cartan matrix. Let $\mathcal{D} = \mathcal{D}(A)$ be the corresponding Dynkin diagram with vertices indexed by $I = \{1, 2, \dots, \ell\}$. We let $\mathcal{D}_*$ denote the graph obtained from $\mathcal{D}$ by deleting all multiple edges, including arrows and edge labels. Let $\mathcal{D}_1, \dots, \mathcal{D}_s$ denote the connected subdiagrams of $\mathcal{D}_*$. We call $\mathcal{D}_*$ the *skeleton* of $\mathcal{D}$. We may describe the graph $\mathcal{D}_*$ as follows

$Vertices(\mathcal{D}_*) = Vertices(\mathcal{D})$ with the same labelling, that is, indexed by $I = \{1, 2, \dots, \ell\}$

$Edges(\mathcal{D}_*) = \bigcup_{i=1}^s Edges(\mathcal{D}_i)$

Vertices i and j are adjacent in $\mathcal{D}_*$ if and only if i and j are connected in $\mathcal{D}$ by a single edge with no arrows or edge labels if and only if $a_{ij} = a_{ji} = -1$.

In the discussions that follow, we let A be a generalized Cartan matrix, $W(A)$ be its Weyl group, Π the set of simple roots indexed by I , Φ the set of real roots, and $\mathcal{D}$ the corresponding

Dynkin diagram. Our main objective is to classify the Weyl orbits on Φ using only the associated Dynkin diagram. This can be accomplished using a lemma of Tits in association with the theorem below, which requires some results regarding symmetric rank 2 root subsystems.

We start with some facts from the theory of finite dimensional Lie algebras, and the notion of ‘vector projection’ from basic linear algebra. First, we know Π is a basis of Φ . Thus if $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$, then any real root $\beta \in \Phi$ can be written as a linear combination of the simple roots,

$$\beta = \sum_{i=1}^{\ell} b_i \alpha_i,$$

where the coordinates $\{b_i\}$ are integers, and are uniquely determined by β . Moreover, the coordinates are either all positive or all negative. In the results that follow, we will need to focus our attention on specific coordinates of a given real root, hence we make the following definition.

Definition 3.1. Let $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$. Given any real root $\beta \in \Phi$, where $\beta = \sum_{i=1}^{\ell} b_i \alpha_i$, we define the *i-projection* of β to be

$$\rho_i(\beta) := b_i.$$

In other words, $\rho_i(\beta)$ is the i^{th} coordinate of β with respect to the root basis Π . For simple root α_j ,

$$\rho_i(\alpha_j) = \delta_{ij}$$

where δ_{ij} is the Kronecker delta function.

Now consider the rank 2 symmetric generalized Cartan matrix A ,

$$A = \begin{pmatrix} 2 & -a \\ -a & 2 \end{pmatrix}.$$

Figure 1, which shows the root systems corresponding to $a = 2$ and $a = 3$, will help to illustrate the sequences described in the next lemma. Figure 3 also shows an example of the 1- and 2-projections of a root β lying on the root lattice.

Lemma 3.2. Let $A = \begin{pmatrix} 2 & -a \\ -a & 2 \end{pmatrix}$ for $a \geq 2$, $\Pi = \{\alpha_1, \alpha_2\}$ simple roots. Let $\beta_n = (w_2 w_1)^n(-\alpha_1)$ and $\gamma_n = (w_1 w_2)^n(-\alpha_1)$ for some $n \in \mathbb{Z}_{\geq 0}$. Denote by $(\rho_1(\beta_n))_{n \in \mathbb{Z}_{\geq 0}}$ and $(\rho_1(\gamma_n))_{n \in \mathbb{Z}_{\geq 0}}$ the sequences of 1-projections of real roots β_n and γ_n for all $n \in \mathbb{Z}_{\geq 0}$. Then

- (i) $(\rho_1(\beta_n))_{n \in \mathbb{Z}_{\geq 0}}$ increasing sequence of integers.
- (ii) $(\rho_1(\gamma_n))_{n \in \mathbb{Z}_{\geq 0}}$ is a strictly decreasing sequence of integers.

Proof. (i) We proceed by induction on n , and use the following general calculation for simple root reflection w_i acting on simple root α_j :

$$w_i \alpha_j = \alpha_j - \langle \alpha_j, \alpha_i^\vee \rangle \alpha_i = \alpha_j - a_{ji} \alpha_i = \alpha_j + a \alpha_i,$$




FIGURE 1. Rank 2 affine and hyperbolic symmetric root systems respectively

where a_{ji} is the $\{j, i\}$ entry of A . For the base case we compare

$$\rho_1(\beta_0) = \rho_1(-\alpha_1) = -1$$

and

$$\rho_1(\beta_1) = \rho_1(w_2 w_1(-\alpha_1)) = \rho_1(\alpha_1 + a\alpha_2) = 1,$$

and see that the sequence increases. Now assume that $(\rho_1(\beta_n))_n$ is an increasing sequence of integers for all n up to some $N \in \mathbb{Z}_{>0}$. Let $\beta_N = b_1\alpha_1 + b_2\alpha_2$. Then

$$w_2 w_1(\beta_N) = (ab_2 - b_1)\alpha_1 + (a(ab_2 - b_1) - b_2)\alpha_2.$$

Since $\rho_1(\beta_1) = 1$, by the induction hypothesis, $b_1 = \rho_1(\beta_N) > \rho_1(\beta_1) = 1$, and so b_1, b_2 are both positive. We recall that when we identify the simple roots Π with the canonical basis vectors of $\mathbb{R}^2$, then the real roots Φ lie on the quadratic form

$$Q(x, y) = 2x^2 - 2axy + 2y^2.$$

For $a \geq 2$, this quadratic form is a hyperbola comprised of two ‘branches’, an upper branch and lower branch, as illustrated in Figure 1. We recall that $w \in W$ is a translation along branches if and only if w is of the form $(w_1 w_2)^n$ or $(w_2 w_1)^n$ for some $n \in \mathbb{Z}$. Therefore, since β_N is a

translation of $-\alpha_1$, β_N lies on the upper branch. Hence, β_N lies strictly above the asymptote $y = \frac{a}{2}x$ in $\mathbb{R}^2$, so $b_2 > \frac{a}{2}b_1$. Thus we have

$$\rho_1(\beta_{N+1}) = \rho_1(w_2 w_1 \beta_N) = ab_2 - b_1 > \left(\frac{a^2}{2} - 1\right) b_1 \geq b_1 = \rho(\beta_N),$$

where the last inequality is due to $a \geq 2$. Hence (1) is true for all n . The proof of (ii) is similar, and we leave it to the reader. $\square$

We are now ready to show that the Weyl group W is not transitive on the symmetric rank 2 root system investigated above if $a \geq 2$.

Lemma 3.3. *Let A be the rank 2 generalized Cartan matrix $A = \begin{pmatrix} 2 & -a \\ -a & 2 \end{pmatrix}$, and $\mathcal{D}$ its associated Dynkin diagram with 2 vertices. Let $\Pi = \{\alpha_1, \alpha_2\}$ be the simple roots. If $a \geq 2$ (equivalently if $\mathcal{D}$ is affine or hyperbolic), then $W\{\alpha_1\} \cap W\{\alpha_2\} = \emptyset$, that is, there is no $w \in W$ such that $w(\alpha_1) = \alpha_2$.*

Proof. Assume there is an element $w \in W$ such that $w(\alpha_1) = \alpha_2$. Without loss of generality, assume w is reduced (or simply replace w with its corresponding unique reduced word). Then w has a first element. Since W is generated by 2 simple Weyl reflections, this element must either be w_1 or w_2 .

Case 1: Assume w begins with w_1 , so $w = u \cdot w_1$ for some Weyl word $u \in W$. Then

$$\alpha_2 = w(\alpha_1) = u \cdot w_1(\alpha_1) = u(-\alpha_1).$$

Both $-\alpha_1$ and α_2 lie on the upper branch of Φ , therefore u must be a translation of the form $u = (w_1 w_2)^m$ or $u = (w_2 w_1)^m$ for some $m \in \mathbb{Z}_{>0}$. However, since w is reduced, u must begin with w_2 , so we omit the latter case. Thus,

$$u(-\alpha_1) = (w_1 w_2)^m(\alpha_1) = \alpha_2.$$

By Lemma 3.2(ii), we now have the following decreasing sequence of integers:

$$\rho_1(-\alpha_1) > \rho_1(w_1 w_2(-\alpha_1)) > \cdots > \rho_1((w_1 w_2)^m(-\alpha_1)) > \cdots.$$

For $n = 0$, $\rho_1(-\alpha_1) = -1$, and for $n = m$, $\rho_1((w_1 w_2)^m(-\alpha_1)) = \rho_1(\alpha_2) = 0$. So the sequence is

$$-1 > \cdots > 0 > \cdots,$$

which is false. Therefore w cannot begin with w_1 .

Case 2: Now assume w begins with w_2 , so $w = v \cdot w_2$ for some Weyl word $v \in W$. Then

$$\alpha_2 = w(\alpha_1) = v \cdot w_2(\alpha_1) = v(\alpha_1 + a\alpha_2).$$

Both α_2 and $\alpha_1 + a\alpha_2$ lie on the upper branch of Φ , and since w is reduced, v must begin with w_1 . Therefore, v is a translation of the form $v = (w_2 w_1)^m$ for some $m \in \mathbb{Z}$. Thus,

$$v(\alpha_1 + a\alpha_2) = (w_2 w_1)^m(\alpha_1 + a\alpha_2) = \alpha_2.$$

Observe that $\alpha_1 + a\alpha_2 = (w_2w_1)(-\alpha_1)$. Therefore, by Lemma 3.2(i), we have the following increasing sequence of integers:

$$\rho_1(\alpha_1 + a\alpha_2) < \rho_1(w_2w_1(\alpha_1 + a\alpha_2)) < \cdots < \rho_1((w_2w_1)^m(\alpha_1 + a\alpha_2)) < \cdots.$$

But $\rho_1(\alpha_1 + a\alpha_2) = 1$ and $\rho_1((w_2w_1)^m(\alpha_1 + a\alpha_2)) = \rho_1(\alpha_2) = 0$, so the sequence is

$$1 < \cdots < 0 < \cdots,$$

which is also false. Thus, w does not begin with w_2 .

Since w has no first element, w does not exist. □

We now formalize the notion of ‘fixing’ root projections.

Lemma 3.4. *Let A be any generalized Cartan matrix, $\beta \in \Phi$, $r \in I$. Then if $i \neq r$,*

$$\rho_i(w_r(\beta)) = \rho_i(\beta).$$

In this sense, w_r can be said to ‘fix’ all but the r -projection b_r of $\beta = \sum_{i=1}^{\ell} b_i\alpha_i$.

Proof. Let w_r act on a root $\beta \in \Phi$:

$$w_r\beta = w_r \sum_{i=1}^{\ell} b_i\alpha_i = \sum_{i=1}^{\ell} b_i(w_r\alpha_i) = \sum_{i=1}^{\ell} b_i(\alpha_i - a_{ir}\alpha_r) = \sum_{i=1, i \neq r}^{\ell} b_i\alpha_i + \left(b_r - \sum_{i=1}^{\ell} b_i a_{ir}\right) \alpha_r.$$

Thus we see that for all $i \neq r$, $\rho_i(w_r(\beta)) = b_i = \rho_i(\beta)$. □

Our objective is to prove that two simple roots lie in the same W -orbit if and only if their corresponding vertices lie in the same connected component of D_* . In other words, their corresponding vertices in the Dynkin diagram are connected by a path of single edges. To prove the forward argument we use the following lemma of Tits, which we will strengthen with the aid of the foregoing results.

Lemma 13.31 ([T]). *Let $\mathcal{D}$ be a Dynkin diagram, W its associated Weyl group, and Π the set of simple roots indexed over a set I . If for some $j, k \in I$, $j \neq k$ there exists $w \in W$ such that*

$$w\alpha_j = \alpha_k,$$

then there exists a sequence $\{u_1, \dots, u_{m-1}\}$ of elements of W , and there exist sequences $\{j = i_1, \dots, i_m = k\}$, $i_t \in I$, and $\{s_1, \dots, s_{m-1}\}$, $s_t \in I$ such that

$$w = u_{m-1}u_{m-2} \cdots u_2u_1$$

where $u_t(\alpha_{i_t}) = \alpha_{i_{t+1}}$ and $u_t \in \langle w_{i_t}, w_{s_t} \rangle$, $t \in \{1, \dots, m-1\}$.

Essentially, if two simple roots α_j and α_k lie in the same Weyl orbit (that is, they are connected by a reduced Weyl element w), then Tits’ lemma concludes that w is a product of elements from the first sequence $\{u_t\}$. Moreover, we observe that the second sequence $\{i_t\}$ corresponds to a sequence of simple roots, starting with α_j and ending with α_k . Since each u_t maps α_{i_t} to the

next simple root $\alpha_{i_{t+1}}$ in this sequence, the Weyl word w maps α_j to α_k through the other $m - 2$ simple roots in sequential order. Figure 2 illustrates this conclusion.




FIGURE 2. The conclusion of Tits' lemma.

We must prove that the hypotheses of the Lemma are sufficient for vertices $j = i_1$ and $k = i_m$ in the Dynkin diagram to be connected by a path of single edges. If the sequence $\{j = i_1, \dots, i_m = k\}$ corresponds to such a path, then we will need a full description of the rank 2 root subsystems generated by each pair of simple roots α_{i_t} and $\alpha_{i_{t+1}}$ in the sequence.

However, the statement of the Lemma does not clearly define the elements u_t which comprise w , that is, for each u_t a full description of the simple reflection w_{s_t} is not provided. In addition, if u_t is a reduced word in $\langle w_{i_t}, w_{s_t} \rangle$, then there is no indication of its length.

In the next theorem, we determine these unknowns without ambiguity, and we follow with a much sharper statement of the lemma of Tits. This later result will give us the sufficiency condition needed to allow us to classify the Weyl group orbits of any hyperbolic Kac–Moody root system based solely on its associated Dynkin diagram.

Theorem 3.5. *Let $\mathcal{D}$ be a Dynkin diagram, Π a basis of its associated root system, I the index set for Π , and W the corresponding Weyl group. Let $\widetilde{W} \subset W$ be generated by 2 simple Weyl reflections. If for some $j, k \in I$ there exists a reduced word $w \in \widetilde{W}$ such that $w(\alpha_j) = \alpha_k$, then*

- 1) $A_{(j,k)} = A_2$,
- 2) j, k are adjacent in $\mathcal{D}_*$,
- 3) $w = w_j w_k$,
- 4) $\widetilde{W} = \langle w_j, w_k \rangle$.

Proof. (1) Denote by $A_{(j,k)}$ the rank 2 submatrix of A ,

$$A_{(j,k)} = \begin{pmatrix} 2 & a_{jk} \\ a_{kj} & 2 \end{pmatrix}.$$

Since W preserves root length, and α_k is a Weyl translate of α_j , $A_{(j,k)}$ must be symmetric, else α_j, α_k are of different lengths. Thus we have

$$A_{(j,k)} = \begin{pmatrix} 2 & -a \\ -a & 2 \end{pmatrix},$$

where $a = -a_{jk} = -a_{kj} \in \mathbb{Z}_{\geq 0}$, and $a \geq 0$. Assume $a \neq 1$.

Case 1: If $a = 0$ then $A_{(j,k)}$ is the Cartan matrix associated to $A_1 \times A_1$. Calculating the $\widetilde{W}$ -orbit of α_j we have

$$\begin{aligned} w_k \alpha_j &= \alpha_j - (\langle \alpha_j, \alpha_k^\vee \rangle) \alpha_k = \alpha_j - (a) \alpha_k = \alpha_j, \\ w_j \alpha_j &= -\alpha_j, \\ w_k(-\alpha_j) &= -w_k(\alpha_j) = -\alpha_j, \\ w_j(-\alpha_j) &= \alpha_j, \end{aligned}$$

and thus, $\widetilde{W}\{\alpha_j\} = \{\pm\alpha_j\}$, which does not contain α_k , a contradiction.

Case 2: If $a \geq 2$, then $A_{(j,k)}$ is of affine or hyperbolic type, so by Lemma 3.3, $W\{\alpha_j\} \cap W\{\alpha_k\} = \emptyset$, contradicting the existence of $w \in \widetilde{W}$ where $w(\alpha_j) = \alpha_k$.

Thus, a must equal 1, and so $A_{(j,k)} = A_2$.

(2) By part (1), α_j and α_k span an A_2 root subsystem. Thus vertices j and k are adjacent in $\mathcal{D}_*$ and thus in $\mathcal{D}$ (Figure 3).




FIGURE 3. Adjacent vertices j and k in $\mathcal{D}_*$

(3) Recalling from Lemma 3.2 that $w_i \alpha_j = \alpha_j + a \alpha_i$ and using $a = 1$, we have

$$\begin{aligned} w_j w_k \alpha_j &= w_j(\alpha_j + \alpha_k) \\ &= \alpha_j + \alpha_k - \langle \alpha_j + \alpha_k, \alpha_j^\vee \rangle \alpha_j \\ &= \alpha_j + \alpha_k - (\langle \alpha_j, \alpha_j^\vee \rangle + \langle \alpha_k, \alpha_j^\vee \rangle) \alpha_j \\ &= \alpha_j + \alpha_k - (a_{jj} + a_{kj}) \alpha_j \\ &= \alpha_j + \alpha_k - (2 - 1) \alpha_j \\ &= \alpha_k. \end{aligned}$$

(4) By part (3) we see that $w_j, w_k \in \widetilde{W}$. By assumption $\widetilde{W}$ is generated by two simple Weyl reflections, thus $\widetilde{W} = \langle w_j, w_k \rangle$, and we are done. $\square$

We now obtain a strengthening of Tits' lemma as a corollary.

Corollary 3.6. *Let $\mathcal{D}$ be a Dynkin diagram, W its associated Weyl group, and Π the set of simple roots indexed over a set I . Let $w \in W$ and $j, k \in I$ be such that $w \alpha_j = \alpha_k$. Then there exists a sequence $\{j = i_1, i_2, \dots, i_m = k\} \subset I$ such that*

$$w = (w_{i_{m-1}} w_k)(w_{i_{m-2}} w_{i_{m-1}})(w_{i_{m-3}} w_{i_{m-2}}) \cdots (w_{i_2} w_{i_3})(w_j w_{i_2}),$$

and i_t, i_{t+1} are adjacent in $\mathcal{D}_*$ for all $t \in \{1, \dots, m-1\}$. Thus, j and k are connected by a path of single edges in $\mathcal{D}$.

Proof. By the lemma of Tits, we have a sequence of Weyl elements $u_1, u_2, \dots, u_{m-1}$ and a sequence $\{s_1, \dots, s_{m-1}\}$ such that

$$w = u_{m-1} \cdots u_2 u_1$$

where

- (i) $u_t(\alpha_{i_t}) = \alpha_{i_{t+1}}$ and
- (ii) $u_t \in \langle w_{i_t}, w_{s_t} \rangle$.

The description of w_{s_t} here is made clear by applying Theorem 3.5. The last two conditions together imply that $s_t = i_{t+1}$. Moreover, they imply that for all $t \in \{1, \dots, m-1\}$, i_t and i_{t+1} are adjacent, and $u_t = (w_{i_t} w_{i_{t+1}})$. Thus we may rewrite the Weyl element w as

$$w = (w_{i_{m-1}} w_k)(w_{i_{m-2}} w_{i_{m-1}})(w_{i_{m-3}} w_{i_{m-2}}) \cdots (w_{i_2} w_{i_3})(w_j w_{i_2}),$$

and so j and k are connected by a path $\{j = i_1, i_2, \dots, i_m = k\}$ in $\mathcal{D}_*$. Thus they are in the same connected component. $\square$

In strengthening the lemma of Tits, we now make clear that the sequence $\{i_1, \dots, i_m\}$ indeed corresponds to a sequence of vertices in D_*^J for some J , where each pair i_t, i_{t+1} is connected by a single edge. Thus the condition that α_j and α_k lie in the same Weyl group orbit is sufficient for vertices $j, k \in \mathcal{D}$ to be connected by a path of single edges. Now we show that this condition is also necessary.

Let $\mathcal{D}$ be a Dynkin diagram, $\mathcal{D}_*$ its skeleton, $\Pi = \{\alpha_1, \alpha_2, \dots, \alpha_\ell\}$ a choice of simple roots and W the Weyl group.

Lemma 3.7. *Suppose vertices i and j are adjacent in $\mathcal{D}_*$. Then $\alpha_i = w_j w_i \alpha_j$, where w_j is the simple root reflection corresponding to α_j .*

Proof. Since i and j are adjacent in $\mathcal{D}_*$, there is a single edge between i and j in $\mathcal{D}$, and hence $a_{ij} = a_{ji} = -1$ in the generalized Cartan matrix A . The proof of Theorem 3.5 part(4) has already established that if $a_{ij} = a_{ji} = -1$, then $w_j w_i$ maps α_j to α_i . $\square$

The following corollary is immediate.

Corollary 3.8. *If $i_1, i_2, \dots, i_m$ are the vertices of a path in $\mathcal{D}_*$, then*

- (i) *There exists $w \in \langle w_{i_1}, \dots, w_{i_m} \rangle$ such that $\alpha_{i_m} = w \alpha_{i_1}$.*
- (ii) *We have $a_{i_1 i_2} = a_{i_2 i_3} = \dots = a_{i_{m-1} i_m} = a_{i_2 i_1} = a_{i_3 i_2} = \dots = a_{i_m i_{m-1}} = -1$.*

Theorem 3.9. *Let $\mathcal{D}$ be a Dynkin diagram, let $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ be the simple roots of the corresponding root system and let W denote the Weyl group. Let $\mathcal{D}_*^J$ denote a connected subdiagram of $\mathcal{D}_*$ whose vertices are indexed by $J \subseteq I$. Then*

- (1) *For all $i, j \in J$ there exists $w \in W$ such that $\alpha_i = w \alpha_j$, thus $W \alpha_i = W \alpha_j$.*
- (2) *If $j \in J$ and $k \notin J$, we have $W \alpha_j \cap W \alpha_k = \emptyset$.*

Proof. (1) Since $i, j \in J$, there exists a connected path of vertices $\{i = i_1, i_2, \dots, i_{m-1}, i_m = j\}$ in $\mathcal{D}_*^J$ and a corresponding subgroup $\langle w_{i_1}, \dots, w_{i_m} \rangle \subset W$. Thus we may use Corollary 3.8(i), and the result follows.

(2) Assume $j \in J$ and $k \notin J$, so j, k are not in the same connected component. Assume that $W\{\alpha_j\} = W\{\alpha_k\}$. Then there exists a $w \in W$ such that $w\alpha_j = \alpha_k$. By Corollary 3.6, there exists a sequence $\{j = i_1, i_2, \dots, i_m = k\} \subset I$ such that

$$w = (w_{i_{m-1}} w_k)(w_{i_{m-2}} w_{i_{m-1}})(w_{i_{m-3}} w_{i_{m-2}}) \cdots (w_{i_2} w_{i_3})(w_j w_{i_2}),$$

and i_t, i_{t+1} are adjacent in $\mathcal{D}_*$ for all $t \in \{1, \dots, m-1\}$. Thus there exists a path of vertices $\{j = i_1, i_2, \dots, i_m = k\}$ connecting j and k . But then $k \in J$, which is a contradiction. Therefore $W\alpha_j \cap W\alpha_k = \emptyset$. $\square$

With Theorem 3.9 and Corollary 3.6, we now have the necessary and sufficient conditions for two distinct simple roots to be in the same Weyl group orbit. This result is summarized in the following corollary, which is immediate.

Corollary 3.10. *Let $\mathcal{D}$ be a Dynkin diagram, let $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ be the simple roots of the corresponding root system and let W denote the Weyl group. Then for $i \neq j$, the simple roots α_i and α_j are in the same W -orbit if and only if vertices i and j in the Dynkin diagram corresponding to α_i and α_j are connected by a path consisting only of single edges.*

Moreover, we have the following stronger statement.

Corollary 3.11. *Let $\mathcal{D}$ be a Dynkin diagram, let $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$ be the simple roots of the corresponding root system and let W denote the Weyl group. Let $\mathcal{D}_*^{J_1}, \mathcal{D}_*^{J_2}, \dots, \mathcal{D}_*^{J_t}$ denote the connected subdiagrams of $\mathcal{D}_*$ whose vertices are indexed by $J_1, J_2, \dots, J_t \subseteq I$. Then the set of real roots $\Phi = W\Pi$ is*

$$\Phi = W\{\alpha_{J_1}\} \Pi W\{\alpha_{J_2}\} \Pi \cdots \Pi W\{\alpha_{J_t}\},$$

where α_{J_s} denotes the subset of simple roots indexed by the subset $J_s \subseteq I$.

Given any Dynkin diagram $\mathcal{D}$, we can therefore determine the disjoint orbits of the Weyl group on real roots by determining the skeleton $\mathcal{D}_*$. Applying the classification of hyperbolic Dynkin diagrams ([CCCMNNP], [KM], [Li], [Sa]), it follows that maximal number of disjoint Weyl group orbits in a hyperbolic root system is 4.

4. A CAYLEY GRAPH ENCODING THE WEYL GROUP ACTION ON REAL ROOTS

Let A be an $n \times n$ generalized Cartan matrix with Kac–Moody algebra $\mathfrak{g} = \mathfrak{g}(A)$, root system $\Delta = \Delta(A)$, real roots $\Phi \subseteq \Delta$, and Weyl group $W = W(A)$. Let $\Pi = \{\alpha_1, \dots, \alpha_n\}$ denote a fixed base of Δ . Using only the action of the simple root reflections on Φ , we define a graph $\mathcal{P}$ associated to Φ , relative to Π , which has the following properties:

- (i) The vertices of $\mathcal{P}$ are in one-to-one correspondences with the elements of Φ .
- (ii) The number of connected components of $\mathcal{P}$ equals the number of disjoint orbits of W on Φ , and W acts transitively on each connected component of $\mathcal{P}$.
- (iii) The vertices of $\mathcal{P}$ in a given connected component are in 1-to-1 correspondence with the roots in the corresponding Weyl orbit.
- (iv) The number of orbits of W on Φ (equivalently the number of connected components of $\mathcal{P}$) can be determined from the generalized Cartan matrix A (equivalently from the Dynkin diagram $\mathcal{D}$ of A).

The graph $\mathcal{P}$ is defined as follows.

Definition 4.1. Let A be a (generalized) Cartan matrix, $A = (A_{ij})_{i,j=1,2,\dots,l}$. Let Π be the root basis for A , W be the Weyl group corresponding to A and $\Phi = W\Pi$ be the real roots of A . We define a labelled graph $(\mathcal{P}, \nu)$ which encodes the action of W on Φ where $\mathcal{P} = (V\mathcal{P}, E\mathcal{P})$ is an unoriented graph (with vertices $V\mathcal{P}$ and edges $E\mathcal{P}$) and $\nu : E\mathcal{P} \rightarrow S$ is a labelling of a simple reflection to each edge of $\mathcal{P}$. Vertices and edges of $\mathcal{P}$ are defined as follows:

$$\text{Vertices}(\mathcal{P}) = \Phi \quad (\text{the set of real roots})$$

$$\text{Edges}(\mathcal{P}) \subseteq \Phi \times \Phi$$

where α and β are adjacent in $\mathcal{P}$ if and only if

$$w_i\alpha = \beta \quad \text{for some simple reflection } w_i \text{ and}$$

$$\nu(e) = w_i \quad \text{where } e = (\alpha, \beta) \text{ is an edge of } \mathcal{P}.$$

We will customarily label the vertices with the corresponding real root, expressed either as a linear combination of simple roots or, equivalently, an n -tuple of i -projections.

It should be noted that, since every $w_i \in W$ is an involution, every edge in $\mathcal{P}$ represents a bigon (Figure 4).




FIGURE 4. The bigon $w_i^2 = 1$

By definition of $\mathcal{P}$, each orbit of W , respectively each connected component of $\mathcal{P}$, contains at least one simple root. Moreover $\mathcal{P}$ is connected if and only if W acts transitively on Φ , and $\mathcal{P}$ has no loops if and only if W acts simply transitively on connected components, that is, with no fixed roots.

Let $\mathcal{P}_j$, ($j = 1, \dots, s$) denote the connected components of $\mathcal{P}$. In the case that W acts simply transitively on the vertices of $\mathcal{P}$, there is a natural isomorphism from $\mathcal{P}$ to the Cayley graph of W . In fact, if W acts simply transitively on a connected component $\mathcal{P}_j$, then there is a natural isomorphism from $\mathcal{P}_j$ to the Cayley graph of W . There are important examples of finite, affine and hyperbolic root systems for which this criterion holds.

By (ii) the action of W is transitive on each connected component $\mathcal{P}_j$. If, however, the action of W is not simply transitive on some $\mathcal{P}_j$, then there is a bijective correspondence between the vertices of $\mathcal{P}_j$ and elements of the coset space $W/W_{\alpha_{i_j}}$, where α_{i_j} is a choice of simple root from the connected component $\mathcal{P}_j$ and $W_{\alpha_{i_j}}$ denotes the subgroup of W stabilizing α_{i_j} , well defined up to conjugacy. This correspondence induces an isomorphism of $\mathcal{P}_j$ with a certain ‘orbifold’ associated to the Cayley graph of W .

We may therefore consider the graph $\mathcal{P}$ as a generalization of the Cayley graph of W , namely we may view $\mathcal{P}$ as ‘the Cayley graph of the action of W on Φ relative to Π .’ This notion may

be generalized further to associate a graph $\mathcal{P}$ to any action of a group G on a set X relative to a choice of generating set for G .

5. THE REFLECTION GEOMETRY OF ROOTS

Here we investigate the geometry governing the action of the Weyl group on roots. Let $\mathfrak{h}$ be a Cartan subalgebra of the Kac–Moody algebra $\mathfrak{g}$ and let α be a root. Associated to the root α is a root vector $h_\alpha \in \mathfrak{h}$, a hyperplane H_α and an involution, s_α such that $s_\alpha|_{H_\alpha} = \text{Id}$ and $s_\alpha(h_\alpha) = -h_\alpha$. Hence, with respect to the invariant form $(\cdot | \cdot)$, the hyperplane H_α is orthogonal to the root vector h_α .

Let now β denote another root. The following observation is clear:

$$s_\alpha(\beta) = \beta \quad \text{if and only if} \quad h_\beta \in H_\alpha.$$

Assume α and β are simple roots in a basis Π of the root system. Then

$$h_\beta \in H_\alpha \quad \text{if and only if} \quad a_{\alpha,\beta} = a_{\beta\alpha} = 0$$

where $a_{\alpha,\beta} = a_{\beta\alpha}$ is the α, β -entry of the generalized Cartan matrix.

If α and β generate a finite root system of type B_2 or C_2 such that α is a short root and β is a long root, then the roots $\pm(\alpha + \beta)$ are in H_α and the roots $\pm(\beta + 2\alpha)$ are in H_β .

The following lemma is easy to establish.

Lemma 5.1. *Let $\mathfrak{g}$ be a symmetrizable Lie algebra or Kac–Moody algebra. Let α be any positive root. Let w_j be a simple root reflection. Then $w_j\alpha$ is also a positive root, unless α equals the simple root α_j in which case $w_j\alpha_j = -\alpha_j$.*

6. SOME FINITE, AFFINE AND HYPERBOLIC EXAMPLES

Consider the Cartan matrix $A_1 = (2)$ for the A_1 root system. Then $\Phi = \{\alpha, -\alpha\}$, and $\mathcal{P}(A_1)$ is given in Figure 5.



FIGURE 5. Orbit of type A_1

For

$$A_2 = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},$$

we have $\Phi = \{\pm\alpha_1, \pm\alpha_2, \pm(\alpha_1 + \alpha_2)\}$, and $\mathcal{P}(A_2)$ is given in Figure 6.

$\mathcal{P}(A_2)$ illustrates that the Weyl group acts transitively on all roots in A_2 : $\Phi(A_2) = W\alpha_1 = W\alpha_2$.




FIGURE 6. Orbit of type A_2

An example in which $\mathcal{P}$ is disconnected is

$$B_2 = \begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix},$$

whose simple roots have two distinct lengths. The Dynkin diagram is given in Figure 7.




FIGURE 7. Dynkin diagram of type B_2

has skeleton as in Figure 8.



FIGURE 8. The skeleton $\mathcal{D}_*(B_2)$

thus by Corollary 3.11,

$$\Phi(B_2) = W\alpha_1 \amalg W\alpha_2.$$

The disjoint Weyl orbits of B_2 , as well as the real roots which are fixed by simple Weyl reflections, are demonstrated in $\mathcal{P}(B_2)$ which is given in Figure 9




FIGURE 9. Orbit of type B_2

The next example is of affine type. Let

$$A_1^{(1)} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}.$$

Its Dynkin diagram is given in Figure 10.



FIGURE 10. Dynkin diagram of type $A_1^{(1)}$

The skeleton $\mathcal{D}_*(A_1^{(1)})$ is given in Figure 11



FIGURE 11. The skeleton $\mathcal{D}_*(A_1^{(1)})$

thus by Corollary 3.11,

$$\Phi(A_1^{(1)}) = W\alpha_1 \amalg W\alpha_2.$$

The disjoint W -orbits are demonstrated in $\mathcal{P}(A_1^{(1)})$ (Figure 12). Here we have chosen to label each root β with its corresponding 2-tuple of projections $(\rho_1(\beta), \rho_2(\beta))$:




FIGURE 12. Root orbit of $A_1^{(1)}$

Consider the hyperbolic example

$$A_2^* = \begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}.$$

The graph $\mathcal{P}(A_2^*)$ is given in Figure 13.

where F_n denotes the n -th Fibonacci number ([ACP]).

The final two examples correspond to rank 3 root systems. Consider $H_2^{(3)}$ in [CCCMNNP], [Li], given by

$$H_2^{(3)} = \begin{pmatrix} 2 & -2 & -1 \\ -2 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}$$




FIGURE 13. Orbit of $\mathcal{H}(3)$

with Dynkin diagram as in Figure 14.

Its Dynkin diagram (Figure 14) contains a double edge and double arrow which, when removed, form a connected skeleton (Figure 15).




FIGURE 14. Dynkin diagram of type H_2^3




FIGURE 15. The connected skeleton $\mathcal{D}_*(H_2^3)$

Thus by Corollary 3.11 there is only one W -orbit, so we may expect that $\mathcal{P}(H_2^{(3)})$ will be a connected graph. Indeed this is the case, as shown in Figure 16. As we will prove in Section 8, this is one of only two rare examples of a hyperbolic root system on which the Weyl group acts both transitively and simply transitively.

Now consider the Feingold-Frenkel rank 3 hyperbolic root system,

$$A_1^{(1)\wedge} = \begin{pmatrix} 2 & -2 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

The Dynkin diagram for $A_1^{(1)\wedge}$ is given in Figure 18. The skeleton $\mathcal{D}_*(A_1^{(1)\wedge})$ is given in Figure 19.

which implies via Corollary 3.11 that the root system $\Phi(A_1^{(1)\wedge})$ is the disjoint union,

$$\Phi = W\alpha_1 \amalg W\{\alpha_2, \alpha_3\}.$$




FIGURE 16. The connected $\mathcal{P}$ -graph for hyperbolic root system $H_2^{(3)}$.

Remark 6.1. The vertex labelling in Figures 17 and 20 is encoded by the labelling of the 4-, 6-, and ∞ -gons of the graph. These polygons correspond to the defining relations of W , and the simple reflections that comprise these relations fix certain root projections according to Corollary 3.4. For instance, the 6-gons correspond to the relation $(w_2 w_3)^3 = 1$. Since w_2 and w_3 fix the root projection ρ_1 , the 1-projections of all 6 roots on a 6-gon are the same. Likewise, the 2-projections of all 4 roots on a




FIGURE 17. Connected component of $P(A_1^{(1)\wedge})$ corresponding to W_{α_1} . To decode the vertex labelling of the roots, see Remark 6.1.




FIGURE 18. Dynkin diagram of type $A_1^{(1)\wedge}$




FIGURE 19. The skeleton $\mathcal{D}_*(A_1^{(1)\wedge})$

4-gon, which corresponds to relation $(w_1 w_3)^3 = 1$, are the same. Thus, one may discern the 1-, 2-, and 3-projections of any root in these figures by reading the labels of the adjacent 6-, 4-, and ∞ -gons, respectively. In this way, given a root γ in these figures, the vertex labelling $(\rho_1(\gamma), \rho_2(\gamma), \rho_3(\gamma))$ is discovered.

Though the orbit graphs $\mathcal{P}$ for $A_1^{(1)\wedge}$ are drawn as if they appear in the Poincaré disk, there is no natural embedding of them in the disk. Since they are orbit graphs of real roots, they naturally live on a hyperboloid of 1 sheet which supports the real roots.

It is an interesting question to compare the combinatorial structure of the $\mathcal{P}$ -graph with the combinatorial structure of the action of the Weyl group W on the root lattice. Let w_i denote a set of generators for W . Then two real roots β_1 and β_2 are connected by a vertex in $\mathcal{P}$ if one is




FIGURE 20. Connected component of $P(A_1^{(1)\wedge})$ corresponding to $W\{\alpha_2, \alpha_3\}$. To decode the vertex labelling of the roots, see Remark 6.1.

transformed into the other by a generator w_i of W . However, the distance between β_1 and β_2 on the root lattice may be arbitrarily large.

Consider for example $A = A_1^{(1)}$ which is a rank 2 subsystem of $A_1^{(1)\wedge}$. Then $W = W(A)$ is the infinite dihedral group $W = \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$. Let β_0 be a real root and let $w_n = (w_1 w_2)^n \in W$ be some Weyl group element. Then $w_n(\beta_0)$ and $(w_2 w_n)(\beta_0)$ are adjacent in $\mathcal{P}$. On the other hand, in the root lattice, the Weyl distance $d((w_n \beta_0), (w_2 w_n) \beta_0)$ is proportional to n .

A similar phenomenon occurs for any hyperbolic Weyl group such that two of its generators w_i and w_j generate a sub-Weyl-group $W_{ij} = \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$.

Note that $\mathcal{P}(A_1^{(1)\wedge})$ shows the 6 roots in this root system which are fixed by simple reflections:

- (i) w_1 fixes $\pm\alpha_3$,
- (ii) w_2 fixes $\pm(\alpha_1 + 2\alpha_2 + 2\alpha_3)$,
- (iii) w_3 fixes $\pm\alpha_1$.

7. DIFFERENT ROOT LENGTHS IN DISTINCT ORBITS

Consider the following symmetric generalized Cartan matrix of noncompact hyperbolic type:

$$A = \begin{pmatrix} 2 & -2 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$$

and let

$$A' = \begin{pmatrix} 2 & -1 & 0 \\ -4 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

Then A' is non symmetric. Also A and A' have different Dynkin diagrams but the same Weyl group $W \cong PGL_2(\mathbb{Z})$ and the same Coxeter matrix

$$\begin{pmatrix} 1 & \infty & 2 \\ \infty & 1 & 3 \\ 2 & 3 & 1 \end{pmatrix}.$$

Let Φ be the real roots of A and Φ' the real roots of A' . The 2 different root lengths in A' justify the 2 distinct W -orbits of real roots in $W\Phi'$ which also occur in $W\Phi$. This is a general phenomenon. We will prove however that this can only occur if A has rank 2 affine type, or A has rank 3 noncompact hyperbolic type. We will make use of the following result from ([CCCMNNP]).

Proposition 7.1. ([CCCMNNP]) *A symmetrizable hyperbolic generalized Cartan matrix contains an $A_1^{(1)}$ or $A_2^{(2)}$ proper indecomposable submatrix if and only if rank $A = 3$ and A has noncompact type.*

Theorem 7.2. *Given a symmetric generalized Cartan matrix A of noncompact hyperbolic type such that real roots of the same length lie in distinct orbits, there is a nonsymmetric generalized Cartan matrix A' with the same Weyl group as A and the same Coxeter matrix as A , such that the roots of different root lengths lie in distinct W -orbits.*

Proof: Let Φ be the real roots of A and Φ' the real roots of A' . If real roots in Φ of the same root lengths lie in distinct W -orbits, then in the skeleton of the Dynkin diagram $\mathcal{D}(A)$, there must be 2 disconnected vertices, say i and j . Hence between vertices i and j in the Dynkin diagram, there must have been ≥ 2 edges. Hence $a_{ij}a_{ji} \geq 4$. Since A has noncompact hyperbolic type, it follows that A contains an $A_1^{(1)} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$ proper indecomposable submatrix.

Applying Proposition 7.1, we see that A must have rank 3 noncompact hyperbolic type. Next we claim that for every Dynkin diagram of rank 3 noncompact hyperbolic type with an $A_1^{(1)}$ subdiagram, there is another Dynkin diagram of rank 3 noncompact hyperbolic type with an $A_2^{(2)} = \begin{pmatrix} 2 & -1 \\ -4 & 2 \end{pmatrix}$ subdiagram. This is easily checked in the tables in the classification of hyperbolic Dynkin diagrams in ([CCCMNNP]). $\square$

It follows that there is a natural bijective correspondence between the orbit structure $W\Phi'$ of the real roots corresponding to A' and the orbit structure $W\Phi$ of the real roots corresponding to A .

From [CCCMNNP], if A_0 is a rank 2 proper submatrix of a symmetrizable hyperbolic generalized Cartan matrix A of rank ≥ 4 then A_0 is of finite type. We have the following corollary which follows easily from the tables in the classification of hyperbolic Dynkin diagrams in ([CCCMNNP]).

Corollary 7.3. *Let A symmetric hyperbolic generalized Cartan matrix of rank ≥ 4 or symmetrizable and rank ≥ 7 . Then any 2 real roots of the same length lie in the same W -orbit.*

If A has rank 4, 5 or 6 and is not symmetric, then the corollary is false.

Example. The generalized Cartan matrix of the Dynkin diagram of rank 4 noncompact hyperbolic type (Figure 21) is symmetrizable but not symmetric. There are 3 distinct lengths of real roots, but 4 W -orbits on real roots. In particular, α_1 and α_3 have the same root length but lie in different W -orbits.




FIGURE 21. Dynkin diagram of type $H_{23}^{(4)}$

8. FIXED POINTS AND ORBIFOLD GRAPHS ASSOCIATED TO THE GRAPH $\mathcal{P}$

In this section we give sufficient conditions in terms of the generalized Cartan matrix A (equivalently $\mathcal{D}$) for a simple reflection in W to stabilize a simple root. We thus obtain necessary conditions for the action of W to be simply transitive, and show that the Weyl group for E_{10} acts transitively but not simply transitively on real roots. We also obtain sufficient conditions for a simple reflection to stabilize a real root that is not simple.

We begin by recalling the notion of complete graph and we apply the definition to Dynkin diagrams.

Definition 8.1. *A Dynkin diagram $\mathcal{D}$ is called **complete** if it is a complete graph, that is, if every pair of distinct vertices $\mathcal{D}$ is connected by an edge.*

We now establish sufficient conditions for the existence of fixed roots.

Lemma 8.2. *Let A be a Cartan matrix (or generalized Cartan matrix), $\mathcal{D}$ its corresponding Dynkin diagram, Φ the set of real roots with basis Π indexed by I , and W the Weyl group. If A contains*

zeros (equivalently, if $\mathcal{D}$ is not complete), then there exist simple roots in Φ that are stabilized by simple reflections in W , and thus W does not act simply transitively on Φ .

Proof. We first verify the equivalency in the hypothesis, namely, that if A contains a zero then $\mathcal{D}$ is not a complete graph. Assume $a_{ij} = 0$. By the definition of a Cartan matrix, this implies that $a_{ji} = 0$. Thus the submatrix $A_{(i,j)}$ corresponds to the Cartan matrix for $A_1 \times A_1$:

$$A_{(i,j)} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix},$$

and this implies vertices i, j in $\mathcal{D}$ are not connected by an edge. Thus $\mathcal{D}$ is not complete.

We now prove the existence of fixed simple roots by observing the action of simple roots w_i and w_j on simple roots α_i and α_j :

$$w_i \alpha_j = \alpha_j - a_{ji} \alpha_i = \alpha_j,$$

and

$$w_j \alpha_i = \alpha_i - a_{ij} \alpha_j = \alpha_i.$$

Thus w_j fixes α_k , and w_k fixes α_j . This is illustrated by the $\mathcal{P}$ -graph for $A_1 \times A_1$ in Figure 22.




FIGURE 22. Orbit structure of $A_1 \times A_1$

The existence of roots fixed by W proves that W does not act simply transitively on Φ . □

Restating the lemma in its contrapositive form reveals the necessary conditions for a Weyl group to act simply transitively.

Corollary 8.3. *If a Weyl group W acts simply transitively on its corresponding root system Φ , then its associated Dynkin diagram is complete.*

Since the Dynkin diagrams for E_{10} and $A_1^{(1)\wedge}$ are not complete, we have the following immediate consequence of Corollary 3.11 and Lemma 8.2.

Corollary 8.4. *The Weyl group for E_{10} acts transitively on its real roots, but not simply transitively. The Weyl group of $A_1^{(1)\wedge}$ is neither transitive nor simply transitive on its real roots.*

These results raise the question, how many hyperbolic root systems have Weyl groups that act both transitively and simply transitively? In the following corollary, we prove that there are only 2.

Corollary 8.5. *Let $\mathcal{D}$ be a Dynkin diagram of hyperbolic type, A its corresponding generalized Cartan matrix, Φ its corresponding root system and W its Weyl group. Assume that W acts both transitively and simply transitively on Φ . Then $\mathcal{D}$ is either $H_2^{(3)}$ or $H_1^{(4)}$ in the classification ([CCCMNNP], [Li], [KM], [Sa]).*

Proof. Since W acts simply transitively on Φ , by Lemma 8.2 $\mathcal{D}$ must be complete. Since W also acts transitively on Φ , $\mathcal{D}_*$ is connected by Corollary 3.11. Thus Φ contains roots of only 1 length, and A is symmetric.

Case 1: $\mathcal{D}$ is comprised of 2 vertices. Thus,

$$A = \begin{pmatrix} 2 & -a \\ -a & 2 \end{pmatrix},$$

where $a \geq 3$. But then $\mathcal{D}$ contains a multiple (bold) edge, which is removed when forming $\mathcal{D}_*$, which is thus disconnected. Hence W does not act transitively on Φ by Corollary 3.11, which is a contradiction.

Case 2: $\mathcal{D}$ is comprised of ≥ 3 vertices. The classification of hyperbolic dynkin diagrams of ranks 3 – 10 (maximal) contains 142 symmetrizable diagrams, of which only 17 diagrams are complete. Of these diagrams, only $H_2^{(3)}$ and $H_1^{(4)}$ have connected skeletons $\mathcal{D}_*$. $\square$

Thus Figure 16 shows the rare case of a hyperbolic root system that consists of only 1 orbit and has no fixed roots.

Note that Lemma 8.2 applies to both finite and infinite root systems. We now reveal a consequence of the lemma for the finite dimensional theory.

Corollary 8.6. *Let $\mathfrak{g}$ be a finite dimensional semisimple Lie algebra of rank ≥ 3 . Then its Weyl group W does not act simply transitively on its corresponding root system.*

Proof. The corresponding Dynkin diagram contains 3 or more vertices. Searching through the classification of finite dimensional simple Lie algebras reveals that $\mathcal{D}$ is not complete, and the result follows. $\square$

Lemma 8.2 showed sufficient conditions for a simple reflection in W to stabilize a simple root. In the following theorem, we provide a more general condition for a simple reflection to stabilize any root, not necessarily simple.

Theorem 8.7. *Let A be a generalized Cartan matrix, and let $W(A)$ and $\Phi(A)$ be its associated Weyl group and set of real roots. Let $A_{(i)}$ denote the i -th column vector of A ,*

$$A_{(i)} = \begin{bmatrix} A_{1i} \\ A_{2i} \\ \vdots \\ A_{li} \end{bmatrix}.$$

Let $\alpha \in \Phi$. If $\alpha \cdot A_{(i)} = 0$, then α is fixed by w_i .

Proof.

$$w_i \alpha = \alpha - (\alpha \cdot A_{(i)}) \alpha_i = \alpha.$$

$\square$

Recall from Section 6 that the hyperbolic root system of $A_1^{(1)\wedge}$ contains 6 fixed roots. Each of these roots is fixed by a simple Weyl reflection, e.g., $w_1 \in \text{Stab}_W(\alpha_3)$ and $w_3 \in \text{Stab}_W(\alpha_1)$. We now further demonstrate that these simple reflections constitute the only nontrivial stabilizers of these roots.

Theorem 8.8. *Consider the fixed points $\pm\alpha_1$, $\pm\alpha_3$, and $\pm(\alpha_1 + 2\alpha_2 + 2\alpha_3)$ in the hyperbolic root system $A_1^{(1)\wedge}$. We have*

- (i) $\text{Stab}_W(\alpha_1) = \langle w_3 \rangle$,
- (ii) $\text{Stab}_W(\alpha_3) = \langle w_1 \rangle$,
- (iii) $\text{Stab}_W(\alpha_1 + 2\alpha_2 + 2\alpha_3) = \langle w_2 \rangle$.

Proof. The reverse inclusions have already been established in Section 6. The following general argument proves the forward inclusions. Let β be a fixed point in $A_1^{(1)\wedge}$. Let $w \in W$ be a stabilizer of β , and assume w is neither the identity nor a simple Weyl reflection. Then w corresponds to a closed path in the $\mathcal{P}$ -graph for $A_1^{(1)\wedge}$ beginning and ending at β . We now label paths in $\mathcal{P}$ by the corresponding Weyl element, extending the notion of labelling single edges by the corresponding simple Weyl reflection. Thus we refer to this closed path also as w .

Case 1: w (as a path) does not contain any fixed points other than β . Then w reduces to the identity since it is a path in the Cayley graph, which is a contradiction.

Case 2: w (as a path) contains a fixed point $\alpha \neq \beta$. Then this path is of the form $w = vhu$, where $h \in \text{Stab}_W(\alpha)$ and $u(\beta) = \alpha$ and $v(\alpha) = \beta$, as shown in Figure 23.




FIGURE 23. $w = vhu$

Since u, v are paths in the Cayley graph, they reduce via the relations of W to some Weyl word g and its inverse. Thus $w = g^{-1}hg$, as illustrated in Figure 24.




FIGURE 24. $w = g^{-1}hg$

However, if we consider this type of Weyl element to be a nontrivial stabilizer of β , then *all* roots in Φ are trivially stabilized by Weyl elements of this type. This would make every root a

fixed root, which is impossible since the action is non-trivial. Thus Weyl elements of the type $g^{-1} \text{Stab}_W(\alpha)g$ are not admissible as stabilizers, and we have another contradiction.

Thus all stabilizers in $A_1^{(1)\wedge}$ must be either trivial or simple reflections, giving us the forward inclusions. □

Thus, given any hyperbolic root system, Theorem 8.7 provides a geometric approach to finding all fixed points, as the stabilizer of any given root contains only simple reflections and the identity.

We may use the example of the Feingold-Frenkel rank 3 hyperbolic root system and its graph $\mathcal{P}(A_1^{(1)\wedge})$ to indicate the role of fixed points. Here, W is the $(\infty, 3, 2)$ triangle group which acts naturally on the Poincaré disk. If W acted simply transitively on the Poincaré disk, the graph $\mathcal{P}(A_1^{(1)\wedge})$ would be a single connected component and would coincide with the Cayley graph, dual to the tessellation. Note that this is the case for $\mathcal{P}(H_2^{(3)})$ (Figure 16), which coincides with the Cayley graph for the triangle group $(\infty, 3, 3)$, dual to the tessellation. However, for $A_1^{(1)\wedge}$ there are 2 orbits for W on real roots, hence 2 connected components of the graph $\mathcal{P}(A_1^{(1)\wedge})$. If there were no fixed points for W on real roots, these 2 components would both be dual to the tessellation. However, each connected component admits distinct fixed points for the action of W on simple roots. The connected components of $\mathcal{P}(A_1^{(1)\wedge})$ are thus orbifold graphs. For transitive Weyl group actions, the graph $\mathcal{P}$ is dual to the W -tessellation of its root space.

9. WEYL GROUP ORBITS ON IMAGINARY ROOTS

In this section we examine the structure of the orbits of the Weyl group on imaginary roots. We use the symmetry properties of the lightcone in the Cartan subalgebra of a hyperbolic Kac–Moody algebra to deduce some natural properties of the structure of Weyl group orbits on imaginary roots.

Let $\mathfrak{g}$ be a Kac–Moody algebra with Cartan subalgebra $\mathfrak{h}$, root space decomposition

$$\begin{aligned} \mathfrak{g} &= \mathfrak{g}^+ \oplus \mathfrak{h} \oplus \mathfrak{g}^-, \\ \mathfrak{g}^+ &= \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}^\alpha, \quad \mathfrak{g}^- = \bigoplus_{\alpha \in \Delta^-} \mathfrak{g}^\alpha \end{aligned}$$

and root spaces

$$\mathfrak{g}^\alpha = \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x, h \in \mathfrak{h}\}.$$

The dimension of the root space $\mathfrak{g}^\alpha$ is called the *multiplicity* of α . We recall that the associated Weyl group acts on the set Δ of all roots, preserving root multiplicities.

Now let $\mathfrak{h}$ be the standard Cartan subalgebra of a symmetrizable hyperbolic Kac–Moody algebra $\mathfrak{g}$ and let $\mathfrak{h}_{\mathbb{R}} \subset \mathfrak{h}$ such that $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{R}} \mathfrak{h}_{\mathbb{R}}$ and $\mathfrak{h}_{\mathbb{R}}$ contains the simple roots of $\mathfrak{h}$. Then the Weyl group has a fundamental chamber $\mathcal{C}$ inside $\mathfrak{h}_{\mathbb{R}}$. The union X of the sets $w(\mathcal{C})$, for $w \in W$

$$X = \bigcup_{w \in W} w(\mathcal{C})$$

is called the Tits cone.

When $\mathfrak{g}$ is of hyperbolic type, we have the following description of the union of the closures of the positive and negative Tits cones

$$\overline{X} \cup -\overline{X} = \{h \in \mathfrak{h}_{\mathbb{R}} \mid (h \mid h) \leq 0\},$$

where $(\cdot \mid \cdot)$ is the symmetric bilinear invariant form on $\mathfrak{g}$. The set

$$\mathcal{L}_{\mathfrak{h}_{\mathbb{R}}} = \{h \in \mathfrak{h}_{\mathbb{R}} \mid (h \mid h) \leq 0\}$$

is called the *lightcone* of the Cartan subalgebra $\mathfrak{h}_{\mathbb{R}}$.

As the Adjoint action (and W) act by isometries, the length of roots is preserved. As a consequence the action of W preserves the set of imaginary roots of zero squared length and of negative squared length.

The imaginary roots of negative squared length lie on hyperboloids of a fixed ‘radius’ inside the lightcone while the imaginary roots of zero squared length lie on the boundary of the lightcone. Moreover, every lattice point inside the lightcone of $\mathfrak{h}$ corresponds to an imaginary root.

Lemma 9.1. *Let $\mathfrak{g}$ be a Kac–Moody algebra. Let α be an imaginary root. Then W acts as ‘translations’ on the hyperboloid of radius $(\alpha \mid \alpha)$. In particular, there is no $w \in W$ such that $w\alpha = -\alpha$.*

Proof: We may assume that α is a positive imaginary root. Then W preserves root length and by Lemma 5.1, $W\alpha$ is positive. It follows that $W\alpha$ is on the ‘sphere’ of radius $(\alpha \mid \alpha)$. Hence there is no $w \in W$ such that $w\alpha = -\alpha$. We call the image $W\alpha$ of α a ‘translation’ along the ‘sphere’ of radius $(\alpha \mid \alpha)$. $\square$

Lemmas 5.1 and 9.1 give us an obvious but useful characterization of real and imaginary roots in terms of the Weyl group orbits.

Corollary 9.2. *Let $\mathfrak{g}$ be a symmetrizable Lie algebra or Kac–Moody algebra. Let α be any root. Then*

$$\alpha \text{ is real if and only if there exists } w \in W \text{ such that } w\alpha = -\alpha,$$

$$\alpha \text{ is imaginary if and only if there does not exist } w \in W \text{ such that } w\alpha = -\alpha.$$

Focusing the attention to sets of roots that lie on the same hyperboloid of fixed squared length, it is possible that these sets contain roots of different multiplicities. If $\alpha, \beta \in \Delta^{imag}$ are on the same hyperboloid of fixed squared length but $mult(\alpha) \neq mult(\beta)$, then there is no element of $Aut(\mathfrak{g})$ that takes α to β . In particular there is no element of W that takes α to β . Hence roots with different multiplicity must lie in different W -orbits.

We make the following self-evident but important observation. Let $\mathfrak{g}$ be a hyperbolic Kac–Moody algebra. Let α and β be any pair of imaginary roots of the same squared length that lie inside the fundamental chamber $\mathcal{F}$ for the Weyl group or on its boundary $\partial\mathcal{F}$. Then α and β may have the same multiplicity but lie in distinct orbits under the action of W .

It follows that imaginary roots are in the same W -orbit, then they must have the same squared length and the same multiplicity but can’t both be in the fundamental chamber for the Weyl group or on its boundary.

Theorem 9.3. *Let $\mathfrak{g}$ be a Kac–Moody algebra of hyperbolic type.*

- (1) *For imaginary roots of fixed squared length on a hyperboloid $\mathcal{H}$, there are finitely many W –orbits of imaginary roots.*
- (2) *The cardinality of each such orbit is infinite.*
- (3) *The number of W –orbits on imaginary roots of fixed squared length is bounded above by the number of lattice points in $(\mathcal{F} \cup \partial\mathcal{F}) \cap \mathcal{H}$.*
- (4) *The number of W –orbits on imaginary roots $\rightarrow \infty$ as squared length $\rightarrow \infty$.*

Proof: Let $\mathcal{H}$ be a hyperboloid supporting the imaginary roots of some fixed squared length. Let S be the set of lattice points in $(\mathcal{F} \cup \partial\mathcal{F}) \cap \mathcal{H}$. Then $|S| < \infty$. The imaginary roots corresponding to lattice points in S cannot be in the same W –orbit. The roots in S may or may not have the same multiplicity. If they have the same multiplicity, they may be in the same W –orbit. If the roots in S do not all have the same multiplicity, then they will fall into finitely many W –orbits. Thus WS is a union of finitely many infinite disjoint sets. This proves (1) and (2) and (3) and (4) follow easily. $\square$

REFERENCES

- [ACP] Andersen, K. K. S., Carbone, L. and Penta, D. *Kac–Moody Fibonacci sequences, hyperbolic golden ratios, and real quadratic fields*, Journal of Number Theory and Combinatorics, Vol 2, No. 3, 245–278 (2011)
- [BGH] Brown, J., Ganor, O. and Helfgott, C., *M-theory and E_{10} : Billiards, Branes, and Imaginary Roots*, J. High Energy Phys. (2004), no. 8, 063, 55 pp. (electronic)
- [CCCMNNP] Carbone, L., Chung, S., Cobbs, L., McRae, R., Nandi, D., Naqvi Y. and Penta, D. *Classification of hyperbolic Dynkin diagrams, root lengths and Weyl group orbits*, Phys. A: Math. Theor. 43 155209 (2010)
- [Ca] Carter, R. W. *Conjugacy classes in the Weyl group*, Compositio Math. 25 (1972), 1–59
- [DHN] Damour, T., Henneaux, M. and Nicolai, H., *Cosmological Billiards*, hep-th/0212256, Classical Quantum Gravity 20 (2003), no. 9, R145–R200
- [EHTW] Englert, F. Houart, L., Taormina, A. and West, P., *The Symmetry of M-Theories*, J. High Energy Phys. 2003, no. 9, 020, 39 pp. (electronic)
- [FF] Feingold, A and Frenkel, I *A hyperbolic Kac- Moody algebra and the theory of Siegel modular forms of genus 2*, Math. Ann. 263, no. 1, 87–144, 1983
- [Ka] Victor G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990.
- [KM] Kobayashi, Z. and Morita, J., *Automorphisms of certain root lattices*, Tsukuba J. Math., 7, 323–336 (1983)
- [Li] Li, W., *Generalized Cartan matrices of hyperbolic type*, Chinese Annals of Mathematics, 9B, (1988), 68–77
- [M1] Moody, R. V. , *Lie algebras associated with generalized Cartan matrices*, Bull. Amer. Math. Soc., 73 (1967) 217-221
- [M2] Moody, R. V. *A new class of Lie algebras*, J. Algebra 10 (1968) 211–230
- [Sa] Sağlıoğlu, C., *Dynkin diagrams for hyperbolic Kac–Moody algebras*, Journal of Physics A: Mathematics and General 22 3753–3769, (1989)
- [T] Tits, J., *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Mathematics, Vol. 386. Springer-Verlag, Berlin-New York, (1974). x+299 pp

DEPARTMENT OF MATHEMATICS, HILL CENTER, BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN RD, PISCATAWAY, NJ 08854-8019

E-mail address: carbonel@math.rutgers.edu

FACHBEREICH MATHEMATIK, TU DARMSTADT, SCHLOSSGARTENSTRASSE 7, 64289 DARMSTADT, GERMANY

E-mail address: walter.freyn@math.tu-darmstadt.de

DEPARTMENT OF MATHEMATICAL SCIENCES, BINGHAMTON UNIVERSITY, BINGHAMTON, NEW YORK 13902-6000

E-mail address: diegopenta@gmail.com