

# AUGMENTATION VARIETIES AND DISK POTENTIALS I

KENNETH BLAKEY, SOHAM CHANDA, YUHAN SUN, AND CHRIS WOODWARD

ABSTRACT. This is the first in a sequence of papers where we develop a field theory for Lagrangian cobordisms between Legendrians in circle-fibered contact manifolds. In particular, we show that Lagrangian fillings such as the Harvey-Lawson filling in any dimension define augmentations of Chekanov-Eliashberg differential graded algebras by counting configurations of holomorphic disks connected by gradient trajectories, as in Aganagic-Ekholm-Ng-Vafa [2]; we also prove that for Legendrian lifts of monotone tori, the augmentation variety is the zero level set of the Landau-Ginzburg potential of the Lagrangian projection, as suggested by Dimitroglou-Rizell-Golovko [22]. In this part, we set up the analytical foundation of moduli spaces of pseudoholomorphic buildings.

## CONTENTS

1. Introduction	2
2. Fibered contact manifolds and Lagrangian cobordisms	5
2.1. Fibered contact manifolds and Legendrians	5
2.2. Lagrangian cobordisms	11
2.3. Reductions of symplectic fillings	19
3. Pseudoholomorphic buildings	22
3.1. Punctured surfaces	23
3.2. Pseudoholomorphic buildings	24
3.3. Treed pseudoholomorphic buildings	26
3.4. Monotonicity assumptions	32
3.5. Homological invariants	37
4. Regularization	39
4.1. Donaldson hypersurfaces	40
4.2. Fredholm theory	45
4.3. Transversality	49
4.4. Intersection multiplicities	56
4.5. Compactness	56
4.6. Orientations	60
References	62

---

Chanda and Woodward were partially supported by NSF grant DMS 2105417 and Sun was partially supported by the EPSRC grant EP/W015889/1. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

## 1. INTRODUCTION

Contact homology as introduced by Eliashberg, Givental, and Hofer [31] is a theory whose generators come from closed Reeb orbits and whose differential counts holomorphic curves in the symplectization of a contact manifold. Its relative version, Legendrian contact homology, was introduced by Chekanov [15] and Eliashberg [30], and has been developed by Ekholm-Etnyre-Ng-Sullivan [28] and related works for the case that the contact manifold fibers over an exact symplectic manifold. This is the first of a sequence of papers in which we develop a version of Legendrian contact homology in case that the contact manifold fibers over a monotone symplectic manifold, extending work of Sabloff [45], Asplund [4], and Dimitroglou-Rizell-Golovko [22]. In particular, we give a definition of augmentation variety of Legendrians such as the lift of a Clifford torus in any dimension.

Our definition of the Legendrian contact homology is related to the work of Aganagic-Ekholm-Ng-Vafa [2]. An important theme in [2] is the relation between *Lagrangian fillings* of a Legendrian and *augmentations* of its Chekanov-Eliashberg algebra. The standard definition of a Lagrangian filling assumes a cylindrical end modelled on the product of the real line with the Legendrian. However, in important examples such as the Harvey-Lawson filling considered here, the Lagrangian is asymptotic to the cylinder, rather than equal to it outside of a compact set. Moreover, the Harvey-Lawson filling is not an exact Lagrangian submanifold, which means it may bound non-trivial holomorphic disks. Therefore Aganagic-Ekholm-Ng-Vafa [2] suggest that in order for such Lagrangian fillings to define augmentations of Chekanov-Eliashberg differential graded algebra, one should count configurations of holomorphic disks connected by gradient trajectories of certain vector fields. Such configurations are otherwise known as treed disks or clusters, in the language of Cornea-Lalonde [20]. By counting such configurations, we give a construction of the Chekanov-Eliashberg dga in which the zeroes of the corresponding vector fields as generators, similar to the definition of immersed Lagrangian Floer homology developed by Akaho-Joyce [3]. The resulting version of the Chekanov-Eliashberg algebra gives rich Legendrian isotopy invariants, which distinguish many Legendrian lifts of Lagrangians in monotone symplectic manifolds such as Vianna's exotic tori; the non-isotopy of these tori was conjectured by Dimitroglou-Rizell-Golovko [22].

The use of the cluster approach might seem like a technical choice, but actually it resolves an obstruction to the differential squaring to zero, as already noted in Aganagic-Ekholm-Ng-Vafa [2]. Namely, in even the basic examples of Lagrangian fillings one has disk bubbling in the one-dimensional moduli spaces of holomorphic disks bounding the filling. In the cluster approach, the moduli space "continues" as a moduli space of configurations where two disks are separated by a gradient line, which eventually breaks to give a true boundary configuration. In this broken configuration, there are two levels, in the sense of symplectic field theory, meaning that this configuration is again a contribution to the square of the differential.

In this first paper in the series, we construct the moduli spaces of treed disks with punctures, and prove results about their compactness, regularization and orientations. In sequels [7, 8], these moduli spaces will be used to:

- (a) Define the Chekanov-Eliashberg algebra, and show the resulting Legendrian contact homology is independent of various choices made;
- (b) For a tame Lagrangian cobordism between Legendrians, see Definition 3.19 below, define a chain map between the Chekanov-Eliashberg algebras. Particularly, define an augmentation variety for a Legendrian and show it is a Legendrian isotopy invariant.
- (c) Establish a concrete relation between the disk potential functions of certain monotone Lagrangians such as Vianna's tori and the augmentation varieties of their Legendrian lifts, hence distinguish infinitely many Legendrians that are mutually not Legendrian isotopic.
- (d) Compute the augmentation varieties for the Clifford-type and Hopf-type Legendrians, and the augmentation variety in particular for disconnected Legendrians, in relation to the conjectures of [2] on the parametrization of components by partitions.

Now we describe our geometric settings and moduli spaces in a more detailed way. Consider a Legendrian  $\Lambda$  in a compact contact manifold  $Z$  that is a negatively-curved circle-fibration over a symplectic base  $Y$ ; we have in mind especially the example that  $Z$  is some unit complex line bundle over a Fano projective toric variety  $Y$ , and  $\Lambda$  is the horizontal lift of a Lagrangian in the base. The contact manifold is equipped with the Boothby-Wang contact form. One fundamental example as in Dimitroglou-Rizell-Golovko in [22] is the horizontal lift of the Clifford torus

$$\Pi_{\text{Cliff}} = \left\{ [z_1, \dots, z_n] \mid \sum_{i=1}^n |z_i|^2 = 1 \right\} \cong (S^1)^{n-1} \subset \mathbb{C}P^{n-1}$$

in complex projective space  $Y = \mathbb{C}P^{n-1}$ . Its horizontal lift to the unit tautological line bundle is a Legendrian torus in the standard contact sphere  $Z = S^{2n-1}$ :

$$(1) \quad \Lambda_{\text{Cliff}} = \left\{ (z_1, \dots, z_n) \in \mathbb{C}^n \mid \begin{array}{l} |z_1|^2 = \dots = |z_n|^2 = \frac{1}{n} \\ z_1 \dots z_n \in (0, \infty) \end{array} \right\} \cong (S^1)^{n-1} \subset S^{2n-1}.$$

The map

$$\mathbb{C}^n - \{0\} \rightarrow \mathbb{C}P^{n-1}, \quad z \mapsto \text{span}(z)$$

restricts an  $n$ -fold cover of  $\Pi_{\text{Cliff}}$ . Similar examples are given by taking  $\Pi$  to be a Lagrangian in a closed monotone symplectic manifold  $Y$ ,  $Z \rightarrow Y$  a circle bundle with connection with rational holonomies and  $\Lambda$  the horizontal lift of  $\Pi$ .

An example of a Lagrangian filling is given by the Harvey-Lawson filling of the Legendrian in the previous paragraph. Let

$$a_1, \dots, a_n \in \mathbb{R}$$

be real numbers with exactly two zero. Define as in Joyce [38, (37)]

$$(2) \quad L_{(1)} = \left\{ (z_1, \dots, z_n) \in \mathbb{C}^n \mid \begin{array}{l} |z_1|^2 - a_1^2 = |z_2|^2 - a_2^2 = \dots = |z_n|^2 - a_n^2 \\ z_1 \dots z_n \in [0, \infty) \end{array} \right\}.$$

A picture of the moment image of this filling is shown in Figure 4 below; the intersections with toric moment fibers are generically tori of codimension one. To

understand this *Harvey-Lawson filling* better, consider the map

$$\pi : \mathbb{C}^n \rightarrow \mathbb{C}, \quad \pi(z_1, \dots, z_n) = z_1 z_2 \dots z_n$$

which is a symplectic fibration away from the critical locus. The symplectic horizontal bundle of the fibration  $\pi$  is the symplectic complement of the vertical bundle. A direct computation shows that the symplectic horizontal bundle  $H_z$  at a point  $z \in \mathbb{C}^n$  is given as

$$H_z = \text{Span}_{\mathbb{C}} \left\{ \left( \frac{z_1}{|z_1|^2}, \dots, \frac{z_n}{|z_n|^2} \right) \right\}.$$

Consider the torus

$$T_1 = \{z \in \pi^{-1}(1) \mid |z_1|^2 - a_1^2 = |z_2|^2 - a_2^2 = \dots = |z_n|^2 - a_n^2\}.$$

Since the derivative of  $|z_i|^2 - |z_j|^2$  along any vector in  $H_z$  vanishes, we conclude that the parallel transport of  $T_1$  along  $[0, \infty)$  lies in  $L_{(1)}$ . Thus  $L_{(1)}$  can be viewed as the union of parallel transports of  $T_1$  along  $[0, \infty)$ .

We briefly describe the moduli spaces of pseudoholomorphic curves needed for the definition of the differential associated to the Legendrian and the chain maps associated to Lagrangian cobordisms. The construction is a special case of symplectic field theory with Lagrangian boundary conditions, where the contact manifold is circle-fibered. The simplifications in this case are comparable to that of genus-zero relative Gromov-Witten theory compared to full symplectic field theory with contact boundary. To orient the reader, we list here two technical parts of our construction.

- (a) The Boothby-Wang contact form on the circle-fibered contact manifold is of Morse-Bott non-degenerate type. We don't perturb it and we introduce a Morse model for the Chekanov-Eliashberg algebra.
- (b) We consider more general tame Lagrangian cobordisms which may bound non-trivial holomorphic disks, to define chain maps.

Now we explain these two points in more details. By a *punctured disk* we mean a surface obtained by removing a finite collection of points on the boundary. More precisely our configurations are *treed punctured disks* obtained by adding a finite collection of trajectories of some vector fields, on both the space of Reeb chords and the Legendrian itself, as in Figure 1, and to be explained below. In Figure 1, the dotted lines represent Reeb chords, and the trajectories can either connect the Reeb chords or connect points on the Lagrangian boundary condition of the disks. The disk shown would contribute to the coefficient of a word of length eight in the differential applied to a Reeb chord. Note that disk components without punctures (of which one is shown in Figure 1) appear in the configurations defining the chain maps associated to cobordisms, but not in the configurations defining the differential on our Chekanov-Eliashberg algebras as we always assume that our one-forms on the boundary are contact and so non-constant holomorphic disks have at least one puncture, see Corollary 3.17 below. We assume the existence of suitable bounding cochains, as in the theory of  $A_\infty$  algebras, so that disk bubbling in our cobordisms does not prevent the counts from defining chain maps. The strata corresponding to disk bubbling are in the interior of the one-dimensional moduli spaces; instead the

true boundary components arise from breaking of edges of the segments connecting the disks. We use Cieliebak-Mohnke [17] perturbations to regularize the moduli spaces. The natural chain complex  $CE(\Lambda)$  is generated by words in chains on the




FIGURE 1. A treed punctured disk

Reeb chords and chains on the Legendrian; the addition of these *classical generators* does not occur in the version of Legendrian contact homology over exact symplectic manifolds in Ekholm-Etnyre-Sullivan [27]. The generators arising from chains on the Legendrian arise because of nodes developing along points in the interior of the cobordism, which then lead to trajectories as in Ekholm-Ng [29]; in particular [29] assigns augmentations to the Harvey-Lawson filling in the three-dimensional setting of knot contact homology while the construction here is intended to work in all dimensions, under some restrictions. In particular, the constructions here do apply to the fillings appearing in knot contact homology. It seems likely that in the cases of overlap the two theories agree, as what we develop here is essentially a Morse-Bott version of that theory, but the relationship remains to be determined.

The current paper first (Section 2) introduces geometric constructions of many circle-fibered contact manifolds and their Legendrians. In particular, we construct Lagrangian fillings of certain Legendrian lifts of Vianna's tori, which may be of independent interest. Then we set up moduli spaces of treed punctured disks solving the Cauchy-Riemann equation (Section 3), and prove their compactness, regularity and orientability (Section 4).

## 2. FIBERED CONTACT MANIFOLDS AND LAGRANGIAN COBORDISMS

In this section, we describe various constructions of Legendrians in circle-fibered contact manifolds and Lagrangian cobordisms. The Harvey-Lawson filling is described both as a (non-exact) asymptotic filling and as a compact filling of a Legendrian with respect to a perturbation of the standard contact structure on the sphere.

**2.1. Fibered contact manifolds and Legendrians.** The contact manifolds we consider are circle fibrations over a compact symplectic manifold. Let  $n$  be a positive

integer and  $Z$  a smooth manifold of dimension  $2n - 1$ . Let

$$\Omega(Z) = \bigoplus_{j=0}^{\dim(Z)} \Omega^j(Z)$$

be the algebra of smooth differential forms. Recall, as in for example Geiges [34], the following definitions.

**Definition 2.1.** Let  $Z$  be a  $2n - 1$ -manifold. A *contact form* on  $Z$  is a one-form

$$\alpha \in \Omega^1(Z), \quad (\alpha \wedge (d\alpha)^{n-1})(z) \neq 0 \quad \forall z \in Z.$$

The kernel of  $\alpha$  is the *contact distribution*

$$\xi := \ker(\alpha) \subset TZ$$

associated to  $\alpha$ . The form  $\alpha$  defining  $\xi$  is unique up to multiplication by a non-zero function.

A *Legendrian submanifold* of  $Z$  is a submanifold  $\iota : \Lambda \subset Z$  of dimension  $n - 1$  on which  $\alpha$  vanishes:

$$\iota^* \alpha = 0 \in \Omega^1(Z).$$

We will assume our submanifolds are embedded, rather than immersed, unless otherwise stated.

More generally, as in Cieliebak-Volkov [18], a *stable Hamiltonian structure* on  $Z$  is a pair of a two-form and a one-form

$$\omega \in \Omega^2(Z), \quad \alpha \in \Omega^1(Z)$$

such that

$$\alpha \wedge \omega^{n-1} \neq 0, \quad \ker(\omega) \subset \ker(d\alpha) \subset TZ.$$

A manifold with a stable Hamiltonian structure will be called a *stable Hamiltonian manifold*.

A *Legendrian submanifold* of a stable Hamiltonian manifold is a maximal isotropic submanifold for  $\omega$  on which  $\alpha$  vanishes. If  $\omega = -d\alpha$  then the pair  $(Z, \alpha)$  is a contact manifold.

Given a manifold  $Z$  with an action of the circle group  $S^1$ , let

$$\partial_\theta \in \text{Vect}(S^1), \quad S^1 = \mathbb{R}/\mathbb{Z}$$

be the rotationally invariant vector field with period 1. A *connection one-form*  $\alpha$  is a one-form invariant under the  $S^1$ -action on  $Z$  and the contraction  $\alpha(\partial_\theta) \in \Omega^0(Z)$  of the generating vector field  $\partial_\theta \in \text{Vect}(Z)$  is equal to 1.

A *fibred contact manifold* is a manifold  $Z$  with a contact form  $\alpha \in \Omega^1(Z)$  so that  $Z$  is the total space of a principal  $S^1$ -fiber bundle

$$p : Z \rightarrow Y, \quad p^{-1}(y) \cong S^1, \quad \forall y \in Y$$

so that  $\alpha$  is a connection one-form. A fibred stable Hamiltonian manifold is defined similarly. This ends the Definition.

**Lemma 2.2.** *Let  $Z$  be a fibered contact manifold. Any Legendrian submanifold  $\Lambda$  in a  $Z$  projects to a (possibly immersed) Lagrangian submanifold*

$$\Pi := p(\Lambda) \subset Y$$

*with respect to the symplectic form  $\omega_Y := -\text{curv}(\alpha)$  on  $Y$ .*

*Proof.* By the constant rank theorem,  $\Pi \subset Y$  is an immersed submanifold of dimension  $\dim(\Pi) = n = \dim(Y)/2$ . Since  $\alpha$  vanishes on  $\Lambda$ , so does  $d\alpha$ . Thus the symplectic form  $\omega_Y = -\text{curv}(\alpha)$  vanishes on  $\Pi$  as desired.  $\square$

*Example 2.3.* The following example arises from truncating the Harvey-Lawson Lagrangian by cutting off the complement of the unit ball. For

$$\epsilon = (\epsilon_1, \dots, \epsilon_n) \in \mathbb{R}^n$$

let

$$\Lambda_\epsilon := \{ |z_1|^2 + \epsilon_1 = \dots = |z_n|^2 + \epsilon_n \mid z_1 \dots z_n \in (0, \infty) \} \cap S^{2n-1}.$$

The subset  $\Lambda_\epsilon$  is a Legendrian submanifold for the stable Hamiltonian structure  $(\pi^*\omega_Y, \alpha_\epsilon)$  defined by a connection one-form  $\alpha_\epsilon$  on  $S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$  that is trivial on  $\Lambda_\epsilon$ . For example, one could take

$$\alpha_\epsilon := \frac{1}{1 + \sum_{i=1}^n \epsilon_i} \sum_{j=1}^n (r_j^2 + \epsilon_j) d\theta_j \in \Omega^1(S^{2n-1})$$

in an open neighborhood of  $\Lambda_\epsilon$ . Indeed, the contraction of the vector fields spanning the tangent space of  $T\Lambda_\epsilon$

$$\frac{\partial}{\partial \theta_i} - \frac{\partial}{\partial \theta_j} \in \text{Vect}(S^{2n-1}), \quad i, j \in \{1, \dots, n\}$$

with the one-form  $\alpha_\epsilon$  is

$$\iota \left( \frac{\partial}{\partial \theta_i} - \frac{\partial}{\partial \theta_j} \right) \alpha_\epsilon = \frac{(r_i^2 + \epsilon_i) - (r_j^2 + \epsilon_j)}{1 + \sum_{i=1}^n \epsilon_i} = 0$$

for all  $i, j$ . If  $\alpha_\epsilon$  is sufficiently close to  $\alpha$  in  $C^1$  then  $d\alpha_\epsilon$  is non-degenerate. So  $\alpha_\epsilon$  defines a contact structure.

A condition for a Lagrangian to admit a lift to an embedded Legendrian involves the subgroup generated by areas of surfaces bounding the Lagrangian.

**Definition 2.4.** For any compact oriented surface  $u : S \rightarrow Y$  bounding  $\Pi$  let

$$A(u) = \int_S u^* \omega$$

denotes its area. Denote by

$$\mathcal{A}(\Pi) = \langle \{A(u), u : S \rightarrow Y, u(\partial S) \subset \Pi\} \rangle \subset \mathbb{R}$$

the *area subgroup* generated by the areas  $A(u)$ .

**Proposition 2.5.** *Suppose that  $p : Z \rightarrow Y$  is a fibered contact manifold,  $Y$  is simply connected and  $\Pi \subset Y$  is a connected embedded Lagrangian submanifold, with respect to the symplectic form  $\omega_Y := -\text{curv}(\alpha)$  on  $Y$ . There exists an embedded Legendrian submanifold  $\Lambda$  projecting to  $\Pi$  if and only if the area subgroup  $\mathcal{A}(\Pi)$  is discrete. In this case, suppose that the areas in  $\mathcal{A}(\Pi)$  are integer multiples of  $1/k$  for some non-zero integer  $k \in \mathbb{Z}$ . Then the order of any fiber divides  $k$ .*

*Proof.* The proof is an application of the usual relationship between integral of the curvature and holonomy. Let  $y \in \Pi$  be a base point,  $Z_y \subset Z$  the fiber over  $y$  and  $z \in Z_y$ . The restriction of  $Z$  to  $\Pi$  is flat and so

$$\ker(\alpha) \subset TZ$$

is an integrable distribution on  $TZ|_{\Pi}$ . Let

$$\Lambda \subset Z, \quad T\Lambda = \ker(\alpha)|_{\Lambda}$$

be the leaf of the associated foliation through  $z$ . The holonomy of  $\Lambda$  around any loop  $\gamma$  bounding a disk  $u : D \rightarrow Y$  may be computed as follows. Let  $\psi$  denote a trivialization of the pull-back bundle.

$$\psi : D \times S^1 \rightarrow u^*Z.$$

The trivializing section is section  $\psi(\cdot, 1) : D \rightarrow u^*Z$ . Then

$$\begin{aligned} \text{Hol}_{\Lambda}(\gamma) &= \exp \left( 2\pi i \int_{\partial D} \psi(\cdot, 1)^* \tilde{u}^* \alpha \right) \\ &= \exp \left( 2\pi i \int_D d\psi(\cdot, 1)^* \tilde{u}^* \alpha \right) \\ &= \exp \left( 2\pi i \int_D \psi(\cdot, 1)^* \tilde{u}^* d\alpha \right) \\ &= \exp \left( 2\pi i \int_D u^* \omega \right) \end{aligned}$$

where  $\tilde{u} : u^*Z \rightarrow Z$  is the natural lift. Suppose the area of any disk  $u : D \rightarrow Y$  is divisible by  $k$ . Let  $\gamma : S^1 \rightarrow Y$  be a loop. Since  $Y$  is simply connected,  $\gamma$  bounds a disk smooth disk  $u : D \rightarrow Y$ , and the holonomy of  $\partial u$  is the exponential of  $2\pi i$  times the area of  $u$ . Since the area is divisible by  $k$ , the holonomy is an  $k$ -th root of unity by assumption. It follows that the fiber  $\Lambda_y$  contains  $\mathbb{Z}_k z$ , so  $k$  divides  $|\Lambda_y|$ .  $\square$

*Example 2.6.* The Clifford Legendrian of (1) is a Legendrian in the standard contact sphere  $S^{2n-1}$  that is an  $n$ -to-1 cover of the Clifford Lagrangian in projective space  $\mathbb{C}P^{n-1}$ . Consider the fibration  $S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$ . The symplectization of  $S^{2n-1}$  may be identified with punctured affine space

$$\mathbb{R}^n \times S^{2n-1} \cong \mathbb{C}^n - \{0\}.$$

The Maslov index two disks bounding the Clifford torus  $\Pi_{\text{Cliff}}$  are those induced by the maps to  $\mathbb{C}^n - \{0\}$

$$u(z) = \frac{1}{\sqrt{n}}(1, \dots, 1, z, 1, \dots, 1)$$



and have areas  $1/n$ , as in Cho-Oh [16]. The Clifford Legendrian  $\Lambda_{\text{Cliff}}$  is an  $n$ -to-1 cover of  $\Pi_{\text{Cliff}}$ . For example, in the case  $n = 2$  the disks correspond to the two hemispheres of  $\mathbb{C}P^1 = S^2$  and the Legendrian lift is

$$\Lambda_{\text{Cliff}} = \left\{ |z_1| = |z_2| = \frac{1}{\sqrt{2}}, z_1 z_2 \in (0, \infty) \right\}$$

is a double cover of the equator  $\Pi_{\text{Cliff}}$  with  $\mathbb{Z}_2$  action on the fiber given by scalar multiplication by  $-1$ .

The following are standard constructions on fibered contact/stable Hamiltonian manifolds, that is, circle bundles with connection.

**Lemma 2.7.** (a) (Unions) *Let  $Z_1 \rightarrow Y_1$  and  $Z_2 \rightarrow Y_2$  be circle-fibered contact resp. stable Hamiltonian manifolds. The disjoint union, written  $Z_1 \sqcup Z_2 \rightarrow Y_1 \sqcup Y_2$  is also a circle-fibered contact resp. stable Hamiltonian manifolds. If  $\Lambda_1 \subset Z_1$  and  $\Lambda_2 \subset Z_2$  then are immersed resp. embedded Legendrian submanifold then the disjoint union  $\Lambda_1 \sqcup \Lambda_2$  is an immersed resp. embedded Legendrian submanifold of  $Z_1 \sqcup Z_2$ .*

(b) (Exterior tensor products) *If  $Z_1 \rightarrow Y_1$  and  $Z_2 \rightarrow Y_2$  are circle-fibered contact/stable Hamiltonian manifolds then so is the exterior tensor product*

$$(Z_1 \boxtimes Z_2) = (Z_1 \times Z_2)/S^1 \rightarrow Y_1 \times Y_2$$

*where the  $S^1$ -action is the anti-diagonal action. If  $\Lambda_1 \subset Z_1$  and  $\Lambda_2 \subset Z_2$  are immersed Legendrian submanifolds then the image*

$$\Lambda_1 \boxtimes \Lambda_2 = (\Lambda_1 \times \Lambda_2)/S^1 \subset Z_1 \boxtimes Z_2$$

*of  $\Lambda_1 \times \Lambda_2$  in  $Z_1 \boxtimes Z_2$  is also an immersed Legendrian submanifold. If each projection  $\pi_j|_{\Lambda_j}, j = 1, 2$  is an embedding, so that in particular  $\Lambda_j$  is embedded, then  $\Lambda_1 \boxtimes \Lambda_2$  is also embedded.*

(c) (Finite covers) *Let  $\mathbb{Z}_m \subset S^1$  denote the finite subgroup of order  $m$ . If  $Z \rightarrow Y$  is a circle-fibered contact/stable Hamiltonian manifold then so is the quotient  $Z/\mathbb{Z}_m \rightarrow Y$ . Note that the transition maps of  $Z/\mathbb{Z}_m$  are the  $m$ -th powers of those of  $Z$  and so the Chern classes  $c_1(Z/\mathbb{Z}_m)$  and  $c_1(Z)$  in  $H^2(Y)$  are related by*

$$c_1(Z/\mathbb{Z}_m) = mc_1(Z).$$

*If  $\Lambda \subset Z$  is an immersed Legendrian submanifold then the image  $\underline{\Lambda}$  of  $\Lambda$  in  $Z/\mathbb{Z}_m$  is also an immersed Legendrian submanifold.*

(d) (Tensor products) *If  $Z_1 \rightarrow Y$  and  $Z_2 \rightarrow Y$  are circle-fibered contact/stable Hamiltonian manifolds*

$$Z_1 \otimes Z_2 := \Delta^*(Z_1 \boxtimes Z_2 \rightarrow Y \times Y)$$

*where  $\Delta : Y \rightarrow Y \times Y$  is the diagonal, is a circle-fibered contact manifold over  $Y$ . As a special case, if  $Z \rightarrow Y$  is a circle-fibered contact/stable Hamiltonian manifold then so is the  $k$ -fold tensor product*

$$Z^{\otimes k} := (Z \times_Y \dots \times_Y Z)/(S^1)^{k-1}.$$

- (e) (Symplectic quotients) *Let  $Z \rightarrow Y$  be a circle-fibered contact/stable Hamiltonian manifold with an action of a Lie group  $H$  preserving the contact form  $\alpha$  and commuting with the action of  $S^1$ . Let  $\mathfrak{h}$  be the Lie algebra of  $H$  and  $\mathfrak{h}^\vee$  its dual. The moment map for the action of  $H$  is the map*

$$\Phi : Z \rightarrow \mathfrak{h}^\vee, \quad z \mapsto (\xi \mapsto \alpha(\xi_Z(z))).$$

*The map  $\Phi$  is the pull-back of a moment map  $\bar{\Phi} : Y \rightarrow \mathfrak{h}^\vee$  for the action of  $H$  on  $Y$ . The symplectic quotient of  $Z$  is*

$$Z//H = \Phi^{-1}(0)/H.$$

*Suppose  $H$  acts properly and freely on  $\bar{\Phi}^{-1}(0)$ . Then  $Z//H$  is a circle-fibered contact manifold over*

$$Y//H := \bar{\Phi}^{-1}(0)/H.$$

*Let  $\Lambda \subset Z$  be an  $H$ -invariant immersed resp. embedded Legendrian submanifold. Then  $\Lambda$  is contained in  $\Phi^{-1}(0)$  and  $\Lambda/H$  is an immersed resp. embedded Legendrian submanifold of  $Z//H$ .*

*Proof.* Most of these claims are immediate and left to the reader. In the case of products, suppose that the connection forms are  $\alpha_1 \in \Omega^1(Z_1)$  and  $\alpha_2 \in \Omega^1(Z_2)$ . The one-form

$$\pi_1\alpha_1^* + \pi_2\alpha_2^* \in \Omega^1(Z_1 \times Z_2)$$

is basic (that is, invariant and vanishes on the vertical subspace) for the anti-diagonal  $S^1$  action. Thus it descends to a one-form on  $Z_1 \boxtimes Z_2$ . An easy check shows that this one-form is contact. Since  $\pi_1\alpha_1^* + \pi_2\alpha_2^*$  vanishes on  $\Lambda_1 \times \Lambda_2$ , its image  $\Lambda_1 \boxtimes \Lambda_2$  is Legendrian.  $\square$

*Example 2.8.* Continuing Example 2.6, let  $\Pi = \Pi_{\text{Cliff}}$  be the Clifford Lagrangian in  $\mathbb{C}P^{n-1}$ . Let  $Z \rightarrow \mathbb{C}P^{n-1}$  be the unit canonical bundle, that is, the set of unit vectors in the total space of the line bundle

$$\mathcal{O}(-n) = \Lambda_{\mathbb{C}}^{n-1} T\mathbb{C}P^{n-1}$$

with respect to the metric induced by the Fubini-Study metric on  $\mathbb{C}P^{n-1}$ . The horizontal lift  $\Lambda$  is diffeomorphic to  $\Pi$  under the projection; the horizontal lift  $\Lambda \rightarrow \Pi$  is defined by setting

$$s(y) \in \Lambda_{\mathbb{R}}^{n-1} T_y \Pi \cong \Lambda_{\mathbb{C}}^{n-1} T_y Y$$

the unique unit volume form defining the orientation on  $\Pi$ .

For a given contact/stable Hamiltonian form  $\alpha$  on  $Z$ , the *Reeb vector field*  $R_\alpha$  is the unique vector field on  $Z$  defined by

$$\omega(R_\alpha) = 0, \quad \alpha(R_\alpha) = 1.$$

A *Reeb orbit* is a closed orbit of the Reeb vector field

$$\gamma : [0, T] \rightarrow Z, \quad d\alpha\left(\frac{d}{dt}\gamma\right) = 0, \quad \alpha\left(\frac{d}{dt}\gamma\right) = 1, \quad \gamma(0) = \gamma(T).$$

A *Reeb chord* with boundary on a Legendrian  $\Lambda$  is a path starting and ending at the Legendrian:

$$\gamma : [0, T] \rightarrow Z, \quad d\alpha\left(\frac{d}{dt}\gamma\right) = 0, \quad \alpha\left(\frac{d}{dt}\gamma\right) = 1, \quad \gamma(\{0, T\}) \subset \Lambda.$$

The number  $T$  is the *action* of the Reeb orbit/chord; here the action is simply the change in angle in the fiber.

In our circle-fibered case, all Reeb orbits are multiple covers of circle fibers. Denote by  $\mathcal{R}(\Lambda)$  the set of Reeb chords:

$$\mathcal{R}(\Lambda) = \{ \gamma : [0, 1] \rightarrow Z \mid \gamma(\{0, 1\}) \subset \Lambda \}.$$

**Lemma 2.9.** *Suppose that as above  $\Lambda \subset Z$  is a compact Legendrian whose image  $\Pi \subset Y$  is an embedded Lagrangian submanifold. There is a diffeomorphism  $\mathcal{R}(\Lambda) \rightarrow (\Lambda \times_{\Pi} \Lambda) \times \mathbb{Z}_{\geq 0}$ .*

*Proof.* Given a Reeb chord  $\gamma$ , the starting and ending point  $\gamma(0), \gamma(1)$  of the chords together with the number of times  $\gamma(t)$  crosses the endpoint  $\gamma(0)$  as  $\gamma$  winds around the fiber  $Z_{p(\gamma)}$  uniquely specifies the chord.  $\square$

**Corollary 2.10.** *If  $\Pi$  is embedded then each component of  $\mathcal{R}(\Lambda)$  is diffeomorphic to  $\Lambda$ . If  $\Pi$  is immersed with transverse self-intersection then the components are either points or diffeomorphic to  $\Lambda$ . In either case, the fibers of the starting points resp. ending point map  $\mathcal{R}(\Lambda) \rightarrow \Lambda$  are identified with discrete subsets of  $\mathbb{R}_{>0}$  via the length map.*

**2.2. Lagrangian cobordisms.** In this section, we define symplectic and Lagrangian cobordisms and introduce assumptions that guarantee that the counts of pseudo-holomorphic disks give rise to chain maps between Legendrian dga's.

**Definition 2.11.** Let  $(Z_{\pm}, Y_{\pm}, \alpha_{\pm}, \omega_{\pm})$  be circle-fibered stable Hamiltonian manifolds. A *symplectic cobordism* from  $(Z_+, Y_+, \alpha_+, \omega_+)$  to  $(Z_-, Y_-, \alpha_-, \omega_-)$  is a symplectic manifold with boundary  $(\tilde{X}, \partial\tilde{X}, \omega)$  whose boundary is equipped with a partition

$$\partial\tilde{X} = \partial_+\tilde{X} \cup \partial_-\tilde{X}$$

and diffeomorphisms

$$\iota_{\pm} : Z_{\pm} \rightarrow \partial_{\pm}\tilde{X}$$

so that

- (a) the two forms satisfy the proportionality relation

$$\iota_{\pm}^* \omega \mid_{\partial_{\pm}\tilde{X}} = \lambda_{\pm} \omega_{\pm}$$

for some positive constants  $\lambda_{\pm} \in \mathbb{R}$ , and

- (b) the natural isomorphism between the normal bundle of  $Z_{\pm}$  in  $\tilde{X}$  and the kernel of  $Dp : TZ_{\pm} \rightarrow TY_{\pm}$  is orientation preserving resp. reversing for the positive resp. negative boundary.

Denote by  $X$  the interior of  $\tilde{X}$ , viewed as a manifold with cylindrical ends, and, abusing terminology, call  $X$  the cobordism.

*Remark 2.12.* In the definition of cobordism the one-forms  $\alpha_{\pm} \in \Omega^1(Z_{\pm})$  are not required to extend over  $X$ , and will not, in our basic example of the Harvey-Lawson filling. For the most part, the reader could restrict to the case that  $Z_{\pm}$  are contact manifolds, keeping in mind that the truncation of asymptotically-cylindrical manifolds such as the Harvey-Lawson filling are only Legendrian with respect to stable Hamiltonian structures on the boundary of the truncation.

*Remark 2.13.* Because of the assumption on the orientations, the notion of symplectic cobordism is not symmetric. That is, a cobordism from  $Z_-$  to  $Z_+$  is not also a cobordism from  $Z_+$  to  $Z_-$ .

*Example 2.14.* Truncations of symplectizations are particular examples of symplectic cobordisms. The symplectization of a contact manifold with one-form  $(Z, \alpha)$  is the symplectic manifold

$$\mathbb{R} \times Z, \quad \omega := d(e^{-s}\alpha)$$

where  $s$  is the coordinate on  $\mathbb{R}$ . Here we use the negative sign in the exponential  $e^{-s}$ , which is the opposite of the usual convention, since the incoming ends in our holomorphic maps will be at the large-volume end of the cobordism. For real numbers  $\sigma_- < \sigma_+$  The truncation

$$X = (\sigma_-, \sigma_+) \times Z$$

is a symplectic cobordism from  $(Z, \alpha)$  to itself, with proportionality constants for the two-forms given by  $e^{-\sigma_{\pm}}$ .

**Definition 2.15.** The *compactification* of a symplectic cobordism  $X$  is the symplectic manifold  $\bar{X}$  obtained by collapsing the null foliation on the boundary:

$$(3) \quad \bar{X} = \tilde{X} / \sim = (\tilde{X} - \partial\tilde{X}) \sqcup (\partial\tilde{X} / S^1)$$

where  $\sim$  is the trivial equivalence relation on  $\tilde{X} - \partial\tilde{X}$  and the equivalence relation

$$z \sim e^{2\pi i\theta} z, \quad z \in Z_{\pm}, \theta \in \mathbb{R}$$

generated by the circle action on the boundary.

*Example 2.16.* Continuing Example 2.14, the compactification of the symplectization  $X = \mathbb{R} \times Z$  of a contact manifold  $Z$  is the smooth manifold  $X$  obtained as the associated sphere bundle. That is,  $X$  is diffeomorphic to the quotient of  $Z \times S^2$  by the diagonal  $S^1$ -action.

**Lemma 2.17.** *Let  $X$  be a symplectic bordism from  $Z_-$  to  $Z_+$ . The compactification  $\bar{X}$  is a smooth symplectic manifold containing  $Y_-$  and  $Y_+$  as codimension-two symplectic submanifolds.*

*Proof.* The statement of the Lemma is a special case of the symplectic cut construction in Lerman [39]: Let  $U_{\pm} \subset X$  be an open neighborhood of  $Z_{\pm}$  in  $X$ , which may be identified with a subset of  $\mathbb{R} \times Z_{\pm}$  by the coisotropic embedding theorem. Consider the product  $\mathbb{R} \times Z_{\pm} \times \mathbb{C}$  with the diagonal  $S^1$ -action with moment map

$$\Phi : \mathbb{R} \times Z_{\pm} \times \mathbb{C} \rightarrow \mathbb{R}, \quad (s, z, w) \mapsto s \pm |w|^2/2.$$

The quotient of the zero level set of the moment map is naturally the quotient described in (3):

$$(U_{\pm} \times \mathbb{C}) // S^1 \cong (Z_{\pm} \times \{0\}) / S^1 \cup (U_{\pm} - Z_{\pm}) \cong \overline{U_{\pm}}.$$

The compactification  $\overline{X}$  is obtained from  $\mathbb{R} \times Z_{\pm}$  near the boundary by taking the symplectic quotient by a free action. The quotient is smooth by the Meyer-Marsden-Weinstein theorem.  $\square$

*Remark 2.18.* The unit normal bundle  $N_{\pm,1}$  of the codimension-two submanifold  $Y_{\pm} = Z_{\pm} / S^1$  in  $\overline{X}$  is naturally identified with  $Z_{\pm}$  up to a reversal of the action for the incoming boundary. It follows that  $N_{\pm,1}$  as a circle bundle with connection is positively resp. negatively curved for the incoming resp. outgoing boundary.

Conversely, compact symplectic manifolds equipped with disjoint symplectic hypersurfaces induce cobordisms, under the assumption of suitable positivity/negativity of the normal bundles:

**Lemma 2.19.** *Let  $(\overline{X}, \overline{\omega})$  be a closed symplectic manifold with two disjoint symplectic hypersurfaces  $Y_{\pm}$ . Suppose that the (possibly inverted) unit normal bundles*

$$Z_{\pm} := N_{\pm,1}^{\pm 1} \rightarrow Y_{\pm}$$

*are equipped with connection one-forms  $\alpha_{\pm}$  so that*

$$d\alpha_{\pm} = \pm \lambda_{\pm} p_{\pm}^* \omega_{Y_{\pm}} \in \Omega^2(Z_{\pm})$$

*for some positive constants  $\lambda_{\pm} > 0$ . Then the real blow up of  $\overline{X}$  at  $Y_{\pm}$  has the structure of a symplectic cobordism from  $(Z_{-}, \alpha_{-}, p_{-}^* \omega_{Y_{-}})$  to  $(Z_{+}, \alpha_{+}, p_{+}^* \omega_{Y_{+}})$ .*

*Proof.* The real blow-up  $\tilde{X}$  is obtained by replacing the  $Y_{\pm}$  by their unit normal bundles and admits a blow-down map to  $\overline{X}$ . The two-form  $\tilde{\omega}$  on  $\tilde{X}$  is obtained by pull-back of the form  $\overline{\omega}$  on  $\overline{X}$ .  $\square$

We view the interiors of the cobordisms as manifolds with cylindrical ends. Denote the interior of the cobordism  $X = \tilde{X} - \partial \tilde{X}$ . Tubular neighborhoods give rise to proper embeddings

$$\kappa_{\pm} : (\sigma_{\pm,0}, \sigma_{\pm,1}) \times Z_{\pm} \rightarrow X.$$

Equip  $(\sigma_{\pm,0}, \sigma_{\pm,1}) \times Z_{\pm}$  with the two form

$$\omega_{\pm} := \pi_{Y_{\pm}}^* \omega_{Y_{\pm}} + d(e^{-s} \alpha_{\pm}).$$

**Lemma 2.20.** *For intervals  $(\sigma_{\pm,0}, \sigma_{\pm,1})$  that are sufficiently small, the forms  $\omega_{\pm}$  are symplectic and the maps  $\kappa_{\pm}$  may be taken to be symplectomorphisms.*

*Proof.* The local model for constant rank embeddings in Marle [40] implies the existence of the claimed symplectomorphism, since the symplectic forms by definition agree at  $Z_{\pm}$ .  $\square$

**Definition 2.21.** For a symplectic cobordism  $X$  from  $Z_{+}$  to  $Z_{-}$  and two Legendrian submanifolds  $\Lambda_{\pm} \subset Z_{\pm}$  so that  $X$  is equipped with cylindrical ends, a *Lagrangian*

*cobordism cylindrical near infinity* from  $\Lambda_+$  to  $\Lambda_-$  is a Lagrangian submanifold  $L \subset X$  such that

$$L \cap ((\sigma_{\pm,0}, \sigma_{\pm,1}) \times Z_{\pm}) = (\sigma_{\pm,0}, \sigma_{\pm,1}) \times \Lambda_{\pm}$$

for some intervals  $(\sigma_{\pm,0}, \sigma_{\pm,1}) \subset \mathbb{R}$ . We call the pair  $(X, L)$  a *cobordism pair*.

Given a Lagrangian cobordism that is cylindrical near infinity, we form a Lagrangian in the compactified symplectic cobordism by closure. Let  $L \subset X$  be a Lagrangian cobordism, and  $\bar{L} \subset \bar{X}$  the closure. We say that an immersed submanifold  $\iota : K \rightarrow \bar{X}$  containing  $\bar{L}$  is cleanly self-intersecting if the fiber product  $K \times_{\iota, \iota} K \subset K \times K$  has the structure of a smooth submanifold so that at any points  $k_1, k_2 \in K$  with  $\iota(k_1) = \iota(k_2) = x$

$$D\pi(T_{k_1}K) \cap D\pi(T_{k_2}K) = D\pi(T_{k_1, k_2}K \times_{\iota, \iota} K).$$

**Proposition 2.22.** *Let  $(X, L)$  be a cobordism pair from  $(Z_-, \Lambda_-)$  to  $(Z_+, \Lambda_+)$  in the sense of Definition 2.21. There exists a manifold with boundary  $\bar{L}$  and an immersion  $\phi : \bar{L} \rightarrow \bar{X}$  with clean self-intersection so that  $\phi(\phi^{-1}(X)) = L$ .*

*Proof.* Suppose that  $\Lambda$  intersects some fiber  $Z_y \cong S^1$  over  $y \in Y$  in a subset  $\Lambda \cap Z_y = \{\theta_1, \dots, \theta_k\}$ . Then the subset

$$\mathbb{R}e^{2\pi i\theta_1} \cup \dots \cup \mathbb{R}e^{2\pi i\theta_k} \subset \mathbb{C} \cong Z_y \times_{\mathbb{C}^\times} \mathbb{C}$$

is a cleanly-intersecting subset of the fiber of the normal bundle of  $Y$  in  $\bar{X}$ . The union of such fibers over  $\Pi$  is locally a cleanly-intersecting Lagrangian in the normal bundle  $N_Y$ . Via the identification of an open neighborhood  $U$  of  $Y$  with an open neighborhood of the zero section in  $N_Y$ , denote the union of these fibers with  $L$  by  $\hat{L}$ . Then  $\hat{L}$  is a non-compact, cleanly-self-intersecting Lagrangian containing  $\bar{L} \cap U$  described locally as

$$L \cap U \cong \mathbb{R}_+ e^{2\pi i\theta_1} \cup \dots \cup \mathbb{R}_+ e^{2\pi i\theta_k}.$$

□

In particular, Proposition 2.22 implies that  $\bar{L}$  is a closed subset of a cleanly-self-intersecting Lagrangian  $\tilde{L} \rightarrow \bar{X}$ .


**Definition 2.23.** Let  $\bar{X}$  be a compact symplectic manifold. A compact subset  $\bar{L} \subset \bar{X}$  has a *branch point* at  $p \in \bar{L}$  if there exists a Darboux chart

$$\phi : U \rightarrow (\mathbb{C}^n, \omega_0)$$

where  $U$  is a neighborhood of  $p$  and  $\phi$  is a symplectomorphism sending  $p$  to 0, such that  $\phi(L)$  is the union of a collection of disjoint open subsets of a finite collection

$$W_1, \dots, W_k \subset \mathbb{C}^n$$

of Lagrangian subspaces of  $(\mathbb{C}^n, \omega_0)$  and an isotropic subspace  $W_\cap \subset \mathbb{C}^n$ . A subset  $\bar{L} \subset \bar{X}$  is a *branched Lagrangian* for any  $p \in \bar{L}$ , either  $\bar{L}$  is a smooth Lagrangian in an open neighborhood of  $p$  or  $p$  is a branch point of  $\bar{L}$  as defined above.


 FIGURE 2. Local model for the Lagrangian  $\bar{L}$  near  $\Pi$ 

We have in mind the case that the Lagrangian is locally homeomorphic to the product of a smooth (lower dimensional) Lagrangian with a union of half-lines ending at the origin in the complex plane, as in Figure 2. If  $\bar{L} \subset \bar{X}$  is a branched Lagrangian then the set of branch points  $L_\cap$  is an isotropic submanifold, being locally modelled on the subspace  $W_\cap$  near any branch point.

**Lemma 2.24.** *Let  $(Z, \alpha)$  be a fibered contact manifold. If  $\Lambda_\pm$  are isotopic Legendrians in  $(Z, \alpha)$  then there exists a Lagrangian cobordism from  $\Lambda_+$  to  $\Lambda_-$ .*

*Proof.* (c.f. Chantraine [11] and Golovko [35, Example 4.1].) Consider a Legendrian isotopy from  $\Lambda_-$  to  $\Lambda_+$ . By definition, such an isotopy consists of a family of Legendrians

$$\Lambda_\rho \subset Z, \rho \in \mathbb{R}, \quad \Lambda_\rho = \Lambda_-, \rho \ll 0, \quad \Lambda_\rho = \Lambda_+, \rho \gg 0.$$

Choose a family of diffeomorphisms

$$\psi_s : Z \rightarrow Z, \quad s \in \mathbb{R}$$

so that

$$\psi_s(\Lambda_-) = \Lambda_s, \forall s, \quad \partial_s \psi_s = 0, |s| \gg 0.$$

Consider the family of contact forms

$$\alpha_\rho = \psi_\rho^* \alpha \in \Omega^1(Z).$$

Define a one-form  $\tilde{\alpha} = (\alpha_s)_{s \in \mathbb{R}} \in \Omega^1(\mathbb{R} \times Z)$  and a two-form

$$\omega| = d_{\mathbb{R} \times Z}(e^{-s} \tilde{\alpha}) = -(ds) \wedge (e^{-s} \alpha_s + e^{-s} \partial_s \alpha_s) + e^{-s} d_Z \alpha_s \in \Omega^2(\mathbb{R} \times Z).$$

The top exterior power of the two-form is

$$(4) \quad \omega^n = -n(ds) \wedge (e^{-s} \alpha_s + e^{-s} \partial_s \alpha_s) \wedge (d_Z \alpha_s)^{n-1}.$$

By taking  $\alpha_s$  to be slowly varying, we may assume that the following inequality holds point-wise with respect to some Riemannian metric on the bundle of forms:

$$(5) \quad \|ds \wedge \alpha_s \wedge d_Z(\alpha_s)^{n-1}\| > \|ds \wedge \partial_s \alpha_s \wedge d_Z(\alpha_s)^{n-1}\|.$$

Since  $ds \wedge \alpha_s \wedge d_Z(\alpha_s)^{n-1}$  is non-vanishing, equations (4) and (5) imply that  $\omega^n$  is non-vanishing. Thus,  $\omega$  is a symplectic form on  $\mathbb{R} \times Z$ . Denote the inclusion

$$\iota : \mathbb{R} \times \Lambda \rightarrow \mathbb{R} \times Z.$$

The pull-back two-form is

$$\iota^*\omega = d(e^{-s}\iota^*\tilde{\alpha}) = 0.$$

Thus  $(\mathbb{R} \times Z, \omega)$  is a symplectic manifold containing  $\mathbb{R} \times \Lambda_-$  as a Lagrangian submanifold. The diffeomorphisms  $\psi_\rho$  identify the ends with the symplectizations  $\pm(|\rho|, \infty) \times Z$  for  $|\rho|$  sufficiently large. By symplectic cutting on  $\mathbb{R} \times Z$ , we obtain a compact symplectic manifold  $\bar{X}$  containing  $Y_\pm$  as symplectic submanifolds, up to re-scaling of the symplectic forms  $\omega_\pm$ .  $\square$

Our motivating examples of Lagrangian cobordisms are asymptotically-cylindrical rather than cylindrical near infinity. To explain the asymptotically-cylindrical definition, recall that the space of submanifolds has a natural  $C^0$ -topology. In this topology, open neighborhoods of a given submanifold are those submanifolds corresponding to sections of the normal bundle contained in some tubular neighborhood. Choose tubular neighborhoods of  $\Lambda_\pm$  in  $Z$  given by diffeomorphisms of the normal bundle  $N\Lambda_\pm$  onto an open neighborhood of  $\Lambda_\pm$ . The tubular neighborhoods of  $\Lambda_\pm$  induce tubular neighborhoods of  $\mathbb{R} \times \Lambda_\pm$  given by maps  $\mathbb{R} \times N\Lambda_\pm \rightarrow \mathbb{R} \times Z_\pm$ . Let

$$\tau_\sigma : \mathbb{R} \times Z \rightarrow \mathbb{R} \times Z, \quad (s, z) \mapsto (s + \sigma, z)$$

be the translation map.

**Definition 2.25.** Let  $(Z, \alpha)$  be a fibered contact manifold. Let  $L_\nu \subset \mathbb{R} \times Z, \nu \in \mathbb{N}$  be a sequence of submanifolds given as the graphs of sections

$$s_{L_\nu} : \mathbb{R} \times \Lambda \rightarrow \mathbb{R} \times N\Lambda.$$

The sequence  $L_\nu$   $C^\ell$ -converges to  $(\sigma_-, \sigma_+) \times \Lambda$  if the sequence of sections  $s_{L_\nu}$  converges to 0 in the  $C^\ell$ -norm uniformly on  $(\sigma_-, \sigma_+) \times \Lambda$ .

A non-compact Lagrangian  $L \subset X$  is *asymptotically cylindrical* if the submanifolds

$$\tau_{-\sigma}(L \cap ((\sigma, \sigma + 1) \times Z)) \subset (0, 1) \times Z$$

converge in  $C^1$  on  $(0, 1) \times Z$  in the sense above. This ends the Definition.

*Example 2.26.* For the Clifford Legendrian  $\Lambda_{\text{Cliff}}$  in (1), a natural asymptotically-cylindrical filling was introduced by Harvey-Lawson [37], see also Joyce [38]. Let

$$a_1, \dots, a_n \geq 0$$

be non-negative constants exactly two of which are zero (without loss of generality, the last two  $a_{n-1} = a_n = 0$ .) The *Harvey-Lawson asymptotic filling* denoted  $L_{(1)}$  of the Clifford Legendrian is given by (2), and is asymptotic to  $\mathbb{R}_{>0} \times \Lambda_{\text{Cliff}}$  at infinity in  $\mathbb{C}^n$ .

By truncation we obtain fillings of Legendrians in the spheres of fixed norm. Namely, each intersection with a ball  $L_1 \cap B_{e^{-\sigma}(0)}$  fills the Legendrian

$$\Lambda_\epsilon = \left\{ \begin{array}{l} |z_1|^2 - a_1^2 = |z_2|^2 - a_2^2 = \dots = |z_n|^2 - a_n^2 \\ z_1 z_2 \dots z_n \in (0, \infty) \end{array} \right\} \cap e^{-\sigma} S^{2n-1}.$$

The projection is the Lagrangian torus orbit torus

$$(6) \quad \{|z_1|^2 - a_1^2 = |z_2|^2 - a_2^2 = \dots = |z_n|^2 - a_n^2\} \cap e^\sigma S^{2n-1} / S^1.$$



This Lagrangian is a toric moment fiber over a point close, but not equal to, the barycenter  $e^{-\sigma}(1, 1, \dots, 1)/n$  of the moment polytope  $\Delta_{n-1}$  of  $\mathbb{C}P^{n-1}$

$$\Delta_{n-1} = \{\lambda_1 + \dots + \lambda_n = 1\} \cap \mathbb{R}_{\geq 0}^n.$$

In particular, if  $L$  is an asymptotically cylindrical Lagrangian with limits  $\Lambda_{\pm}$  that have monotone projections to  $Y_{\pm}$ , the truncations  $\Lambda_{\sigma_{\pm}}$  may not have monotone projections.

For technical reasons, in the construction of moduli spaces of pseudoholomorphic curves it is convenient to have cylindrical-near-infinity Lagrangians rather than asymptotically cylindrical Lagrangians. Presumably, the gluing and compactness results in symplectic field theory hold equally well for asymptotically cylindrical Lagrangians, but these results have yet to appear in the literature. The following lemma allows us to replace asymptotically cylindrical Lagrangians with those cylindrical near infinity.

*Remark 2.27.* (Outline of straightening construction) Since the Lemma is somewhat technical, we give an outline of the construction first. We are given a Lagrangian asymptotically-cylindrical cobordism that we wish to make cylindrical-near-infinity. The intersection  $\Lambda_{\sigma}$  of the Lagrangian with any slice  $\{\sigma\} \times Z_{\pm}$  on the neck is not necessarily Legendrian. Instead, we construct a nearby contact structure  $\alpha_{\sigma}$  for which the slice is Legendrian. The cylinder  $L_{\sigma} = \mathbb{R} \times \Lambda_{\sigma}$  is Lagrangian for the symplectization of  $Z$  with respect to  $\alpha_{\sigma}$ . Now by the equivariant coisotropic embedding theorem  $L$  may be assumed to agree with  $L_{\sigma}$  on an open subset, and replacing the end of  $L$  with half of  $L_{\sigma}$  gives the desired cobordism that is cylindrical-near-infinity. The price of this construction is that the cylindrical end of the Lagrangian cobordism is now modelled on a different contact structure. However, the new contact structure and Legendrian may be taken arbitrarily close to the original, and then invariance of holomorphic curves under deformation will show that the resulting Lagrangian cobordism still defines a chain map between Chekanov-Eliashberg algebras, as explained in [7, Remark 3.15].

**Lemma 2.28.** (Straightening Lemma) *Suppose that  $X$  is a smooth symplectic manifold with ends modelled on  $\mathbb{R} \times Z_{\pm}$  with symplectic form  $d(e^{-s}\alpha)$ . Let  $L \subset X$  be an asymptotically cylindrical Lagrangian with limiting Legendrians  $\Lambda_{\pm} \subset Z_{\pm}$ . Suppose that  $\Lambda_{\pm}$  fiber over Lagrangians  $\Pi_{\pm} \subset Y$  with finite covering group*

$$\Upsilon \cong \mathbb{Z}_k \subset S^1$$

*and  $L$  is  $\Upsilon$ -invariant on the ends. For  $\sigma > 0$  let*

$$X_{[\sigma_-, \sigma_+]} := X - \kappa_+^{-1}((\sigma_+, \infty) \times Z_+) - \kappa_-^{-1}((-\infty, \sigma_-) \times Z_-)$$

*denote the result of truncating the cylindrical ends at  $s = \sigma_{\pm}$ . For  $\sigma \in \mathbb{R}$  let*

$$\Lambda_{\sigma} := (\{\sigma\} \times Z_{\pm}) \cap L$$

*be the intersection with the  $\sigma$ -level set. For*

$$\sigma_{\pm} \in (0, \infty)$$

there exist connection one-forms  $\alpha_{\pm}$  on  $Z_{\pm}$  for which  $\Lambda_{\pm}$  are Legendrian and so that

$$L_{[\sigma_-, \sigma_+]} := X_{[\sigma_-, \sigma_+]} \cap L$$

can be isotoped through Lagrangians to a cylindrical-near-infinity Lagrangian cobordism pair  $(X, L')$  from  $(Z_-, \Lambda_{\sigma_-})$  to  $(Z_+, \Lambda_{\sigma_+})$  for some stable Hamiltonian manifolds  $(Z_{\pm}, \alpha_{\pm}, \omega_{\pm})$ .

*Proof.* Moser isotopy can be used to create a cylindrical end for which the Lagrangian is cylindrical near infinity. It suffices to consider the case that  $X$  has only a positive end  $Z_+$ , as the case of a negative end is similar. By assumption, the submanifold  $L$  is  $C^1$  close to  $\mathbb{R} \times \Lambda_+$  on the end. It follows that  $\Lambda_{\sigma_+}$  is smooth and is a codimension one submanifold of  $L$ . The projection

$$\Pi^{\sigma_+} = p(\Lambda_{\sigma_+}) \subset \{\sigma_+\} \times Z$$

is an (in general immersed) isotropic submanifold, since  $\omega = d(e^{-s}\alpha_{\pm})$  vanishes on  $\Lambda_{\sigma_+}$ . Because  $\Lambda_+$  is an  $\Upsilon$ -cover and  $\Lambda^{\sigma_+}$  is  $C^1$ -close to  $\Lambda_+$ , the map  $\Lambda^{\sigma_+} \rightarrow \Pi^{\sigma_+}$  is an  $\Upsilon$ -cover.

The first step is to choose a connection one-form for which the slice is horizontal. At any point  $z \in \Lambda_{\sigma_+}$  there is a unique one-form

$$\alpha_{\sigma_+} \in \text{Hom}(T_z Z_+, \mathbb{R})$$

for which  $T_z \Lambda_{\sigma_+}$  is horizontal and for which  $\alpha_{\sigma_+}(\frac{\partial}{\partial \theta}) = 1$ . By  $S^1$ -invariance,  $\alpha_{\sigma_+}$  extends to a form on the orbit of  $z$ . Since  $\Lambda_{\sigma_+} \subset L$  is by assumption  $\Upsilon$ -invariant,  $T_{hz} \Lambda_{\sigma_+}$  is horizontal for any  $h \in \Upsilon$ . This construction defines a one-form

$$\alpha_{\sigma_+} \in \Omega^1(Z|_{\Pi_{\sigma_+}})$$

of  $Z$  to  $\Pi_{\sigma_+}$ . Using an  $S^1$ -equivariant tubular neighborhood, one may extend  $\alpha_{\sigma_+}$  first to an open neighborhood of  $Z|_{\Pi_{\sigma_+}}$ , then to all of  $Z$  using a patching argument and convexity of the space of connection one-forms. For  $\sigma_+$  is sufficiently large,  $\alpha_{\sigma_+}$  is  $C^1$ -close to  $\alpha$ , and so the form  $d\alpha_{\sigma_+}$  is  $C^0$  close to  $d\alpha = \pi^*\omega$ .

We construct a Lagrangian cobordism that is cylindrical-near-infinity by a patching argument. The cylindrical submanifold  $L_{\sigma_+} = \mathbb{R} \times \Lambda_{\sigma_+}$  (shown in Figure 3) is isotropic with respect to the closed two-form

$$\omega_{\sigma_+} := \pi_{\sigma_+}^*(\omega|_{\{\sigma_+\} \times Z}) + d((s - \sigma_+)\alpha_{\sigma_+}) \in \Omega^2(\mathbb{R} \times Z_+)$$

where  $\pi_{\sigma} : \mathbb{R} \times Z_{\pm} \rightarrow X$  is the map obtained by composing projection onto  $Z_{\pm}$  with inclusion of  $Z_{\pm}$  into  $X$ . Furthermore,  $\omega_{\sigma_+}$  is symplectic and  $S^1$ -invariant in a tubular neighborhood of  $\{\sigma_+\} \times Z_+$ . By construction, the projection on the first factor is a moment map for the  $S^1$ -action. The equivariant coisotropic embedding theorem [36, p. 315] implies that the forms  $\omega$  and  $\omega_{\sigma_+}$  are symplectomorphic on the truncation

$$X_{(\sigma_0, \sigma_1)} := (\sigma_0, \sigma_1) \times Z \subset (0, \infty) \times Z$$

for  $\sigma_0 < \sigma_+ < \sigma_1$  where  $\sigma_i$  is close enough to  $\sigma_+$ , by a symplectomorphism

$$\psi : (X_{(\sigma_0, \sigma_1)}, \omega) \rightarrow (X_{(\sigma_0, \sigma_1)}, \omega_{\sigma})$$




FIGURE 3. Straightening an asymptotically cylindrical Lagrangian

that preserves the moment map for the  $S^1$ -action, which is projection on the first factor. Furthermore,  $\psi$  may be taken to equal the identity on  $\Lambda_{\sigma_+}$ , since the two-forms agree at the tangent space there. The image  $\psi(L)$  of  $L$  under  $\psi$  is a Lagrangian with respect to  $\omega_{\sigma_+}$  and agrees with  $L_{\sigma_+}$  at  $\Lambda_{\sigma_+}$ . It follows that  $\psi(L)$  is an exact deformation of  $L_{\sigma_+}$  in an open neighborhood of  $\Lambda_{\sigma_+}$ , given by the Hamiltonian flow of  $L_{\sigma_+}$  of some function

$$H : X_{(\sigma_0, \sigma_1)} \rightarrow \mathbb{R}.$$

A cylindrical-near-infinity Lagrangian  $\psi'(L) \subset X$  is given as Hamiltonian flow  $\psi'$  of  $L$  for  $\rho H$  where  $\rho$  is a function whose derivative is supported near  $\{\sigma_+\} \times \Lambda_{\sigma_+}$ , equal to 0 in an open neighborhood of  $\{\sigma_1\} \times \Lambda_{\sigma_+}$  and 1 in an open neighborhood of  $\{\sigma_0\} \times \Lambda_{\sigma_-}$ . Gluing together  $\psi'(L)$  with  $L$  (the map  $\psi$  is the identity outside of a small open neighborhood of  $\{\sigma_+\} \times Z_+$ ) gives the required Lagrangian  $L'$ .  $\square$

*Remark 2.29.* We may choose  $\alpha_{\sigma_{\pm}}$  arbitrarily close to  $\alpha$ , so that in particular  $\alpha_{\pm}$  define contact structures on  $Z_{\pm}$ , and so the submanifold  $\Lambda_{\sigma_+}$  is Legendrian.

*Remark 2.30.* We remark on the possibility of obtaining a cobordism between contact manifolds, rather than stable Hamiltonian manifolds. If  $\sigma_+, -\sigma_-$  are sufficiently large then the forms  $\alpha_{\pm}$  are contact forms. However, we may not be able to choose a cobordism from  $(Z_-, \alpha_-)$  to  $(Z_+, \alpha_+)$  in the sense that the forms  $d\alpha_{\pm}$  extend as symplectic forms over the cobordism and  $L$  is Lagrangian with respect to the extension. Indeed, the action of loops in the Legendrian  $\Lambda$  as  $d\alpha_{\pm}$  may not equal those of  $\omega_{\sigma_{\pm}}$ .

*Example 2.31.* We specialize to the case of the Harvey-Lawson filling. The straightening Lemma 2.28 implies that after a Hamiltonian isotopy we may assume that  $L_{(1)} \cong S^1 \times \mathbb{R}^{n-1} \subset \mathbb{C}^n$  is invariant under dilation in a neighborhood of some sphere  $e^{\sigma} S^{2n-1} \cap L_{(1)}$ .

**2.3. Reductions of symplectic fillings.** We introduce various constructions that will be used to produce fillings later. In particular, the following Lemma produces symplectic fillings by symplectic reduction:

**Lemma 2.32.** *Let  $X$  be a symplectic cobordism between  $Z_-$  and  $Z_+$  equipped with a Hamiltonian action of a Lie group  $H$  with moment map  $\Phi$  such that  $H$  acts properly and freely on  $\Phi^{-1}(0)$ . Then  $X//H := \Phi^{-1}(0)/H$  is a symplectic cobordism from  $Y_-//H$  to  $Y_+//H$  (both of which may possibly be empty). If  $L \subset X$  is an invariant Lagrangian cobordism from  $\Lambda_-$  to  $\Lambda_+$  then  $L//H$  is a Lagrangian cobordism from  $\Lambda_-//H$  to  $\Lambda_+//H$ .*

The proof is an immediate consequence of the Meyer-Marsden-Weinstein theorem, applied to Hamiltonian actions on manifolds with boundary for which the proof is the same. We obtain fillings of toric moment fibers in toric varieties by symplectic reduction as follows. Let  $Y$  be a projective toric variety  $Y$  equipped with a symplectic form corresponding to the anticanonical class; later we will assume that  $Y$  is Fano. Let  $H = (S^1)^n$  denote the  $n$ -torus acting in Hamiltonian fashion with moment map

$$\Psi : Y \rightarrow \mathbb{R}^n.$$

The moment polytope  $P = \Psi(Y)$  may be written as the set of points in  $\mathbb{R}^n$  satisfying a finite list of linear inequalities

$$P = \{\lambda \in \mathfrak{h}^\vee \mid \langle \lambda, \nu_i \rangle \geq -1, i = 1, \dots, k\} \subset \mathbb{R}^n.$$

We may assume by translation that  $P$  contains the 0 vector.

A singular symplectic filling is given by the cone construction. By the *cone*  $\text{Cone}(Y)$  on  $Y$  we mean the (usually singular) affine toric variety whose polytope is

$$\text{Cone}(P) = \mathbb{R}_{\geq 0}(P \times \{1\}) \subset \mathfrak{h}^\vee \times \mathbb{R}.$$

The symplectic form on  $Y$  induces a symplectic form on  $\text{Cone}(Y) - \{0\}$  so that the natural action

$$\hat{H} \times \text{Cone}(Y) \rightarrow \text{Cone}(Y)$$

of the torus  $\hat{H} = H \times S^1$  of one dimension higher is completely integrable. The moment map for this action

$$\hat{\Psi} : \text{Cone}(Y) \rightarrow \hat{\mathfrak{h}}^\vee$$

has moment polytope given by  $\text{Cone}(P)$ . The truncation of  $\text{Cone}(P)$ , the convex hull of the origin and  $P \times \{1\}$ , is the projective closure of  $\text{Cone}(Y)$ ; see for example Evans [32, Appendix F] for the correspondence between polytopes and projective toric varieties.

Consider in particular the case of projective space. Let  $Y = \mathbb{P}^{n-1}$ , so that the cone is  $\text{Cone}(\mathbb{P}^{n-1}) = \mathbb{C}^n$ . The unit tautological bundle  $Z = \mathcal{O}(-1)_1$  is then a contact manifold that is circle fibered over  $Y$ , and the complement

$$\text{Cone}^*(Y) \subset \text{Cone}(Y)$$

of the vertex  $0 \in \text{Cone}(Y)$  is a smooth symplectic cobordism from the unit tautological bundle  $Z$  to itself. The compactification  $\overline{X}$  of  $X$  may be taken to be the part of the cone between  $\sigma_- P \times \{\sigma_-\}$  and  $\sigma_+ P \times \{\sigma_+\}$ . For example for  $Y = \mathbb{C}P^{n-1}$  the compactification  $\overline{X}$  would have polytope defined by

$$a_1 + \dots + a_n \in [\sigma_-, \sigma_+], a_1, \dots, a_n \in \mathbb{R}_{\geq 0}.$$

We produce fillings corresponding to the choice of codimension two faces; this generalizes the Harvey-Lawson construction which depends on a choice of two coordinates, see Figure 4. Let  $P$  be the moment polytope for the anti-canonical polarization on  $Y$ ; by the Fano assumption,  $P$  contains 0. Let  $Q \subset P$  be a codimension two face, given as the intersection of facets with normal vectors  $\nu_1, \nu_2 \in \mathfrak{h}$ . Let  $\text{hull}(Q, 0) \subset P$  denote the convex hull of  $Q$  with the zero vector. Recall that the moment polytope for the cone on  $Y$  is the cone  $\text{Cone}(P)$  on  $P \times \{1\}$  in  $\hat{\mathfrak{h}} = \mathfrak{h} \times \mathbb{R}$ .

**Lemma 2.33.** *Let  $Y, P, Q$  be as above. For any vector  $\lambda$  in the interior of  $\text{hull}(Q, 0)$  there exists a Lagrangian  $L$  invariant under a codimension one sub-group of  $H$  whose image  $\Psi(L)$  under  $\Psi$  is the intersection of  $P$  with the line  $\{\lambda\} \times \mathbb{R}$ , and whose intersection  $L \cap \hat{\Psi}^{-1}(\lambda)$  projects to the Lagrangian torus orbit  $\Psi^{-1}(\lambda)$  in  $Y$ .*

*Proof.* We construct the Lagrangian by symplectic reduction. Choose a subspace  $\mathfrak{h}_1$  of codimension one in  $\mathfrak{h}$  intersecting the span of  $\nu_1, \nu_2$  in the span of  $\nu_1 + \nu_2$ :

$$\mathfrak{h}_1 \cap \text{span}(\nu_1, \nu_2) = \text{span}(\nu_1 + \nu_2).$$

The subspace  $\mathfrak{h}_1$  generates a torus subgroup  $H_1$ . Let  $\lambda_1 \in \mathfrak{h}_1$  be the projection of some point in  $\text{Cone}(Q) \subset \mathfrak{h}^\vee$ . Choose a splitting  $\mathfrak{h} \cong \mathfrak{h}_1 \oplus \mathbb{R}$  using, for example, the standard metric on  $\mathfrak{h} \cong \mathbb{R}^n$ . Let

$$\Psi_1 : \text{Cone}^*(Y) \rightarrow \mathfrak{h}_1^\vee$$

denote the moment map for the action of  $H_1$  obtained by composing the moment map  $\Psi$  for  $H$  with the projection  $\mathfrak{h}^\vee \rightarrow \mathfrak{h}_1^\vee$ . Let

$$Y_1 = \Psi_1^{-1}(\lambda_1) = \text{Cone}^*(Y) // H_1$$

denote the symplectic reduction of  $\text{Cone}(Y)^*$ . For generic values  $\lambda_1$ , the symplectic quotient  $Y_1$  is a toric orbifold of dimension two, and has a residual action of  $H/H_1$  with moment polytope  $P_1$  some slice of  $\text{Cone}^*(P)$ . The reduction of  $\text{Cone}^*(Y)$  at the locus  $\Psi^{-1}(\text{Cone}(Q))$  is smooth. As in Delzant [21], smoothness of the symplectic quotient follows from the fact that the vectors

$$(\nu_1 + \nu_2, 0), (\nu_1, 1), (\nu_2, 1) \in \mathfrak{h}_{\mathbb{Z}} \oplus \mathbb{Z}.$$

form a lattice basis for their real span. In particular,  $(\nu_1 + \nu_2, 0)$  generates a circle subgroup that acts freely on inverse image of the cone on  $Q$ . Since reductions by free actions are smooth, the claim follows. The polytope  $P_1$  has a vertex at some point  $v \in \text{Cone}^*(P)$  with normal vectors  $(1, 0)$ ,  $(0, 1)$ , and the line segment  $(\lambda, \lambda)$ ,  $\lambda \in [0, 1]$  is entirely contained in  $P_1$ . It follows that  $Y_1$  may be viewed as a symplectic cut of  $\mathbb{C}P^2$  by circle actions.

There is a natural bijection between invariant Lagrangians and Lagrangians in the symplectic quotient. An invariant Lagrangian is obtained by pulling back any Lagrangian in the four-dimensional quotient. The inverse image in  $\text{Cone}^*(Y)$  is a Lagrangian in  $\text{Cone}^*(Y)$  with the stated property.  $\square$

*Example 2.34.* (Harvey-Lawson via reduction) We explain how to obtain the Harvey-Lawson filling via symplectic reduction to a projective plane in projective three-space. Suppose that  $Y = \mathbb{C}P^2$  is the projective plane. The maximal torus  $\mathfrak{h}$  of

$SU(2)$  acts on  $Y$  with with moment polytope  $Q$  the convex hull of the projections of three standard basis vectors

$$P = \pi(\text{hull}\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}).$$

under the projection  $\pi$  from  $\mathbb{R}^3$  to  $\mathfrak{h} = \{\lambda_1 + \lambda_2 + \lambda_3 = 0\}$ . We identify  $\mathbb{R}^3$  with its dual using the standard inner product, and identify  $\hat{h}$  with  $\mathbb{R}^3$  using the decomposition  $\mathbb{R}^3 = \hat{h} \oplus \mathbb{R}$ , with the last factor being embedded as the diagonal. Let

$$Q = \{(0, 0, 1)\}$$

be the last vertex. The normal vectors to the edge with direction  $(1, 0, -1)$  resp.  $(0, 1, -1)$  are

$$\nu_1 = (1, -2, 1), \quad \nu_2 = (-2, 1, 1).$$

The sub-torus used for reduction has Lie algebra

$$\mathfrak{h}_1 = \text{span}(\nu_1 + \nu_2), \quad \nu_1 + \nu_2 = (-1, -1, 2).$$

The symplectic reduction  $\mathbb{C}P^3 // H_1$  may be identified with  $\mathbb{C}P^2$  via projection on the first two variables. The Harvey-Lawson Lagrangian is

$$\{|z_1|^2 + \epsilon = |z_2|^2 + \epsilon = |z_3|^2, z_1 z_2 z_3 \in [0, \infty)\}$$

and is the lift of the antidiagonal  $\overline{\{[z, \bar{z}, 1]\}}$  in the reduction  $\mathbb{C}P^2 \cong \mathbb{C}P^3 // H_1$ . See Figure 4.




FIGURE 4. Obtaining the Harvey-Lawson filling via reduction

### 3. PSEUDOHOLOMORPHIC BUILDINGS

In this section, we construct the moduli spaces of holomorphic buildings used for both the differential in the Chekanov-Eliashberg algebra and the chain maps used associated to Lagrangian cobordisms. Because our stable Hamiltonian manifolds are circle-fibered over symplectic manifolds and we consider only holomorphic disks rather than curves of higher genus, we may use Cieliebak-Mohnke [17] perturbations to regularize.

**3.1. Punctured surfaces.** We recall basic terminology for surfaces with strip-like ends at punctures.

**Definition 3.1.** A surface with strip-like and cylindrical ends  $S$  is obtained from a closed oriented surface-with-boundary  $\bar{S}$  by removing a finite collection of boundary points  $z_{e,\circ}, e = 1, \dots, e(\circ)$  and interior points  $z_{e,\bullet}, e = 1, \dots, e(\bullet)$

$$S = \bar{S} - \{z_{e,\circ}, e = 1, \dots, e(\circ), \quad z_{e,\bullet}, e = 1, \dots, e(\bullet)\}.$$

We call  $S$  a *punctured surface* for short, and the removed points *punctures*.

Suppose  $S$  is equipped with a conformal structure, giving rise to an almost complex structure

$$j : TS \rightarrow TS, \quad j^2 = -\text{Id}_{TS}.$$

An open neighborhood  $U_{e,\circ} \subset S$  of each puncture  $z_{e,\circ}$  is assumed to be equipped with a local coordinate, and similarly for open neighborhoods  $U_{e,\bullet}$  of  $z_{e,\bullet}$ . Each such coordinate gives holomorphic embeddings called *strip-like* resp. *cylindrical ends*

$$\kappa_{e,\circ} : U_{e,\circ} - \{z_{e,\circ}\} \cong \pm\mathbb{R}_{>0} \times [0, 1], \quad \kappa_{e,\bullet} : U_{e,\bullet} - \{z_{e,\bullet}\} \cong \pm\mathbb{R}_{>0} \times S^1.$$

The set of ends is denoted  $\mathcal{E}(S)$  and is equipped with a partition into incoming resp. outgoing ends

$$\mathcal{E}(S) = \mathcal{E}_+(S) \cup \mathcal{E}_-(S).$$

To keep with the terminology standard in the field, the punctures corresponding to incoming ends are called *positive* while the outgoing punctures are called *negative*.

In the Gromov compactification of the moduli space of holomorphic maps, nodal surfaces will appear as degenerations.

**Definition 3.2.** A *nodal surface*  $S$  is obtained from a surface with boundary  $\tilde{S}$  with strip-like and cylindrical ends by gluing along disjoint pairs of points  $w_{\pm}(e) \in \tilde{S}$ , where  $e$  ranges over some index set, which may be either a pair of boundary points  $w_{\pm}(e) \in \partial\tilde{S}$  or a pair of interior points  $w_{\pm}(e) \in \text{int}(\tilde{S})$ .

A nodal surface  $S$  is a *nodal disk* if each component  $S_v$  is a disk, the boundary  $\partial S$  is connected, and the combinatorial type of the curve (defined below) is a tree. .

A *marking* is a pair  $\underline{z} = (z_{\bullet}, z_{\circ})$  consisting of a tuple of points  $z_{\circ}$  on the boundary  $\partial S$  disjoint from the nodes and a tuple of points  $z_{\bullet} \in \text{int}(S)$  in the interior, with the property that the ordering of points on the boundary  $\partial S_v \cap z_{\circ}$  is cyclic.

A marked nodal surface  $(S, \underline{z})$  is *stable* if the group of automorphisms  $\text{Aut}(S)$  is finite. Equivalently, for each sphere component there are at least three special (nodal or marked points), while for each disk component there are at least three special boundary points or at least one special boundary point and one special interior point.

A marked nodal surface  $(S, \underline{z})$  has a combinatorial type  $\Gamma = \Gamma(S)$ , which is a graph whose vertices  $v \in \text{Vert}(\Gamma)$  correspond to components  $S_v$  and edges  $e \in \text{Edge}(\Gamma)$  correspond to nodes  $w_{\pm}(e) \in S$  or markings  $z(e) \in S$ ; we define the subset  $\text{Edge}_{\rightarrow}(\Gamma)$  to be the set of edges corresponding to markings and call them *leaves*.

For nodal disks  $(S, \underline{z})$ , this combinatorial type  $\Gamma$  is required to be a tree.

**3.2. Pseudoholomorphic buildings.** We define pseudoholomorphic buildings in symplectic cobordisms with ends modelled on cylinders over stable Hamiltonian manifolds.

**Definition 3.3.** Let  $Z \rightarrow Y$  be a circle-fibered stable Hamiltonian manifold as above and  $X = \mathbb{R} \times Z$  be the trivial cobordism. An almost complex structure  $J : TX \rightarrow TX$  is *cylindrical* if there exists an almost complex structure  $\bar{J} : TY \rightarrow TY$  so that the projection  $p_X : X \rightarrow Y$  is almost complex and  $J$  is the standard almost complex structure on any fiber. More precisely, let

$$\partial_s \in \text{Vect}(\mathbb{R} \times Z), \quad \partial_\theta \in \text{Vect}(\mathbb{R} \times Z)$$

denote the translational vector field on  $\mathbb{R}$  resp. rotational vector field on  $Z$ . The almost complex structure  $J$  is determined on the vertical part of  $TX$  by

$$J\partial_s = \partial_\theta, \quad J\partial_\theta = -\partial_s \text{ in } \text{Vect}(Z).$$

On the other hand, the projection to  $Y$  is required to be almost complex:

$$Dp_X J = \bar{J} Dp_X \text{ in } \text{Map}(TX, TY).$$

Suppose now  $X$  is an arbitrary cobordism from  $Z_-$  to  $Z_+$ . An almost complex structure  $J$  on  $X$  is called *cylindrical* if it is the restriction of cylindrical almost complex structures on  $\mathbb{R} \times Z_\pm$  on the cylindrical ends  $\pm(0, \infty) \times Z_\pm \rightarrow X$ . The space of cylindrical almost complex structures is denoted  $\mathcal{J}_{\text{cyl}}(X)$ . This ends the Definition.

We recall some notions of energy for maps from a punctured surface  $S$ .

**Definition 3.4.** (a) (Horizontal energy) The *horizontal energy* of a holomorphic map  $u = (\phi, v) : (S, j) \rightarrow (\mathbb{R} \times Z, J)$  bounding  $\mathbb{R} \times \Lambda$  is ([9, 5.3])

$$E^{\text{h}}(u) = \int_S v^* \omega.$$

(b) (Vertical energy) The *vertical energy* of a holomorphic map  $u = (\phi, v) : (S, j) \rightarrow (\mathbb{R} \times Z, J)$  is ([9, 5.3])

$$(7) \quad E^{\text{v}}(u) = \sup_{\zeta} \int_S (\zeta \circ \phi) d\phi \wedge v^* \alpha$$

where the supremum is taken over the set of all non-negative  $C^\infty$  functions

$$\zeta : \mathbb{R} \rightarrow \mathbb{R}, \quad \int_{\mathbb{R}} \zeta(s) ds = 1$$

with compact support.

(c) (Hofer energy) The *Hofer energy* of a holomorphic map

$$u = (\phi, v) : (S, j) \rightarrow (\mathbb{R} \times Z, J)$$

is ([9, 5.3]) is the sum

$$E(u) = E^{\text{h}}(u) + E^{\text{v}}(u).$$



- (d) (Generalization to manifolds with cylindrical ends) Suppose that  $X$  is a manifold with cylindrical ends modelled on  $\pm(0, \infty) \times Z_{\pm}$ . The vertical energy  $E^v(u)$  is defined as before in (7). The Hofer energy  $E(u)$  of a map  $u : S^{\circ} \rightarrow X^{\circ}$  from a surface  $S^{\circ}$  with cylindrical ends to  $X^{\circ}$  is defined by dividing  $X^{\circ}$  into a compact piece  $X^{\text{com}}$  and cylindrical ends diffeomorphic to  $\pm(0, \infty) \times Z_{\pm}$ . Then we set

$$E(u) = E(u|X^{\text{com}}) + E(u|(0, \infty) \times Z_+) + E(u|(0, \infty) \times Z_-).$$

In symplectic field theory, the compactification of punctured holomorphic curves is given by *holomorphic buildings* in which bubbling occurs on the cylindrical ends. Let  $(S, \partial S, j)$  be a bordered Riemann surface, with punctures in the interior and on the boundary. Let  $(X, J)$  be an almost complex manifold. A smooth map  $u : S \rightarrow X$  is *pseudoholomorphic* if the antiholomorphic part of the derivative vanishes:

$$\bar{\partial}_J u := \frac{1}{2}(du + Jduj) = 0.$$

Now suppose that  $X$  is a cobordism from  $Z_-$  to  $Z_+$ , and  $J$  is an almost complex structure on  $X$  that is cylindrical on the ends.

**Definition 3.5.** For integers  $k_-, k_+ \geq 0$  define topological spaces  $\mathbb{X}[k_-, k_+]$  by adding on  $k_-$  resp.  $k_+$  incoming resp. outgoing neck pieces:

$$(8) \quad \mathbb{X}[k_-, k_+] := (\overline{\mathbb{R} \times Z_-}) \cup_{Y_-} \dots \cup_{Y_-} (\overline{\mathbb{R} \times Z_-}) \cup_{Y_-} \overline{X} \\ \cup_{Y_+} (\overline{\mathbb{R} \times Z_+}) \cup_{Y_+} \dots \cup_{Y_+} (\overline{\mathbb{R} \times Z_+}).$$

Denote by  $\overline{\mathbb{X}}[k_-, k_+]_k$  the  $k$ -th topological space in the union above, and  $\mathbb{X}[k_-, k_+]_k$  the complement of the copies of  $Y_{\pm}$ . Thus  $\mathbb{X}[k_-, k_+]_k$  is either  $\mathbb{R} \times Z_{\pm}$  if  $k \neq 0$  or  $X$  if  $k = 0$ . Let  $L$  be a Lagrangian cobordism from  $\Lambda_-$  to  $\Lambda_+$ . Let

$$(9) \quad \overline{L}[k_-, k_+] := \overline{\mathbb{R} \times \Lambda_-} \cup_{\Pi_-} \dots \cup_{\Pi_-} \overline{\mathbb{R} \times \Lambda_-} \cup_{\Pi_-} \overline{L} \\ \cup_{\Pi_+} \overline{\mathbb{R} \times \Lambda_+} \cup_{\Pi_+} \dots \cup_{\Pi_+} \overline{\mathbb{R} \times \Lambda_+}$$

be the union of the closure  $\overline{L}$  with  $k_-, k_+$  cylindrical pieces attached to the ends  $\Pi_{\pm}$ .

A *holomorphic  $(k_-, k_+)$ -building* in  $X$  bounding  $L$  is a continuous map of a nodal curve  $S$  to  $\mathbb{X}[k_-, k_+]$  with the following properties: For each irreducible component  $S_v$ , let  $S_v^{\circ}$  denote the complement of the nodes. Then the restriction  $u_v$  of  $u$  to  $S_v$  is a holomorphic map to some component  $\mathbb{X}[k_-, k_+]_{k(v)}$

$$u_v : S_v \rightarrow \mathbb{X}[k_-, k_+]_{k(v)}, \quad v \in \text{Vert}(\Gamma)$$

with finite Hofer energy, and for each node  $w$  of  $S$  connecting components  $S_{v_-}$  and  $S_{v_+}$  and mapping to some copy of  $Y$ , the limiting Reeb chord or orbit of  $u_{v_-}$  at  $w$  is equal to limiting Reeb chord or orbit of  $u_{v_+}$  at  $w$ . We write  $u : S \rightarrow \mathbb{X}$  for short.

An *isomorphism* between buildings  $u_0, u_1 : S_0, S_1 \rightarrow \mathbb{X}[k_-, k_+]$  is an isomorphism of nodal curves  $\phi : S_0 \rightarrow S_1$  and a combination of translations  $\psi : \mathbb{X}[k_-, k_+]$  on the end pieces so that  $u_1 \circ \phi := \psi \circ u_0$ .

A *trivial cylinder or strip* is a map  $u : S \rightarrow \mathbb{R} \times Z$  mapping to a fiber  $Z_y \cong \mathbb{C}^\times$  of the projection to  $Y$ , with domain  $S$  either an  $\mathbb{R} \times S^1$  or  $\mathbb{R} \times [0, 1]$ , and mapping to  $Z_y$  by an exponential map  $(s, t) \mapsto e^{\mu(s+it+c)}$  for some exponential growth constant  $\mu$  and constant  $c \in \mathbb{C}$ . Any trivial cylinder  $u$  is contained in a fiber of  $p_Y$  and has exactly two punctures. Indeed, the action sums at the incoming and outgoing punctures are equal as in Lemma 3.16.

A holomorphic building  $u$  is *stable* if  $u$  has finitely many automorphisms, or equivalently the map  $u_0$  to the component  $\mathbb{X}[k_-, k_+]_0 \cong \overline{X}$  is stable in the usual sense, and the map  $u_j, j \neq 0$  to each component  $\mathbb{X}[k_-, k_+]_j = \overline{\mathbb{R} \times Z_\pm}$  has at least one connected component  $u_v$  that is not a trivial cylinder or strip.

*Remark 3.6.* Since our stable Hamiltonian manifolds are always circle-fibered, we can consider both the open manifolds and the compact manifolds given by collapsing the circle action. The nodal surfaces should be thought as surfaces before removing boundary and interior punctures.

**3.3. Treed pseudoholomorphic buildings.** Treed disks, also known as cluster configurations, were introduced in the context of Lagrangian Floer homology by Cornea-Lalonde [20]. The domains for treed disks are combinations of disks with strip-like or cylindrical ends and line segments.

**Definition 3.7.** An *closed interval* is a closed connected subset of  $\{-\infty\} \cup \mathbb{R} \cup \{\infty\}$ . Let  $I_1, \dots, I_k$  be a sequence of closed intervals so that  $I_1, \dots, I_{k-1}$  contain  $\infty$ , and  $I_2, \dots, I_k$  contain  $-\infty$ . A *broken interval* is a topological space  $I$  obtained from the union of closed intervals  $I_1, \dots, I_k$  by identifying  $\infty$  in one interval with  $-\infty$  in the next:

$$I = I_1 \cup_{\infty \sim -\infty} I_2 \cup_{\infty \sim -\infty} \dots \cup_{\infty \sim -\infty} I_k.$$

The images of the identified points  $\pm\infty$  in each subinterval are called *breakings*, while the boundary points of  $I$  are *ends* and may be either infinite or non-infinite. The standard metric the complement of infinities in each  $I_j$  induces a metric on the complement in  $I$  of the breakings.

A *treed disk*  $C$  is obtained from a nodal disk  $S$  by replacing each node  $w_\pm(e) \in \partial S$  or marking  $w(e) \in \partial S$  with a broken interval  $T_e$ ; in the case of a marking the corresponding broken interval  $T_e$  is required to have one infinite end and be joined to  $S$  at the finite end; while in the case of a node both ends of the broken interval are required to be finite (although the length of the broken interval may be infinite).

The *combinatorial type*  $\Gamma$  of any treed disk is the graph

$$(\text{Vert}(\Gamma) = \text{Vert}_\circ(\Gamma) \cup \text{Vert}_\bullet(\Gamma), \quad \text{Edge}(\Gamma) = \text{Edge}_\circ(\Gamma) \cup \text{Edge}_\bullet(\Gamma))$$

whose vertices correspond to disk and sphere components, respectively, and whose edges correspond to boundary resp. interior edges. The set of semi-infinite edges is denoted

$$\text{Edge}_\rightarrow(\Gamma) \subset \text{Edge}(\Gamma).$$

The type  $\Gamma$  has the additional data of subsets

$$\text{Edge}_0(\Gamma), \text{Edge}_\infty(\Gamma) \subset \text{Edge}(\Gamma) - \text{Edge}_\rightarrow(\Gamma)$$

describing which edges have zero or infinite length. That is,  $\text{Edge}_\infty(\Gamma)$  consists of those edges that are broken intervals, while  $\text{Edge}_0(\Gamma)$  consists of edges of zero length that are collapsed under the equivalence relation  $\sim$  in (10). See [14, Section 4.2] for more discussion of the combinatorial type. Any treed disk

$$C = \bigcup_{v \in \text{Vert}(\Gamma)} S_v \cup \bigcup_{e \in \text{Edge}(\Gamma)} T_e$$

is the union of disks and spheres  $C_v$  for the vertices  $v \in \text{Vert}(\Gamma)$  and interior and boundary edges  $T_e$  for the edges  $e \in \text{Edge}(\Gamma)$ .

A treed disk  $C$  is *stable* if the surface components  $S_v$  with markings given by  $S_v \cap T$  is the union of stable disks and the each broken interval  $T_e$  has at most one breaking. This ends the Definition.




FIGURE 5. A punctured disk and two treed punctured disks

The moduli space of stable treed disks with a fixed number of punctures has the structure of a compact Hausdorff space as in Charest-Woodward [14]. Let  $\mathcal{M}_\Gamma$  denote the moduli space of treed bordered surfaces of type  $\Gamma$ . If  $\Gamma$  is a type with an edge  $e \in \text{Edge}_0(\Gamma)$ , we declare  $\Gamma$  to be equivalent to the type  $\Gamma'$  obtained by removing the edge  $e$  and identifying the adjacent vertices  $v_\pm \in \text{Vert}(\Gamma)$ . Denote by

$$(10) \quad \overline{\mathcal{M}} = \sqcup_\Gamma \mathcal{M}_\Gamma / \sim$$

the union over stable types  $\Gamma$ , possibly disconnected, modulo the edge-collapse equivalence relation. The space  $\overline{\mathcal{M}}$  has a natural Hausdorff topology, induced by convergence of the surface components in the moduli space of marked disks and convergence of the edge lengths, as in Cornea-Lalonde [20]. Denote by  $\mathcal{M} \subset \overline{\mathcal{M}}$  the subset of strata of top dimension, consisting of configurations  $C$  so that each edge  $T_e$  has




FIGURE 6. Treed building

non-zero length  $\ell(e)$  in  $(0, \infty)$ . The set of types has a natural partial order with  $\Gamma' \preceq \Gamma$  if and only if  $\mathcal{M}_{\Gamma'}$  is contained in the closure of  $\mathcal{M}_{\Gamma}$ .

We will also consider domains with components labelled by integers, describing which *level* they map to the language of symplectic field theory.

**Definition 3.8.** A genus zero *building* is a nodal disk  $S$  equipped with a partition

$$(11) \quad S = S_1 \cup \dots \cup S_k$$

into (possibly disconnected) components  $S_1, \dots, S_k \subset S$  called *levels* so that intersection between two levels  $S_i \cap S_j$  empty unless  $i = j \pm 1$ . Removing the intersections  $S_i \cap S_{i \pm 1}$  gives a surface with cylindrical and strip-like ends

A *treed building* is a treed disk  $C$  equipped with a decomposition

$$C = C_1 \cup \dots \cup C_k$$

so that  $C_i, C_j$  intersect only if  $j \in \{i - 1, i, i + 1\}$ . See Figure 6.

The *combinatorial type* of a treed building is a graph

$$\Gamma = (\text{Vert}(\Gamma), \text{Edge}(\Gamma))$$

equipped with a decomposition into not-necessarily-connected subgraphs

$$\Gamma = \Gamma_1 \cup \dots \cup \Gamma_k$$

so that  $\Gamma_i \cap \Gamma_j$  is empty unless  $j \in \{i - 1, i, i + 1\}$ .

Treed holomorphic buildings in symplectic cobordisms are obtained by adding trajectories of the following kind. Let  $Z$  be a fibered contact or stable Hamiltonian manifold, and  $\Lambda \subset Z$  a Legendrian. We assume that a metric is fixed on  $\Lambda$ , hence on  $\mathcal{R}(\Lambda)$  using the identification of components of  $\mathcal{R}(\Lambda)$  with  $\Lambda$  in Lemma 2.9.

**Definition 3.9.** A *Morse datum* for  $(Z, \Lambda)$  consists of a pair of vector fields on the space of Reeb chords and on the Legendrian

$$\zeta_\circ \in \text{Vect}(\mathcal{R}(\Lambda)), \quad \zeta_{\bullet, \Lambda} \in \text{Vect}(\mathbb{R} \times \Lambda)^\mathbb{R}$$

arising as follows:

- (a) There exists a Morse function on the space of Reeb chords

$$f_{\circ} : \mathcal{R}(\Lambda) \rightarrow \mathbb{R};$$

so that  $\zeta_{\circ}$  is the gradient vector field:

$$(12) \quad \zeta_{\circ} := \text{grad}(f_{\circ}) \in \text{Vect}(\mathcal{R}(\Lambda)).$$

- (b) There exists a Morse function

$$f_{\bullet} : \Lambda \rightarrow \mathbb{R};$$

with gradient vector field

$$\text{grad}(f_{\bullet}) \in \text{Vect}(\Lambda)$$

so that  $\zeta_{\bullet, \Lambda}$  is a translation-invariant lift of  $\text{grad}(f_{\bullet})$ .

Since each component of the space of Reeb chords is isomorphic to the Legendrian, we could take the vector fields to be equal under the identification of the various components; however, this assumption is not necessary.

**Definition 3.10.** A vector field  $\zeta_{\bullet, \Lambda} \in \text{Vect}(\mathbb{R} \times \Lambda)$  is *positive* if in coordinates  $(s, \lambda)$  on  $\mathbb{R} \times \Lambda$  there exists a function

$$a : \Lambda \rightarrow \mathbb{R}_{>0}$$

so that

$$(13) \quad \zeta_{\bullet, \Lambda} = a(\lambda) \partial_s + p^* \text{grad}(f_{\bullet})$$

where

$$p^* : \text{Vect}(\Lambda) \rightarrow \text{Vect}(\mathbb{R} \times \Lambda)^{\mathbb{R}}$$

is the obvious identification of translationally-invariant vector fields trivial in the  $\mathbb{R}$ -direction with vector fields on  $\Lambda$ .

The limit of any Morse trajectory on a compact manifold along any infinite length trajectory is a zero of the gradient vector field. We introduce labels for the possible limits of the trajectories above as follows. Denote by

$$\underline{\mathbb{R}} = (T\mathbb{R}) \times \Lambda$$

the translational factor in  $T(\mathbb{R} \times \Lambda) = T\mathbb{R} \oplus T\Lambda$ . The zeroes of the vector field  $p_*(\zeta_{\bullet, \Lambda})$  correspond to tangencies of  $\zeta_{\bullet, \Lambda}$  with the translational factor:

$$p_*(\zeta_{\bullet, \Lambda})^{-1}(0) = \zeta_{\bullet, \Lambda}^{-1}(\underline{\mathbb{R}}) \subset \mathbb{R} \times \Lambda.$$

Let

$$(14) \quad \mathcal{I}(\Lambda) := \mathcal{I}_{\circ}(\Lambda) \cup \mathcal{I}_{\bullet}(\Lambda), \quad \mathcal{I}_{\circ}(\Lambda) := \zeta_{\circ}^{-1}(0), \quad \mathcal{I}_{\bullet}(\Lambda) := \zeta_{\bullet, \Lambda}^{-1}(0)$$

be the set of zeroes of these vector fields; these will be the generators of our Chekanov-Eliashberg algebras. The inclusion of the generators  $\mathcal{I}_{\bullet}(\Lambda)$  is similar to the inclusion of the chains on the Lagrangian in the definition of immersed Lagrangian Floer theory, while the generators  $\mathcal{I}_{\circ}(\Lambda)$  correspond to the self-intersections.

Treed holomorphic disks in cobordisms are combinations of pseudoholomorphic maps and trajectories of the vector fields above. Let  $(X, L)$  be a cobordism pair from  $(Z_-, \Lambda_-)$  to  $(Z_+, \Lambda_+)$ . Suppose  $\mathbb{R} \times Z_{\pm}$  are equipped with cylindrical almost

complex structures  $J_{\pm}$  and  $\mathbb{R} \times \Lambda_{\pm}$  are equipped with vector fields  $\zeta_{\pm} \in \text{Vect}(\mathbb{R} \times \Lambda_{\pm})$  as above.

**Definition 3.11.** Let  $X$  be a symplectic cobordism from  $Z_-$  to  $Z_+$ . An almost complex structure

$$J : TX \rightarrow TX$$

is *cylindrical-near-infinity* if  $J$  restricts to cylindrical almost complex structure  $J_{\pm}$  on the ends

$$\kappa_{\pm}(\pm(0, \infty) \times Z_{\pm}) \subset X.$$

A function resp. gradient vector field

$$f_L : L \rightarrow \mathbb{R}, \quad \zeta_{\bullet, L} = \text{grad}(f_L) \in \text{Vect}(L)$$

is *cylindrical near infinity* if  $\text{grad}(f_L)$  if the restriction of the vector field to the cylindrical ends is given by

$$\zeta_{\bullet, L}|_{\kappa_{\pm}(\pm(0, \infty) \times Z_{\pm})} = \zeta_{\bullet, \Lambda_{\pm}}$$

for some vector fields  $\zeta_{\bullet, \pm}$  of the form (13).

A *decorated cobordism pair* is a cobordism pair  $(X, L)$  together with a pair  $(J, f_L)$  as above.

In order to use the Cieliebak-Mohnke perturbation, we need pick some Donaldson type hypersurface  $D$  in  $X - L$ . This will be discussed in detail in Section 4.1. Here we only use it to define combinatorial types of treed holomorphic disks.

**Definition 3.12.** Let  $(X, L)$  be a decorated cobordism pair from  $(Z_-, \Lambda_-)$  to  $(Z_+, \Lambda_+)$  as above, so that the Lagrangians  $\Pi_{\pm} = p_{\pm}(\Lambda_{\pm})$  are embedded.

A *treed holomorphic disk* from a treed disk  $C$  to  $X$  is a collection of holomorphic maps and trajectories

$$\begin{aligned} u_v : S_v &\rightarrow X \quad v \in \text{Vert}(\Gamma) \\ u_e : T_e &\rightarrow \begin{cases} L & e \in \text{Edge}_L(\Gamma) \\ \mathcal{R}(\Lambda_{\pm}) & e \in \text{Edge}_{o, \pm}(\Gamma) \\ D & e \in \text{Edge}_D(\Gamma) \end{cases} \end{aligned}$$

for the vector fields  $\zeta_L, \zeta_{o, \pm}, \zeta_D = 0$  for some partition of the edges  $\text{Edge}(\Gamma)$  into subsets

$$\text{Edge}_D(\Gamma), \text{Edge}_L(\Gamma), \text{Edge}_{o, \pm}(\Gamma) \subset \text{Edge}(\Gamma)$$

so that the boundary conditions

$$u_v(\partial S_v) \subset L, \quad \forall v \in \text{Vert}_o(\Gamma)$$

are satisfied and matching conditions hold at the end of each edge according to

$$\text{ev}_z(u_v) = \text{ev}_z(u_e), \quad \forall z \in T_e \cap \overline{S}_v, \quad \forall e \in \text{Edge}(\Gamma), \quad v \in \text{Vert}(\Gamma).$$

We denote by  $T_D$  resp.  $T_L$  resp.  $T_{o, \pm} \subset T$  the union of the edges  $T_e$  with  $e \in \text{Edge}_D(\Gamma), \text{Edge}_L(\Gamma), \text{Edge}_{o, \pm}(\Gamma)$ .

A *treed holomorphic building* in  $\mathbb{X}[k_-, k_+]$  is a treed disk  $C$  equipped with a decomposition  $C = C_{k_-} \cup \dots \cup C_{k_+}$  together with treed disks bounding  $L$  or  $\mathbb{R} \times \Lambda_{\pm}$

$$\begin{aligned} u_j : C_j &\rightarrow \mathbb{X}[k_-, k_+]_j := \mathbb{R} \times Z_- \quad j = k_-, \dots, -1 \\ u_0 : C_0 &\rightarrow \mathbb{X}[k_-, k_+]_0 := X \\ u_j : C_j &\rightarrow \mathbb{X}[k_-, k_+]_j := \mathbb{R} \times Z_+ \quad j = 0, \dots, k_+ \end{aligned}$$

satisfying

- (a) (Balancing conditions) For any two adjacent  $j, j+1$  there exists  $\lambda_j \in [0, \infty]$  such that the following holds: Suppose  $T_e$  is an edge connecting  $C_j$  and  $C_{j+1}$  and  $u|_{T_e} : T_e \rightarrow \mathcal{R}(\Lambda)_{\pm}$  is a trajectory on the space of Reeb chords. Then the length of  $T_e$  is

$$\ell(T_e) = \lambda_j, \quad \forall T_e \text{ connecting } C_j, C_{j+1}.$$

- (b) (Matching conditions) If  $u_i, u_{i+1}$  are adjacent levels joined by an edge  $T_e = T_{e,i} \cup T_{e,i+1}$  with coordinate  $s$  on  $T_{e,i}, T_{e,i+1}$  then

$$\lim_{s \rightarrow \infty} u_i(s) = \lim_{s \rightarrow -\infty} u_{i+1}(s).$$

This limit should be interpreted as an element of the space  $\mathcal{R}(\Lambda_{\pm})$  of Reeb chords, if the trajectory represents a trajectory on  $\mathcal{R}(\Lambda_{\pm})$ , or an element of  $\Lambda_{\pm}$  if the trajectory represents a trajectory on  $\mathcal{R}(\Lambda_{\pm})$ .

An *isomorphism* of treed buildings  $u', u''$  with  $k$  levels is an isomorphism of domains  $\phi : C' \rightarrow C''$  (holomorphic on the surface parts and length-preserving on the segments) together with translations

$$\tau_j : \mathbb{X}[k_-, k_+]_j \rightarrow \mathbb{X}[k_-, k_+]_j, j = 1, \dots, k$$

so that

$$u''_v \circ \phi = \tau_j \circ u'_v$$

for each vertex  $v \in \text{Vert}(\Gamma_j)$ .

The *map type* of a treed building  $u$  is the decorated graph

$$\mathbb{F} = (\Gamma, h, l)$$

where

$$h : \text{Vert}(\Gamma) \rightarrow \pi_2(\overline{X}, \overline{L})$$

(with  $\overline{X}$  resp.  $\overline{L}$  the compactification of  $X = \mathbb{R} \times Z$  resp.  $L = \mathbb{R} \times \Lambda$ ) is the collection of homotopy classes of maps associated to the vertices and

$$l : \text{Edge}_{\rightarrow}(\Gamma) \rightarrow \mathcal{I}(\Lambda) \cup \{D\}$$

is the collection of labels on the leaves; that is, the constraints  $l(e) \in \mathcal{I}(\Pi)$  on constrained edges; or  $D$  for the edge  $T_e, e \in \text{Edge}_D(\Gamma)$  constrained to map to  $D$ .

We introduce the following notation for moduli spaces with fixed limits along the leaves. Denote by  $\mathcal{M}_{\mathbb{F}}(L)$  the moduli space of stable treed buildings bounding  $L$  of type  $\mathbb{F}$ , and  $\mathcal{M}(L)$  the union over types  $\mathbb{F}$ . In case  $L = \mathbb{R} \times \Lambda$ , we use the notation  $\mathcal{M}(\Lambda) = \mathcal{M}(\mathbb{R} \times \Lambda)$ . For each type  $\mathbb{F}$  (possibly disconnected) let  $\mathcal{M}_{\mathbb{F}}(L)$  denote the moduli space of isomorphism classes of finite energy holomorphic maps  $u : C \rightarrow X$

bounding  $L$  of map type  $\mathbb{F}$ , and  $\mathcal{M}_\Gamma(L)$  is the union of  $\mathcal{M}_{\mathbb{F}}(L)$  over map types  $\mathbb{F}$  whose underlying domain type is  $\Gamma$ . Denote

$$\mathcal{I}(L) = \zeta_{\bullet, L}^{-1}(0) = \text{crit}(f_L)$$

denote the zeroes of  $\zeta_{\bullet, L}$ . Taking the limit of any trajectory along a semi-infinite edge  $e$  defines an *evaluation map*

$$\text{ev}_e : \mathcal{M}_{\mathbb{F}}(L) \rightarrow \mathcal{I}(\Lambda_-)^{d_-} \cup \mathcal{I}(L)^d \cup \mathcal{I}(\Lambda_+)^{d_+}.$$

For any given collection

$$\underline{\gamma} = (\underline{\gamma}_-, \underline{\gamma}_\bullet, \underline{\gamma}_+) \in \mathcal{I}(\Lambda_-)^{d_-} \times \mathcal{I}(L)^d \times \mathcal{I}(\Lambda_+)^{d_+}$$

of generators  $\mathcal{I}(\Lambda_\pm)$  and critical points  $\underline{\gamma}_\bullet$  in the interior, let

$$\mathcal{M}(L, \underline{\gamma}) \subset \mathcal{M}(L)$$

the moduli space of treed disks with limits along the leaves given by  $\underline{\gamma}$ . If the domain  $C$  is disconnected, then we assume the set of components  $C_1, \dots, C_k$  of the domain is equipped with an ordering and the incoming and outgoing labels  $\underline{\gamma}_\pm$  are compatible with the ordering in the sense that the incoming labels for each component are in cyclic order around the boundary and the labels for the various components are in the same order; similarly for the outgoing labels. Also, the incoming labels are ordered before the outgoing labels. This ends the Definition.

**3.4. Monotonicity assumptions.** We make several assumptions on the cobordism to allow us to restrict to Chekanov-Eliashberg differentials counting rational curves with one incoming boundary puncture. These assumptions guarantee that spherical components do not bubble off in one-dimensional components of the moduli space: Such bubbling would force us to include Reeb orbits in the set of generators of the Chekanov-Eliashberg complex.

Our assumptions also force each component of the domain of a rigid treed disk to have at least one incoming puncture; otherwise, we would have to consider curves with multiple incoming punctures and the resulting differential would no longer respect the algebra structure given by concatenation of words.

Notably, we do not put any monotonicity assumption on the Lagrangian projection of the Legendrian, or on the Lagrangian cobordisms we consider. Possible disk bubbles will be studied with the theory of bounding chains in the sequel [7].

**Definition 3.13.** A fibered contact manifold  $(Z, \alpha)$  with base  $(Y, -\text{curv}(\alpha))$  is *tame* if the curvature is a negative multiple of an integral symplectic form  $\omega_Y \in \Omega^2(Y, \mathbb{Z})$ : for some constant  $\tau_Z \geq 1$

$$-\text{curv}(\alpha) = \tau_Z \omega_Y$$

and the base is monotone in the sense that there exists a *monotonicity constant*  $\tau_Y \geq 3$  so that

$$c_1(Y) = \tau_Y [\omega_Y] \in H^2(Y, \mathbb{R}).$$

For any compact spin Legendrian  $\Lambda \subset Z$  we call  $(Z, \Lambda)$  a *tame pair*.

The following is immediate from the Definition:



**Lemma 3.14.** *Suppose that  $(Z, \alpha)$  is tame. For any  $J$ -holomorphic map  $u : C \rightarrow Y$ , the Chern numbers satisfy*

$$\int_C u^* c_1(Y) \geq 2, \quad \int_C u^* c_1(Z) \geq 1.$$

*Example 3.15.* The odd sphere  $S^{2n-1}$  fibered over base  $\mathbb{C}P^{n-1}$  has  $\tau_Z = 1$  and  $\tau_Y = n$ , and so is tame if  $n - 1 \geq 2$ .

**Lemma 3.16.** *Let  $(Z, \alpha)$  be a fibered contact/stable-Hamiltonian manifold and  $\Lambda \subset Z$  a Legendrian. Let  $u : S \rightarrow \mathbb{R} \times Z$  be a punctured surface with boundary mapping to  $\mathbb{R} \times \Lambda$  and limiting to Reeb chords and orbits  $\gamma_e$  on the ends  $e \in \mathcal{E}(S)$  with actions  $\theta_e \in \mathbb{R}$ . Then the incoming and outgoing actions are related by*

$$\sum_{e \in \mathcal{E}_+(S)} \theta_e - \sum_{e \in \mathcal{E}_-(S)} \theta_e + \int_S u^* d\alpha = - \int_S u_Y^* \omega_Y.$$

*Proof.* Let  $\hat{S}$  denote the compactification of  $S$  obtained by adding in intervals along each strip-like end. Denote by  $u^* \alpha$  the pull-back connection form on the manifold with corners  $\hat{S}$ . By Stokes' formula, the difference in incoming and outgoing actions is given by

$$\sum_{e \in \mathcal{E}_+(S)} \theta_e - \sum_{e \in \mathcal{E}_-(S)} \theta_e + \int_{\partial S} u^* \alpha = \int_S du^* \alpha = - \int_S u_Y^* \omega_Y.$$

Here, the integral over  $\partial S$  vanishes since  $\alpha$  restricts to zero on  $\Lambda$ .  $\square$

The following corollary is used to show that the count of curves with a single input defines a differential.

**Corollary 3.17.** *Suppose that  $(Z, \Lambda)$  is a tame pair and  $J$  is a cylindrical almost complex structure on  $\mathbb{R} \times Z$ . Let  $u : S \rightarrow \mathbb{R} \times Z$  be a punctured  $J$ -holomorphic curve with boundary on  $\mathbb{R} \times \Lambda$ . The sum of the actions of Reeb chords at outgoing punctures is smaller than the sum of the actions at the incoming punctures:*

$$(15) \quad \sum_{e \in \mathcal{E}_+(S)} \theta_e \leq \sum_{e \in \mathcal{E}_-(S)} \theta_e$$

*with equality only if the map  $u$  projects to a constant map  $u_Y$  to  $Y$ . In particular, any non-constant punctured holomorphic curve  $u : S \rightarrow \mathbb{R} \times Z$  has at least one incoming puncture.*

*Proof.* The claim (15) follows from Lemma 3.16. To prove the last claim, let  $u$  be a map as in the statement of the Lemma. Suppose that  $u$  projects to a constant map  $u_Y$  but is non-constant in  $\mathbb{R} \times Z$ . The equation (15) implies that there are both positive and negative punctures. On the other hand, if  $u_Y$  is non-constant then (15) implies that there is at least one incoming puncture, that is, a point in  $\bar{u}^{-1}(Y_-)$ .  $\square$

We assume the following conditions on our cobordisms to guarantee the chain maps are well-defined. Recall that  $(Z_\pm, \alpha_\pm)$  are fibered contact/stable-Hamiltonian

manifolds with base the symplectic manifolds  $(Y_{\pm}, -d\alpha_{\pm})$ . Let  $(X, L)$  be a (cylindrical near infinity) cobordism pair from  $(Z_-, \Lambda_-)$  to  $(Z_+, \Lambda_+)$  in the sense of Definition 2.21. The normal bundle to  $Y_{\pm}$  in  $\overline{X}$  is the line bundle associated to  $Z_{\pm}$ .

**Definition 3.18.** Let  $\overline{X}, Y_{\pm}$  be as above. A *Thom form* for the inclusion  $Y_{\pm} \rightarrow \overline{X}$  is a two-form on  $\overline{X}$  given as follows. Let  $\rho : [0, \infty) \rightarrow [0, 1]$  be a compactly supported bump function equal to 1 in an open neighborhood of 0. The corresponding Thom form is

$$\text{Thom}_{Y_{\pm}} = -d(\rho\alpha_{\pm}) \in \Omega^2(\overline{X}, \overline{L})$$

where  $\Omega^2(\overline{X}, \overline{L})$  denotes the space of two-forms on  $X$  vanishing on each stratum of  $\overline{L}$ . The Thom form is closed and the *Thom class*  $[\text{Thom}_{Y_{\pm}}] \in H^2(\overline{X}, \overline{L})$  is independent of the choice of bump function  $\rho$ . The *logarithmic Chern class* of  $\overline{X} - Y_{\pm}$  is

$$c_1^{\log}(\overline{X} - Y_{\mp}) := c_1(\overline{X} - Y_{\mp}) - [Y_{\pm}]^{\vee} \in H^2(\overline{X} - Y_{\pm}).$$

**Definition 3.19.** The pair  $(X, L)$  is a *tame cobordism pair* from  $(Z_-, \Lambda_-)$  to  $(Z_+, \Lambda_+)$  if and only if

- (P1) (Integrality) The symplectic classes  $[\omega_Y] \in H^2(Y)$  and  $[\overline{\omega}] \in H^2(\overline{X})$  are integral; this is a technical assumption for our regularization scheme which presumably could be removed and the Lagrangians are rational in the sense of Definition 4.3 below;
- (P2) (No-cap condition) The logarithmic first Chern class of  $\overline{X} - Y_+$  is a very positive multiple of the symplectic class on the complement of  $Y_+$  in the sense that there exists a constant  $\lambda_- > 0$  so that

$$c_1^{\log}(\overline{X} - Y_+) = (1 + \lambda_-)[\overline{\omega}] \in H^2(\overline{X} - Y_+).$$

- (P3) (Monotonicity for the outgoing end) If the outgoing end  $\Lambda_+$  of the cobordisms  $L$  is non-empty then the relative Thom class of the outgoing end  $[Y_+]^{\vee} \in H^2(\overline{X} - Y_-, \overline{L} - Y_-)$  is non-positive in the sense that there exists a constant  $\lambda_+ \geq 0$  so that

$$[Y_+]^{\vee} = -\lambda_+[\overline{\omega}] \in H^2(\overline{X} - Y_-, \overline{L} - Y_-).$$

*Example 3.20.* In the case of the cobordism  $X = \mathbb{C}^n - \{0\} \cong \mathbb{R} \times S^{2n-1}$  from  $S^{2n-1}$  to itself and  $L$  is a cylinder  $\mathbb{R} \times \Lambda$ , the compactification  $\overline{X}$  is the blow-up of  $\mathbb{C}P^n$  at 0 with  $Y_{\pm} \cong \mathbb{C}P^{n-1}$ , the logarithmic first Chern class is

$$c_1^{\log}(\overline{X} - Y_+) = n[Y_-]^{\vee}$$

and the outgoing divisor has dual class

$$[Y_+]^{\vee} = -[\overline{\omega}] \in H^2(\overline{X} - Y_-, \overline{L} - Y_-).$$

Therefore the constants are

$$\lambda_- = n - 1, \quad \lambda_+ = 1,$$

hence the cobordism pair is tame if  $n > 2$ .

We now discuss our motivating example given by the Harvey-Lawson filling. According to Lemma 2.28, the Harvey-Lawson filling defines a cylindrical-near-infinity filling  $L \subset X$  of a Legendrian  $\Lambda_\epsilon$  with respect to some stable Hamiltonian triple  $(Z, \alpha_\epsilon, \omega_Z)$  with  $Z \cong S^{2n-1}$  and  $\Lambda_\epsilon$  a lift of a perturbation of the Clifford torus. Although the form  $\alpha_\epsilon$  for which  $\Lambda_\epsilon$  is Legendrian, is contact, the symplectic form  $\omega$  on  $X$  is an extension of  $\alpha_0$  so that  $d\alpha_0 = -\omega_Z$ , rather than the perturbed form  $\alpha_\epsilon$ . From now on, by the *Harvey-Lawson filling* we mean the filling  $L$  of  $\Lambda_\epsilon \subset S^{2n-1}$  produced by Lemma 2.28 applied to the Harvey-Lawson Lagrangian (2).

**Lemma 3.21.** *The Harvey-Lawson filling  $L \cong T^{n-2} \times \mathbb{R}^2$  of  $\Lambda_\epsilon \cong T^{n-1} \subset S^{2n-1}$  is tame.*

*Proof.* The compactified filling is  $\overline{X} = \mathbb{C}P^n$  which has Chern class

$$c_1(\overline{X}) = (n+1)[\mathbb{C}P^{n-1}]^\vee = (n+1)[Y_-]^\vee.$$

Hence

$$c_1(\overline{X}) - [Y_-]^\vee = n[Y_-]^\vee = (1 + \lambda_-)[\overline{\omega}]$$

for  $n \geq 2$ . The outgoing end  $\Lambda_+$  is empty, so the condition (P3) is vacuous. Hence  $(X, L)$  is a tame cobordism pair.  $\square$

We now give an example of a cobordism that is not tame. Let  $X$  be the blow-up of the unit ball  $B_1(0)$  in  $\mathbb{C}^n$ . Since the Harvey-Lawson filling  $L$  of  $\Lambda_\epsilon$  is disjoint from  $0 \in B_1(0)$ , it defines a Lagrangian filling in  $X$  as well, denoted with the same notation.

**Lemma 3.22.** *Let  $X, L$  be as above so that  $X$  is the blow-up of  $B_1(0)$  at 0. The Harvey-Lawson filling  $L \subset X$  is not tame as a filling.*

*Proof.* We check that the condition (P2) fails. The compactification  $\overline{X}$  is the blow-up  $\text{Bl}_0 \mathbb{C}P^n$  of  $\mathbb{C}P^n$  at  $0 \in \mathbb{C}^n \subset \mathbb{C}P^n$ . The space  $\overline{X}$  fibers over  $\mathbb{C}P^{n-1}$  via a map

$$\pi : \text{Bl}_0 \mathbb{C}P^n \rightarrow \mathbb{C}P^{n-1}.$$

The blow-up  $\text{Bl}_0 \mathbb{C}P^n \cong \mathbb{P}(\mathcal{O}(0) \oplus \mathcal{O}(1))$  is isomorphic to the projectivization of the sum of the hyperplane and trivial bundles  $\mathcal{O}(1), \mathcal{O}(0)$ . The Chern numbers of the fibers of the projection are

$$c_1(p^{-1}(\ell)) = 2, \quad \forall \ell \in \mathbb{C}P^{n-1}.$$

The evaluation of the logarithmic Chern class on a fiber is

$$\langle c_1^{\log}(\overline{X} - Y_+), [p^{-1}(\ell)] \rangle = \langle c_1(\overline{X}) - [Y_-], [p^{-1}(\ell)] \rangle = 2 - 1 = 1$$

which is less than the required bound in (P2).  $\square$

The moduli spaces of holomorphic disks in the example in Lemma 3.22 have boundary components where a sphere representing a fiber has bubbled off. Thus, for a well-defined theory of Legendrian contact homology in which such fillings define augmentations, one would have to include the Reeb orbits in the set of generators.

**Lemma 3.23.** *Let  $\Lambda$  be a Legendrian in a tame fibered contact manifold  $Z$ . Suppose that  $X = \mathbb{R} \times Z$  is the symplectization of  $Z$  and*

$$\overline{L} = (\mathbb{R} \times \Lambda)_{[\sigma_1, \sigma_2]} \cong \Pi \sqcup ((\sigma_1, \sigma_2) \times \Lambda) \sqcup \Pi$$

(where  $\Pi \cong \Pi_- \cong \Pi_+$ ) the symplectic cut at  $\sigma_1, \sigma_2$ . For choices of  $\sigma_1, \sigma_2 \in \mathbb{R}$  so that  $[e^{-\sigma_1}\omega_Y], [e^{-\sigma_2}\omega_Y] \in H_2(Y)$  are integral and  $e^{-\sigma_2} - e^{-\sigma_1} \in \mathbb{Z}$ ,  $(X, L)$  is a tame cobordism pair.

*Proof.* We check the integrality of the symplectic class. The degree two homology of  $\overline{X}$  is generated by the images of classes under either embedding  $Y \rightarrow \overline{X}$  together with classes in the fibers of  $\pi : \overline{X} \rightarrow Y$ , by the Leray-Serre spectral sequence (or, more simply, the Gysin sequence.) The fiber  $\pi^{-1}(y)$  over any  $y \in Y$  has area  $e^{-\sigma_2} - e^{-\sigma_1}$  which is an integer by assumption. On the other hand, the degree two homology classes in  $Y$  have integral areas by assumption. Hence, **(P1)** is satisfied.

To check the positivity of the logarithmic first Chern class, note that the complement  $\overline{X} - Y_+$  contracts to  $Y_-$ , compatible with the Lagrangian boundary condition, so it suffices to check **(P2)** on spheres in  $Y_-$ . The class  $c_1(\overline{X} - Y_+)$  restricts to  $(\tau_Y + \tau_Z)[\omega_{Y_-}]$  on  $Y_-$ , where it is equal to the restriction of  $(\tau_Y + \tau_Z)[\omega_Y]$ . So

$$c_1^{\log}(\overline{X} - Y_+) = c_1(\overline{X} - Y_+) - [Y_-]^\vee = p^*c_1(Y) = \tau_Y[\omega_Y].$$

So the condition **(P2)** holds with proportionality constant  $\lambda_- = \tau_Y + \tau_Z - 1$ .

Finally, we check the negativity of the outgoing end. We compute

$$[Y_+] = p^*c_1(Z) = -\tau_Z p^*[\omega_Y] = -\tau_Z[\overline{\omega}] \in H_2(\overline{X} - Y_-, \overline{L} - Y_-).$$

So **(P3)** holds with proportionality constant  $\lambda_+ = \tau_Z$ .  $\square$

The conditions in Definition 3.19 could presumably be relaxed by dealing with more complicated curve counts; for example, including Reeb orbits in the set of generators would allow relaxation of the second inequality while presumably allowing all-genus counts would allow relaxation of the first. However, allowing these possibilities introduces further technical difficulties. Note that the conditions here are weaker than the exact conditions in, for example, Chantraine [12].

**Lemma 3.24.** *Let  $(X, L)$  be a cobordism satisfying **(P3)**. Any holomorphic punctured disk  $u : S \rightarrow X$  bounding  $L$  with an outgoing strip-like end has at least one incoming end.*

*Proof.* We argue by contradiction. Let  $u : S \rightarrow X$  be a punctured disk bounding  $L$  with only outgoing ends. The intersection number of the map  $u$  with the outgoing divisor  $Y_+$  is

$$[u] \cdot [Y_+] = [u] \cdot \lambda_+[\omega] = -\lambda_+A(u) \leq 0.$$

Since  $u$  is holomorphic with non-positive area,  $u$  must be constant. Since by definition  $u$  maps to the interior  $X$ , the map  $u$  has no outgoing ends, which is a contradiction.  $\square$

We now explain the meaning of the no-cap condition **(P2)**. By a *rigid holomorphic sphere* we mean a map  $u : S \rightarrow X$  from a sphere  $S$  of some type  $\mathbb{I}$  which lies in a moduli space of holomorphic maps of expected dimension zero. In section 4.2 we

show that the moduli spaces made be made regular by suitable perturbations, so that in fact such a map is rigid if and only if it is isolated.

**Lemma 3.25.** *Suppose  $(X, L)$  is a tame cobordism. There are no rigid holomorphic spheres  $u : S \rightarrow X$  with a single puncture  $z \in \bar{S}$  constrained to map to a fixed Reeb orbit over  $Y_-$ .*

*Proof.* Consider a punctured sphere  $u : S \rightarrow X$  of type  $\mathbb{I}$  whose closure  $\bar{u}$  meets the incoming divisor  $Y_-$  but not the outgoing divisor  $Y_+$ , and  $\bar{u}$  its extension to a map to  $\bar{X}$ . The dimension of the moduli space  $\mathcal{M}_{\mathbb{I}}(X)$  of spheres constrained to intersect  $Y_-$  at a single puncture with multiplicity

$$m = (\bar{u}_*[\mathbb{P}^1], [Y_-])$$

is (similar to Cieliebak-Mohnke [17, Proposition 6.9])

$$\begin{aligned} \dim \mathcal{M}_{\mathbb{I}}(X) &= 2(c_1(\bar{X}), \bar{u}_*[\mathbb{P}^1]) - 2 - 2m \\ &= 2((1 + \lambda_-)[\bar{\omega}], \bar{u}_*[\mathbb{P}^1]) - 2 > 0. \end{aligned}$$

Here we used that  $X$  satisfies **(P2)**. Hence such spheres cannot be rigid. On the other hand, if  $\bar{u}$  has domain a sphere  $S$  with a single outgoing puncture and no incoming punctures then by **(P3)**

$$0 > -[Y_+].[\bar{u}] = \lambda_+[\bar{\omega}].[\bar{u}].$$

This is a contradiction.  $\square$

**3.5. Homological invariants.** In the definition of contact homology we will work with coefficient rings that are completed group rings for various homology groups. The standard in the field is to use a group ring on first homology. The standard approach is to assign homology classes to punctured disks by closing up the Reeb chords at infinity. Denote by

$$\Lambda_1, \dots, \Lambda_k \subset \Lambda$$

the connected components of the Legendrian  $\Lambda$ . For each connected component  $\Lambda_i$  of  $\Lambda$  choose a base point

$$\lambda_i \in \Lambda_i, \quad i = 1, \dots, k.$$

**Definition 3.26.** (a) (Capping paths) For each Reeb chord  $\gamma : [0, 1] \rightarrow Z$  a *capping path* is a path

$$(16) \quad \hat{\gamma}_b : [0, 1] \rightarrow \Lambda, \quad \hat{\gamma}_b(0) = \gamma(b), b \in \{0, 1\}, \quad \hat{\gamma}_b(1) = \lambda_i.$$

(b) (Boundary cycles) Let  $u : S \rightarrow \mathbb{R} \times Z$  be a holomorphic map bounding  $\mathbb{R} \times \Lambda$  with limits on the strip-like ends given by the collection of Reeb chords

$$\underline{\gamma} = (\gamma_0, \dots, \gamma_k) \in \mathcal{R}(\Lambda)^{k+1}$$

with capping paths  $\hat{\gamma}_{i,b}, i = 0, \dots, k+1$ . The boundary  $\partial S$  is divided into components

$$\partial S = (\partial S)_0 \cup \dots \cup (\partial S)_k$$

by the strip-like ends. The restriction  $(\partial u)_m$  of  $u$  to  $(\partial S)_m$  after projection to  $\Lambda$  concatenates with the capping paths for chords in  $\underline{\gamma}$  to produce a sequence of *boundary cycles*

$$\overline{\partial u}_m := \hat{\gamma}_{m+1,0} \circ (\partial u)_m \circ \hat{\gamma}_{m,1}^{-1} : S^1 \rightarrow \Lambda_{i(m)}, m = 0, \dots, k$$

in the components  $\Lambda_{i(m)}$ .

- (c) (Total homology) The total homology class of the combined boundary cycles is denoted

$$(17) \quad [\partial u] = \sum_k [\overline{\partial u}_j] \in H_1(\Lambda).$$

This ends the Definition.




FIGURE 7. Capping paths to make a cycle

*Remark 3.27.* Similarly, in the case that the Legendrian is connected and the contact manifold is simply-connected, second homology classes are associated to punctured disks by a capping procedure. For each Reeb chord, the loop obtained by concatenating  $\gamma, \hat{\gamma}_0, \hat{\gamma}_1$  is the boundary of some continuous map  $v_\gamma : S_\gamma \rightarrow Z$  from a disk  $S_\gamma$  to  $Z$  by the assumption that  $Z$  is simply-connected. The homology class obtained from a punctured disk  $u : S \rightarrow Z$  by gluing in the disk  $v_\gamma$  at each Reeb chord or orbit is denoted

$$(18) \quad [u] = \left[ u_*(S) + \sum_\gamma v_{\gamma,*} S_\gamma \right] \in H_2(Z, \Lambda)$$

This ends the Remark.

Finally, we define homology classes of disks in cobordisms.

**Definition 3.28.** Let  $(X, L)$  be a symplectic cobordism. Let  $\overline{L}$  denote the closure of  $L$  in the manifold with boundary  $\tilde{X}$ . For each component  $\tilde{L}_i$  of a cobordism  $\tilde{L}$  choose a choice of base point  $\lambda_i$  in each component and a collection of paths  $\hat{\gamma}_b, b \in \{0, 1\}$  from the base point to the start and end points of any Reeb chord  $\gamma$ . Let  $u : S \rightarrow X$

bounding  $L$  punctured holomorphic curve. By concatenating with the disks  $v_{\gamma_b}$  and paths  $\hat{\gamma}_b$  for each strip-like end of  $S$  one obtains homology classes denoted

$$[\partial u] \in H_1(\tilde{L}) \cong H_1(L).$$

If  $L$  is connected and  $X$  is simply connected, one obtains second homology classes

$$[u] \in H_2(\tilde{X}, \tilde{L}) \cong H_2(X, L)$$

by a choice of capping disk at each Reeb chord. If  $u = (u_1, u_2, \dots, u_k)$  is a holomorphic building with levels  $u_1, u_2, \dots, u_k$  then define

$$[\partial u] := [\partial u_1] + [\partial u_2] + \dots + [\partial u_k], \quad [u] := [u_1] + [u_2] + \dots + [u_k].$$

The *boundary homology class* of a treed holomorphic disk

$$u : C \rightarrow \mathbb{R} \times Z \quad \text{or} \quad u : C \rightarrow X$$

is defined as follows: In the first case of maps to  $\mathbb{R} \times Z$ , the boundary components  $(\partial S)_i, i = 1, \dots, l(i)$  of each disk component  $u_v : S_v \rightarrow X$  define paths

$$(\partial u_v)_i : (\partial S)_i \rightarrow \Lambda_{k(i)}, \quad i = 1, \dots, l(i)$$

where  $\Lambda_{k(i)} \subset \Lambda$  is some connected component. Each trajectory  $u_e : T_e \rightarrow \mathcal{R}(\Lambda)$  defines by composition a pair of maps

$$u_{e,b} : T_e \rightarrow \Lambda_{k_b(e)}, \quad b \in \{0, 1\}$$

by composing  $u_e$  with evaluation  $\mathcal{R}(\Lambda) \rightarrow \Lambda$  at the start or endpoint of the path. The boundary homology class  $[\partial u] \in H_1(\Lambda)$  is defined as the class of the sum of the chains above. In the case of a treed disk in  $X$  bounding a cobordism  $L$ , one obtains a class in  $H_1(\tilde{L}) \cong H_1(L)$  by a similar construction.

The *boundary class* of a treed building  $u : C \rightarrow \mathbb{X}$  is the element  $[\partial u] \in H_1(L)$  obtained by concatenating the restrictions  $\partial u_v$ , the compositions of the trajectories  $u_e : T_e \rightarrow \mathcal{R}(\Lambda)$  with the projections  $\mathcal{R}(\Lambda) \rightarrow \Lambda$  given by evaluation at the end point or starting point, and the capping paths to the base points  $\lambda_i$  in the corresponding components  $\Lambda_i$  of  $\Lambda$ . This ends the Definition.

#### 4. REGULARIZATION

In this section, we discuss the necessary Fredholm theory, compactness results, and regularization techniques to make the moduli spaces of treed buildings of dimension at most one compact manifolds with boundary. The compactness results in symplectic field theory are similar to those covered in [9], [1]. We take as a regularization scheme the stabilizing divisors scheme of Cieliebak-Mohnke [17]; the adaptation to the case at hand requires a small extension of the results in Pascaleff-Tonkonog [44] to the case of clean intersection. The reader could presumably substitute their favorite regularization scheme.

**4.1. Donaldson hypersurfaces.** To achieve manifold structures on the moduli spaces of treed holomorphic maps we use Cieliebak-Mohnke perturbations [17]. Since the compactification of our Lagrangians are only cleanly-intersecting, we extend the construction to that case, generalizing the construction in Pascaleff-Tonkonog [44].

**Definition 4.1.** A *Bertini hypersurface* in a complex projective subvariety  $M \subset \mathbb{C}P^{n-1}$  is the intersection of  $M$  with a generic hyperplane. By Bertini's theorem, such a hypersurface is a smooth subvariety.

A *Donaldson hypersurface* in a symplectic manifold  $M$  with symplectic form  $\omega$  is a codimension two symplectic submanifold  $D \subset M$  such that the dual cohomology class  $[D]^\vee$  satisfies

$$[D]^\vee = k[\omega] \in H^2(M)$$

for some positive  $k > 0$  called the *degree* of  $D$ .

Now we assume that  $X$  is a symplectic cobordism with circle-fibered boundary and let  $\bar{X}$  be its compactification, as in Section 3. Results of Cieliebak-Mohnke [17] and Charest-Woodward [13] imply that there exist Donaldson hypersurfaces which intersect holomorphic disks and spheres in only finitely many points:

**Proposition 4.2.** *There exists a  $k_0 \in \mathbb{Z}_+$  so that if  $D \subset \bar{X}$  is a Donaldson hypersurface  $D$  of degree at least  $k_0$  then there exists a tame almost complex structure on  $\bar{X}$  so that  $D$  contains no rational curves  $u : S^2 \rightarrow \bar{X}$ . Furthermore, for any such  $J_0$  and energy bound  $E$ , there exists an open neighborhood,*

$$\mathcal{J}^E(\bar{X}) \subset \mathcal{J}(\bar{X}),$$

*in the space of tame almost complex structures on  $\bar{X}$  with the property that if  $J_1 \in \mathcal{J}^E(\bar{X})$  then any non-constant  $J_1$ -holomorphic sphere  $u : S^2 \rightarrow \bar{X}$  of energy at most  $E$  meets  $D$  in at least three but finitely many points  $u^{-1}(D)$ .*

In the case that  $X$  is a complex projective variety, a result of Clemens [19] implies that we may take  $D$  to be a Bertini hypersurface, and leave the complex structure unperturbed.

We wish to find Donaldson hypersurfaces that intersect each non-constant punctured holomorphic disk. Let  $\bar{L} \subset \bar{X}$  be the closure of a Lagrangian cobordism, so that  $\bar{L}$  is the union of a smooth Lagrangian  $L$  and Lagrangians  $\Pi_\pm$  in  $Y_\pm$ , with angles between the branches at the self-intersections bounded from below, as in Definition 2.23.

**Definition 4.3.** A cleanly-self-intersecting Lagrangian  $\bar{L} \subset \bar{X}$  is *rational* if there exists a line-bundle-with-connection

$$\hat{X} \rightarrow \bar{X}$$

whose curvature is  $k\omega \in \Omega^2(\bar{X})$  for some integer  $k > 0$  such that the pull-back of  $\hat{X}$  to the Lagrangian  $\bar{L}$  has a flat section.



The Lagrangian  $L$  is *exact* on a subset  $U \subset X$  with  $\bar{U} \subset \bar{X}$  open if there exists a one-form  $\alpha \in \Omega^1(\bar{U})$  with  $d\alpha = \omega|_{\bar{U}}$  and a continuous function  $f : \bar{L} \rightarrow \mathbb{R}$  so that  $df|_L = \alpha|_L$ .

**Lemma 4.4.** *Suppose that  $\bar{X}$  is simply connected and every disk  $u : S \rightarrow \bar{X}$  bounding  $\bar{L}$  has rational area  $\langle [u], [\bar{\omega}] \rangle \in \mathbb{Q}$ . Then  $\bar{L}$  is rational.*

*Proof.* Let  $\hat{X}$  be a line bundle-with-connection with curvature  $k\omega$ . After replacing  $k$  with a sufficiently large integer multiple, every disk bounding  $\bar{L}$  has integer  $k\omega$ -area. It follows from the assumption that  $X$  is simply connected that the holonomy around any loop in  $\bar{L}$  is trivial, as in the computation in the proof of Proposition 2.5. Parallel transport of an arbitrary point in  $\hat{X}$  along paths in  $\bar{L}$  then defines the desired flat section  $s : \bar{L} \rightarrow \hat{X}^k|_{\bar{L}}$ .  $\square$

The idea of creating symplectic divisors in rational symplectic manifolds as introduced in [23] was to take zero locus of an “almost” holomorphic section of the line bundle  $\hat{X}^{\otimes k}$ . We recall the construction of Donaldson divisor in the complement of an isotropic submanifold as done in [6]. The idea is to find a sequence of almost holomorphic sections over tensor powers of the line bundle  $\hat{X}^{\otimes k}$  which is concentrated away from zero on the isotropic submanifold. Thus, the zero-locus of such a section will not intersect the isotropic submanifold. In [44, Theorem 3.3], it proves that one can chose a Donaldson divisor on the complement of two isotropic which are cleanly intersecting. We extend the ideas introduced there to construct Donaldson divisors on the complement of a branched rational Lagrangian.

**Proposition 4.5.** *Let  $\bar{L}$  be a branched rational Lagrangian submanifold then for large  $k$ , there exists a section  $\sigma_{k,L}$  of  $\hat{X}^{\otimes k}$  which is concentrated over  $\bar{L}$  in the sense of [6].*

In preparation for the proof, we cover the singular locus by Darboux balls as follows. Let  $\bar{L}$  be a branched rational Lagrangian. Choose Darboux charts at  $p \in L_\cap$  so that the intersection with  $L$  is the union of Lagrangian subspaces. We will use these charts to construct peak sections  $\sigma_{k,p}$  of  $\hat{X}^{\otimes k}$  as done in Auroux-Gayet-Mohsen [6]. Define

$$d_k : \bar{L} \times \bar{L} \rightarrow \mathbb{R}_{\geq 0}$$

to be the distance with respect to the scaled metric  $kg$  where  $g$  is the compatible metric on  $X$  induced from  $J$ . Since  $\bar{L}$  is a rational Lagrangian, there exists a flat section

$$\tau_k : \bar{L} \rightarrow \hat{X}^{\otimes k}|_{\bar{L}}.$$

We construct an approximately holomorphic section  $\sigma_{k,L_\cap}$  by the process described in [6]. Near any  $p \in L_\cap$ , there is the symplectomorphism  $\phi$  from Definition 2.23, under which  $\bar{L}$  is a union of  $k$  Lagrangian subspaces in the local model. Cover  $L_\cap$  with such Darboux balls

$$B_1, \dots, B_N \subset L_\cap$$

of radius 1 with respect to  $d_k$  which are centered  $\frac{2}{3}$  distance away from each other. Let

$$P_k \subset L_\cap$$

denote the center of the balls. Define  $\sigma_{k,L_\cap}$  as follows:

$$(19) \quad \sigma_{k,L_\cap} = \sum_{p \in P_k} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_{k,p} : X \rightarrow \hat{X}^k$$

**Lemma 4.6.** *For large enough integers  $k$ , if  $d_k(x, L_\cap) < k^{\frac{1}{30}}$  for  $x \in \bar{L}$ , then*

$$|\sigma_{k,L_\cap}(x)| > \frac{1}{2} e^{-2(d_k^2(x, L_\cap) + d_k(x, L_\cap))}.$$

*Proof.* We run through the construction of asymptotically holomorphic sections again, paying careful attention to the differences in phases. Fix some  $x \in \bar{L}$  such that  $d_k(x, L_\cap) < k^{\frac{1}{30}}$ . By the proof of Lemma 4 of [6] verbatim, when  $p \in L_\cap$  and  $x \in \bar{L}$  are at most  $k^{1/10}$  distance apart, their phases are close in the sense that

$$(20) \quad \left| \arg \left( \frac{\sigma_{k,p}(x)}{\tau_k(x)} \right) - \arg \left( \frac{\sigma_{k,p}(p)}{\tau_k(p)} \right) \right| \leq \frac{\pi}{4}.$$

Let  $p_f \in P_k$  be the center of the ball which contains the closest point in  $\mathcal{L}_\cap$  to  $x$ . For a large enough  $k$ , the distance from  $x$  to the center  $p_f$  satisfies the bound

$$d_k(p_f, x) < k^{\frac{1}{20}}.$$

By the proximity of the phases  $\arg(\frac{\sigma_{k,p}}{\tau_k})$  to each other in (20),

$$(21) \quad \left| \sum_{p \in P_k | d_k(x,p) < k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) \right| \geq |\sigma_{k,p_f}(x)|.$$

The cutoff function to define the section  $\sigma_{k,p_f}$  is equal to one at  $x$ . From the definition of  $p_f$  and the triangle inequality

$$d_k(x, L_\cap) + 1 > d_k(x, p_f).$$

Thus for any  $x \in \bar{L}$  with  $d_k(x, L_\cap) < k^{\frac{1}{30}}$ ,

$$(22) \quad |\sigma_{k,p_f}(x)| \geq e^{-d_k^2(x, p_f)} \geq e^{-(d_k(x, L_\cap) + 1)^2}.$$

Lemma 3 from [6] implies that there exists  $\tilde{c}, \tilde{r} > 0$  independent of  $k, P_k$  such that

$$(23) \quad \left| \sum_{p \in P_k | d_k(x,p) > k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) \right| \leq \tilde{c} e^{-\tilde{r} k^{\frac{1}{20}}} \quad \forall x \in X.$$

From the inequalities (22) and (23), we get that for large  $k$ ,

$$\begin{aligned}
|\sigma_{k,L_\cap}(x)| &= \left| \sum_{p \in P_k | d_k(x,p) < k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) + \sum_{p \in P_k | d_k(x,p) > k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) \right| \\
\text{so } |\sigma_{k,L_\cap}(x)| &\geq \left| \sum_{p \in P_k | d_k(x,p) < k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) \right| - \left| \sum_{p \in P_k | d_k(x,p) > k^{\frac{1}{20}}} \frac{\tau_k(p)}{\sigma_{k,p}(p)} \sigma_p(x) \right| \\
\text{so } |\sigma_{k,L_\cap}(x)| &\geq e^{-(d_k(x,L_\cap)+1)^2} - \tilde{c}e^{-\tilde{r}k^{1/10}} \\
\text{so } |\sigma_{k,L_\cap}(x)| &\geq \frac{1}{2}e^{-2(d_k^2(x,L_\cap)+d_k(x,L_\cap))}.
\end{aligned}$$

The last inequality follows from the chain of inequalities

$$\frac{1}{2}e^{-d_k^2(x,L_\cap)-d_k(x,L_\cap)} > \frac{1}{2}e^{-k^{1/15}-k^{\frac{1}{30}}} > \tilde{c}e^{-\tilde{r}k^{1/10}}$$

for large  $k$ . This gives the desired inequality

$$|\sigma_{k,L_\cap}(x)| \geq \frac{1}{2}e^{-2(d_k^2(x,L_\cap)+d_k(x,L_\cap))}.$$

□

*Proof of Proposition 4.5.* We aim to construct an approximately holomorphic section concentrated on the Lagrangian by summing the locally Gaussian sections  $\sigma_{k,L_\cap}$  as defined in Equation (19) and Lemma 4.6. We define  $\sigma_{k,L_i}$  by summing peak sections on the  $m$  Lagrangian branches  $L_i$ , while including the sections on the points  $P_k$  on  $L_\cap$ . We start with balls centered around  $P_k$  and extend it to a finite cover of  $L_i$  by radius  $\frac{1}{3}$  balls. Denote the set of centers of these balls as

$$P_k^{\text{pre}} \subset \bar{L}.$$

Iterated removal of points from  $P_k^{\text{pre}}$  ensures that any two points in  $P_k^{\text{pre}}$  are at least  $\frac{2}{3}$  distance apart. That is, if  $p, q \in P_k^{\text{pre}}$  are closer than  $\frac{2}{3}$  we drop either  $p$  or  $q$  from the set  $P_k^{\text{pre}}$ , without remove points from  $P_k$  while doing the iterated removal. The proof of Lemma 4 of [6] goes over verbatim to show that for large  $k$ , close points we have that their phases are close: That is, if

$$d_k(p, x) \leq k^{1/10}, \quad p, x \in \bar{L}$$

then

$$\left| \arg \left( \frac{\sigma_{k,p}(x)}{\tau_k(x)} \right) - \arg \left( \frac{\sigma_{k,p}(p)}{\tau_k(p)} \right) \right| \leq \frac{\pi}{4}.$$

We imitate the argument of Theorem 3.3 in Pascaleff-Tonkonog [44]. To match with the notation in [44] do the following renaming: call  $\sigma_{k,L_i}$  as  $s_i$  and  $\sigma_{k,L_\cap}$  as  $s_\cap$ . We will drop  $k$  from the subscript in the notation since the following bounds for some fixed parameters  $c_l, c_M, r_l, r_M$  do not depend on  $k$ :

- $|s_\cap(x)| \geq c_l \exp(-r_l(d_k^2(x, L_\cap) + d_k(x, L_\cap)))$  for  $x \in \bar{L}$ , with  $d_k(x, L_\cap) < k^{\frac{1}{30}}$  from Lemma 4.6,

- $c_l \leq |s_i| < 1$  for all  $i$ ,
- $|s_i(x)| \leq c_M \exp(-r_M(d_k^2(x, L)))$ .

Define  $c_\theta \in \mathbb{R}$  such that for  $p \in L_i, q \in L_j$  with  $i \neq j$  which are at a distance  $d$  from  $L_\cap$ , then  $d_k(p, q) \geq c_\theta d$ . Such a constant  $c_\theta$  exists because the angles between  $L_i, L_j$  have a lower bound, by assumption. Let  $\delta > 0$  be large enough so that

$$c_M \exp(-r_M c_\theta^2 \delta^2) < \frac{c_l}{10m^2}.$$

This choice of  $\delta$  does not depend on  $k$ . Set

$$(24) \quad c := \frac{c_l \exp(-r_l(\delta^2 + \delta))}{10m^2}.$$

and

$$s := s_\cap + c \sum s_i : \bar{X} \rightarrow \hat{X}^k.$$

For  $k$  large, we have  $\delta < k^{\frac{1}{30}}$ . Thus, in a  $\delta$  neighborhood of  $L_\cap$ , from Lemma (4.6), we have that

$$|s_\cap(x)| > c_l \exp(-r_l(\delta^2 + \delta)) \quad \forall x \in \bar{L}$$

on  $L_i$ . By (24) the sections

$$|cs_j| < \frac{1}{10m^2} c_l \exp(-r_l(\delta^2 + \delta)).$$

Thus, their sum is bounded above in a  $\delta$  neighborhood of  $L_\cap$  as follows:

$$\left| \sum_{j=1}^m cs_j \right| \leq m \frac{c_l \exp(-r_l(\delta^2 + \delta))}{10m^2} = \frac{c_l \exp(-r_l(\delta^2 + \delta))}{10m}.$$

Thus we have in a  $\delta$  neighborhood of  $L_\cap$ ,

$$|s| \geq |s_\cap| - \left| \sum_{j=1}^m cs_j \right| \geq \left( 1 - \frac{1}{10m} \right) c_l \exp(-r_l(\delta^2 + \delta)).$$

When  $x \in L_i$  and outside an open neighborhood of  $\delta$  distance from  $L_\cap$  we have that  $s_\cap + cs_i$  is bounded below by  $cc_l$  on  $L_i$  whereas we have an upper bound

$$|cs_j| < cc_M \exp(-r_M c_\theta \delta^2) < \frac{cc_l}{10m^2}.$$

Thus, after adding the  $m-1$  sections  $cs_j$ , for  $j \neq i$ , to  $s_\cap + cs_i$  we have the following lower bound for any point on  $L_i$ :

$$\left| s_\cap + cs_i + \sum_{j \neq i} cs_j \right| \geq |s_\cap + cs_i| - \sum_{j \neq i} |cs_j| \geq cc_l - cc_l \frac{m-1}{10m^2} > \frac{cc_l}{10m^2}.$$

We get a global lower bound of  $s$  on any point in  $\bar{L}$ :

$$|s(x)| \geq \min \left\{ \frac{cc_l}{10m^2}, c_l \exp(-r_l(\delta^2 + \delta)) \right\}.$$

□

**Corollary 4.7.** *If  $L$  is a rational Lagrangian cobordism then there exists a Donaldson hypersurface  $D$  of arbitrarily large degree in  $\overline{X}$  with the property that  $\overline{L}$  is exact in the complement of  $D$ . In particular, any holomorphic disk in  $\overline{X}$  bounding  $\overline{L}$  with positive area intersects  $D$  in at least one (interior) point.*

*Proof.* Since  $\overline{L}$  is a branched Lagrangian in  $\overline{X}$ , the claim follows from Lemma 4.5 and the perturbation technique described in Donaldson [23] and Auroux-Gayet-Mohsen [6].  $\square$

**4.2. Fredholm theory.** The moduli space of holomorphic disks bounding a Lagrangian cobordism is cut out locally by a Fredholm map. Note that the domains of our maps are treed disks with punctures, we need to consider linearizing PDE's over domains with cylindrical ends and over intervals. The Fredholm theory for treed disks without punctures can be found in [14, Section 4.3]. The Fredholm theory for maps with cylindrical ends can be found in [48, Lecture 4]. Our following construction is a combination of these two approaches.

We state our results in the context of a Lagrangian cobordism in a symplectic cobordism as in Section 2.2. Recall that  $(\overline{X}, \omega)$  is a closed symplectic manifold,  $Y_{\pm}$  are two disjoint symplectic hypersurfaces in  $\overline{X}$  and  $X = \overline{X} - (Y_- \sqcup Y_+)$ . Moreover  $\overline{L}$  is a Lagrangian in  $\overline{X}$  such that  $\Pi_{\pm} = \overline{L} \cap Y_{\pm}$  are Lagrangians in  $Y_{\pm}$ . Then  $L = \overline{L} - \Pi_{\pm}$  is a Lagrangian cobordism in  $X$  between  $\Lambda_+$  and  $\Lambda_-$ .

Our Sobolev maps are perturbations of a class of model maps defined as follows. Let  $S$  be a punctured disk with ends  $\mathcal{E}(S)$ .

**Definition 4.8.** A *model map* is a smooth map  $u : (S, \partial S) \rightarrow (X, L)$  such that, with coordinates given by some collection of cylindrical/strip-like ends  $\kappa_e : \pm(0, \infty) \times [0, 1] \rightarrow S$  or  $\kappa_e : \mathbb{R} \times S^1 \rightarrow S$ , the map  $u$  is linear in the sense that

$$(25) \quad u(\kappa_e(s, t)) = (\mu s, \gamma(\mu t)) \in \mathbb{R} \times Z, \quad \forall (s, t) \in \pm(0, \infty) \times [0, 1]$$

where  $\gamma$  is a Reeb chord with action  $\mu$  or similarly in the case that  $\gamma$  is a Reeb orbit. The set of model maps is defined as

$$\text{Map}_0(S, X, L) = \{u : (S, \partial S) \rightarrow (X, L) \mid (25) \text{ holds for all } e \in \mathcal{E}(S)\}.$$

Our Sobolev completion is defined as a space of maps that are obtained by geodesic exponentiation of a Sobolev-class section of the pull-back of the tangent bundle. To define the class of sections, Let  $\beta$  be a *good bump function* on  $\mathbb{R}$  such that

- (a)  $\beta$  is non-decreasing;
- (b)  $\beta(s) = 1$  when  $s \geq 1$ ;
- (c)  $\beta(s) = 0$  when  $s \leq 0$ .

For any *Sobolev decay constant*  $\delta > 0$  we define a *Sobolev weight function*  $\kappa^{\delta}$  on  $S$  such that

- (a)  $\kappa^{\delta} = 0$  outside cylindrical/strip-like ends;
- (b) on each cylindrical/strip-like end, there is a constant  $M > 0$  with  $\kappa^{\delta}(s, t) = \delta\beta(|s| - M)$ .

Pick a Riemannian metric  $g$  on  $X$  which is cylindrical near infinity and a metric connection  $\nabla$ . We further assume that  $L$  is totally geodesic with respect to  $g$ . Equip  $S$  with a metric which is of standard form on the cylindrical and strip-like ends.

**Definition 4.9.** For Sobolev constants  $k, p$  with  $kp > 2$  and a model map  $u \in \text{Map}_0(S, X, L)$ , define an *enlarged weighted Sobolev space* limiting to Reeb chords  $\gamma_e, e \in \mathcal{E}(S)$  as follows: A *model section* of  $u^*TX$  is a section  $\xi : S \rightarrow u^*TX$  taking values in  $\partial u^*TL$  on the boundary so that on the complement of a compact set on the ends,  $\xi(s, t) = \mathcal{T}_t \xi(e)$  is given by parallel transport  $\mathcal{T}_t$  of an element of  $TL$  along the Reeb chord  $\gamma_e$  denoted  $\xi(e) \in \gamma_e(0)$ . For such a model section, define the Sobolev  $(k, p, \lambda)$ -norm as

$$\|\xi\|_{k,p,\lambda}^p := \sum_{e \in \mathcal{E}(S)} |\xi(e)|^p + \int_S (|\xi|^p + |\nabla \xi|^p + \cdots + |\nabla^k \xi|^p) e^{p\kappa_\lambda} ..$$

The completion of the space of model sections in this norm is denoted

$$\Omega^0(u^*TX, \partial u^*TL)'_{k,p,\lambda} := \{\xi \mid \|\xi\|_{k,p,\lambda} < +\infty, \xi|_{\partial S} \in TL\}.$$

Let  $\exp : TX \rightarrow X$  denote the geodesic exponentiation map defined using the metric  $g$  on  $X$ .

Define the *weighted Sobolev space of maps*

$$\text{Map}_{k,p,\lambda}(S, X, L) := \{\exp_u \xi \mid u \in \text{Map}_0(S, X, L), \xi \in W^{k,p,\lambda}(u^*TX, \partial u^*TL)\}.$$

This ends the Definition.

There is a similar discussion for Sobolev spaces of maps from broken intervals. Let  $I$  be a broken interval with subintervals  $I_j$  as in Definition 3.7. For each sub-interval, define a space of model maps from  $I_j$  to  $L$  that are constant near infinity. The space  $\text{Map}_{k,p}(I_j, L)$  is the space of maps obtained by geodesic exponentiation of a  $W^{k,p}$  section of  $u^*TL$  from a model map. The tangent space is denoted  $\Omega^0(T, (u|_{T^\circ})^*T^\circ L)_{k,p}$ , by which we mean  $W^{k,p}$  maps on each subinterval whose limits match at any breaking. By definition, such maps have limits along the infinite ends of any sub-interval  $I_j$ . Let  $\text{Map}_{k,p}(I, L)$  be the subset of the product of spaces  $\text{Map}_{k,p}(I_j, L)$  so that the limit along the positive end of the  $j$ -th interval is equal to that along the negative end of the  $j+1$ -th interval.

We now define similar Sobolev completion of maps from a treed disk. Let  $C = S \cup T$  be a treed disk, where  $S$  is the surface part and  $T$  is the edge part. Let  $\text{Map}_{k,p,\lambda}(C, X, L)$  denote the subset of the product of maps from the surface part and tree part that agree on the intersection:

$$\begin{aligned} \text{Map}_{k,p,\lambda}(C, X, L) &:= \{(u_S, u_T) : u_S|_{S \cap T} = u_T|_{S \cap T}\} \\ &\subset \text{Map}_{k,p,\lambda}(S, X, L) \times \text{Map}_{k,p}(T, L). \end{aligned}$$

We claim that the moduli space is cut out locally over the moduli space of domains by a Fredholm map. Recall from Section 3.3 that  $\mathcal{M}_\Gamma$  is the moduli space of treed

disks  $C = S \cup T$  of type  $\Gamma$ . Let  $\mathcal{M}_\Gamma^i, i = 1, \dots, m$  be an open cover of  $\mathcal{M}_\Gamma$  over which the universal treed disk

$$\mathcal{U}_\Gamma = \{(C, z \in C)\} \rightarrow \mathcal{M}_\Gamma$$

is trivial as a stratified bundle in the following sense: There exists a homeomorphism

$$\mathcal{U}_{\Gamma,i} := \mathcal{U}_\Gamma|_{\mathcal{M}_\Gamma^i} \rightarrow \mathcal{M}_\Gamma^i \times C$$

which induces a homeomorphism from any treed disk  $C' \in \mathcal{M}_\Gamma^i$  with  $C$  that is smooth on each surface component  $S_v \subset C$  and on each edge  $T_e \subset C$ . Thus we may view  $C'$  as the smooth space  $C$  equipped with an almost complex structure on the surface part  $j(C') \in \mathcal{J}(S)$  and metric  $g(C') \in \mathcal{G}(T)$  depending on  $C'$ . Define

$$\mathcal{B}_\Gamma^{i,k,p,\lambda} := \mathcal{M}_\Gamma^i \times \text{Map}_{k,p,\lambda}(C, X, L).$$

A chart near a pair  $(C', u')$  is given by the product of a chart for  $\mathcal{M}_\Gamma^i$  near  $C'$  and a neighborhood of 0 under the map constructed as follows: Suppose for simplicity that  $S$  has a single strip-like end and no cylindrical ends. Define a map

$$(26) \quad \begin{aligned} \Omega^0(S, (u|_S)^*TX)_{k,p,\lambda}' \oplus \Omega^0(T, (u|_{T^\circ})^*T^L)_{k,p} \oplus \Omega^0(T_{\circ,-}, (u|_{T_{\circ,-}})^*\mathcal{R}(\Lambda_-))_{k,p} \\ \oplus \Omega^0(T_{\circ,+}, (u|_{T_{\circ,+}})^*\mathcal{R}(\Lambda_+))_{k,p} \\ \oplus \Omega^0(T_D, (u|_{T_D})^*TX)_{k,p} \rightarrow \text{Map}_{k,p,\lambda}(C, X, L) \end{aligned}$$

by geodesic exponentiation, for some metric for which  $L$  is totally geodesic and we use these maps as charts to give a Banach manifold structure on  $\text{Map}_{k,p,\lambda}$ . The fiber of the vector bundle  $\mathcal{E}_\Gamma^{i,k-1,p,\lambda}$  over any map  $u$  is the bundle of one-forms of Sobolev differentiability class one less:

$$(27) \quad \begin{aligned} \mathcal{E}_{\Gamma,u}^{k-1,p,\lambda} := \Omega^{0,1}(S, (u|_S)^*TX)_{k-1,p,\lambda} \oplus \Omega^1(T_L, (u|_{T_L})^*TL)_{k-1,p,\lambda} \\ \oplus \Omega^1(T_{\circ,-}, (u|_{T_{\circ,-}})^*\mathcal{R}(\Lambda_-))_{k-1,p} \oplus \Omega^1(T_{\circ,+}, (u|_{T_{\circ,+}})^*\mathcal{R}(\Lambda_+))_{k-1,p} \\ \oplus \Omega^1(T_D, (u|_{T_D})^*TX)_{k-1,p} \end{aligned}$$

and in the case of broken edges, we mean the space of one-forms on the complement of the breakings. We remark that in general, the vector spaces  $\mathcal{E}_{\Gamma,u}^{k-1,p,\lambda}$  do not form a smooth Banach vector bundle over  $\mathcal{B}_\Gamma^{k,p,\lambda}$  for reasons explained in [13, Remark 4.2]: The transition maps between local trivializations identifying the curves with, say, fixed curves  $C_1, C_2$  involve identifications

$$\mathcal{M}_\Gamma^i \times \Omega^{0,1}(S_1, (u|_{S_1})^*TX)_{k-1,p,\lambda} \rightarrow \mathcal{M}_\Gamma^i \times \Omega^{0,1}(S_2, (u|_{S_2})^*TX)_{k-1,p,\lambda}$$

induced from varying diffeomorphisms  $S_1 \rightarrow S_2$  depending on the curve  $[C'] \in \mathcal{M}_\Gamma^i$ . Differentiating with respect to  $[C']$  induces a map to a Sobolev class of one lower, and so the transition maps are not differentiable. However, the union of  $\mathcal{E}_{\Gamma,u}^{k-1,p,\lambda}$  for maps  $u'$  in  $\mathcal{B}_\Gamma^{i,k,p,\lambda}$  is a smooth Banach vector bundle with local trivialization induced by parallel transport.

In order to achieve regularity, we assume that some of the edges map to a Donaldson hypersurface. Let  $D \subset \bar{X}$  be a Donaldson hypersurface so that  $\bar{L}$  is exact

in the complement of  $D$ . Recall from Corollary 4.7 that when  $\bar{L}$  is rational, we can construct such a hypersurface. For notational convenience, we set

$$\zeta_D = 0 \in \text{Vect}(D)$$

so that trajectories on  $D$  are constant maps. As in Cieliebak-Mohnke [17]: Each component of  $u^{-1}(D)$  contains an edge  $T_e, e \in \text{Edge}_D(\Gamma)$ .

The moduli spaces of treed disks are cut out locally by Fredholm sections: Define for each  $i$  a section

$$(28) \quad \mathcal{F}_\Gamma^i : \mathcal{B}_\Gamma^{i,k,p,\lambda} \rightarrow \mathcal{E}_\Gamma^{i,k-1,p,\lambda} \times X^{\#\text{Edge}_D(\Gamma)},$$

$$([C'], u') \mapsto \left( \bar{\partial}_{j(C')} u' | S, \frac{d}{dt} u' | T + \text{grad}(\zeta(u'), (\text{ev}_e(u))_{e \in \text{Edge}_D(\Gamma)}) \right)$$

where  $\zeta$  denotes either  $\zeta_\circ, \zeta_{\bullet, \Lambda_\pm}, \zeta_{\bullet, L}$  or  $\zeta_D$  depending on the type of edge. We drop the Sobolev constants  $k, p, \lambda$  to simplify notation. Embed

$$\underline{0} : \mathcal{B}_\Gamma^i \times D^{\#\text{Edge}_D(\Gamma)} \rightarrow \mathcal{E}_\Gamma^i \times X^{\#\text{Edge}_D(\Gamma)}$$

by the product of the zero section and a number of copies of the Donaldson hypersurface  $D$ . The moduli space of maps over  $\mathcal{M}_\Gamma^i$  is

$$\mathcal{M}_\Gamma^i(L) = (\mathcal{F}_\Gamma^i)^{-1}(\underline{0}),$$

that is, the inverse image of the zero section times a number of copies of the Donaldson hypersurface. The global moduli space  $\mathcal{M}_\Gamma(L)$  is obtained by patching together the local spaces  $\mathcal{M}_\Gamma^i(L)$ .

We introduce the following terminology for linearized operators and regularity.

**Definition 4.10.** The *linearized operator* at any solution  $u$  in some trivialization of the universal bundle is denoted

$$(29) \quad \tilde{D}_u : T_{(C,u)} \mathcal{B}_\Gamma^i \rightarrow \mathcal{E}_{\Gamma,u}^i \times (TX/TD)^{\#\text{Edge}_D(\Gamma)}, \quad \xi \mapsto \frac{\partial}{\partial \rho} \Big|_{\rho=0} \mathcal{F}_\Gamma^i(\exp_u(\rho \xi)).$$

The definition is independent of the choice of local trivialization of the universal bundle used.

A map  $u$  is *regular* if  $\tilde{D}_u$  is surjective, in which case  $\mathcal{M}_\Gamma(L)$  is a smooth manifold of dimension  $\text{Ind}(\tilde{D}_u)$  in an open neighborhood of  $u$ .

The *virtual dimension* of the moduli space  $\mathcal{M}_\Gamma(L)$  at  $u$  is the index of the linearized operator

$$\text{vdim}(\mathcal{M}_\Gamma(L)) := \text{Ind}(\tilde{D}_u)$$

which is the honest dimension if every element is regular. This ends the Definition.

The linearized operator is Fredholm, as for example shown in Wendl [48, Lecture 4]. The restriction of  $\tilde{D}_u$  to variations of the map  $\xi \in \Omega^0(S, u^*TX)$  is the linearized operator  $D_u$  studied in McDuff-Salamon [41] for holomorphic maps to closed symplectic manifolds.



The moduli space of treed maps is naturally a union over combinatorial type. For any integer  $d \geq 0$ , denote

$$(30) \quad \overline{\mathcal{M}}(L)_d = \bigcup_{\mathbb{F}} \mathcal{M}_{\mathbb{F}}(L)$$

the union of types with

$$(31) \quad \text{vdim } \mathcal{M}_{\mathbb{F}}(L) + \text{codim}(\mathcal{M}_{\mathbb{F}}) = d.$$

That is,  $\mathcal{M}_{\mathbb{F}}(L)$  is in the locus of expected dimension  $d$ , allowing for deformations of domain which change the combinatorial type.

The locus of regular maps has good smoothness properties. Denote by  $\mathcal{M}_{\mathbb{F}}^{\text{reg},i}(L) \subset \mathcal{M}_{\mathbb{F}}^i(L)$  the locus of regular maps. By the implicit function theorem for smooth maps between Banach manifolds, the locus of regular maps  $\mathcal{M}_{\mathbb{F}}^{\text{reg},i}(L)$  is a smooth manifold of expected dimension. The transition maps between the various moduli spaces  $\mathcal{M}_{\mathbb{F}}^{\text{reg},i}(L), \mathcal{M}_{\mathbb{F}}^{\text{reg},i'}(L)$  for two choices of  $i, i'$  so that  $\mathcal{M}_{\mathbb{F}}^i \cap \mathcal{M}_{\mathbb{F}}^{i'}$  are nonempty, are smooth as all derivatives exist by elliptic regularity. .

We define the contact differential and cobordism maps by counting rigid maps, defined as follows:

**Definition 4.11.** The locus of *rigid* configurations  $\mathcal{M}(L)_0$  is the locus of maps  $u : C \rightarrow X$  for which the expected dimension  $\text{Ind}(\tilde{D}_u)$  vanishes and  $\mathcal{M}_{\Gamma}$  represents a stratum of top dimension in the moduli space of domains.

The last condition is equivalent to requiring that the lengths  $\ell(e)$  of all edges  $e \in \text{Edge}(\Gamma)$  are finite and non-zero.

The case of maps in the symplectization is a special case of maps in the cobordism above. For a Legendrian  $\Lambda$  in  $Z$ , let  $\mathcal{M}(\Lambda)$  be the moduli space  $\mathcal{M}(\mathbb{R} \times \Lambda)$  up to the equivalence given by the translations in the  $\mathbb{R}$ -direction. Thus  $\mathcal{M}(\Lambda)_0$  is the locus in  $\mathcal{M}(\Lambda)$  that are rigid (after modding out by translation) and whose domains  $C = S \cup T$  are in a top-dimensional strata  $\mathcal{M}_{\Gamma} \subset \overline{\mathcal{M}}$  of treed disks.

**4.3. Transversality.** We consider the moduli space of treed disks with interior edges mapping to a Donaldson hypersurface as constructed in the previous section. The domains of stable treed buildings are automatically stable, except for trivial cylinders: For a non-constant sphere component  $u_v : S_v \rightarrow X$  there must be at least three edges  $T_e \subset C, e \in \text{Edge}_D(\Gamma)$  incident to  $S_v$ . Any holomorphic disk  $u : S \rightarrow X$  not contained in a fiber of the projection to  $Y$  meets  $D$  in at least one interior point, by the exactness of  $\Pi$  in  $X - D$ .

Domain-dependent perturbations are defined as maps from the universal curve. Suppose that  $\Gamma$  is a stable type of domain and denote by  $\mathcal{M}_{\Gamma}$  the moduli space of disks of type  $\Gamma$ . Over  $\mathcal{M}_{\Gamma}$  we have a universal curve  $\mathcal{U}_{\Gamma}$  which decomposes into tree and surface parts

$$\mathcal{U}_{\Gamma} = \{[C, z], z \in C\}, \quad \mathcal{U}_{\Gamma} = \mathcal{S}_{\Gamma} \cup \mathcal{T}_{\Gamma}$$

depending on whether the point  $z$  lies in the surface  $S$  or tree  $T$  part of  $C$ .

**Definition 4.12.** Let  $(X, L)$  be a decorated cobordism as above. A *domain-dependent perturbation* is a collection of maps

$$\begin{aligned} J_\Gamma &: \mathcal{S}_\Gamma &\rightarrow \mathcal{J}_{\text{cyl}}(X) \\ \zeta_{\bullet, \Gamma} &: \mathcal{T}_{\bullet, \Gamma} &\rightarrow \text{Vect}(L) \\ \zeta_{\circ, \pm, \Gamma} &: \mathcal{T}_{\circ, \pm, \Gamma} &\rightarrow \text{Vect}(\mathcal{R}(\Lambda_\pm)) \end{aligned}$$

consist of a domain-dependent cylindrical almost complex structure  $J_\Gamma$  and domain-dependent vector fields  $\mathcal{T}_{\bullet, \Gamma}$  and  $\mathcal{T}_{\circ, \Gamma}$  so that the corresponding maps

$$\begin{aligned} \mathcal{S}_\Gamma \times TX &\rightarrow TX \\ \mathcal{T}_{L, \Gamma} \times L &\rightarrow TL \\ \mathcal{T}_{\circ, \pm, \Gamma} \times \mathcal{R}(\Lambda_\pm) &\rightarrow T\mathcal{R}(\Lambda_\pm) \end{aligned}$$

are smooth,  $\zeta_\Gamma(z)$  is of the form in (b), (12) on each edge depending on the label of the edge. Later, we will assume that the perturbations vanish on an open subset containing the zeros of the vector fields and the intersections  $\mathcal{S}_\Gamma \cap \mathcal{T}_\Gamma \subset \mathcal{U}_\Gamma$ .

One may also allow Hamiltonian perturbations. These are not necessary for regularization of the moduli spaces in the case of Legendrians with embedded projection but is necessary for those with immersed Lagrangian projections. The use of Hamiltonian perturbations is also useful for our later localization computations. Let

$$\text{Vect}_h(X, U) \subset \text{Vect}(X)$$

denote the space of  $\mathbb{R}$ -invariant Hamiltonian vector fields vanishing on an open subset  $U$  (which later will be taken to include the Donaldson hypersurface used for regularization). A domain-dependent Hamiltonian perturbation is a one-form

$$H_\Gamma \in \Omega^1(\mathcal{S}_\Gamma, \text{Vect}_h(X, U)).$$

A *domain-dependent perturbation* is a tuple

$$P_\Gamma = (J_\Gamma, \zeta_{L, \Gamma}, \zeta_{\circ, -, \Gamma}, \zeta_{\circ, +, \Gamma}, H_\Gamma)$$

as above.

Domain-dependent perturbations help us obtain transversality of moduli spaces by breaking the symmetry of the auxiliary choices such as the almost complex structure and Morse functions. Although beneficial for obtaining transversality, it comes with a cost: We lose expected symmetry such as the divisor axiom for curve counts. To overcome this difficulty we use multi-valued domain dependent perturbation which allows us to restore part of the expected symmetry back while still preserving transversality.

**Definition 4.13.** A *multi-valued domain-dependent perturbation* is a formal sum of domain-dependent perturbations with real coefficients

$$(32) \quad \tilde{P}_\Gamma := \sum_{i=1}^{k_\Gamma} c_i P_{\Gamma, i}$$

where

$$\sum_{i=1}^{k_\Gamma} c_i = 1, \quad c_i > 0.$$

We call the terms  $P_{\Gamma,i}$  *sheets* of the multi-valued perturbation and the terms  $c_i$  are the *weights* of the corresponding sheets. Two multi-valued perturbations are considered equivalent if they are related by replacements of the form

$$(33) \quad c_i P_{\Gamma,i} + c_j P_{\Gamma,j} = (c_i + c_j) P_{\Gamma,i} = (c_i + c_j) P_{\Gamma,j}$$

whenever  $P_{\Gamma,i} = P_{\Gamma,j}$ . The number of sheets is the minimum of the number  $k_\Gamma$  over all representatives, that is, the number of distinct  $P_{\Gamma,j}$ .

Our perturbations of the almost complex structure are required to be small in the following sense.

**Definition 4.14.** Define a (discontinuous) function recording the number of interior edges meeting the surface component containing the given point:

$$n_D : \mathcal{S}_\Gamma \rightarrow \mathbb{Z}_{\geq 0}, \quad z \mapsto \max_{z \in S_v} \# \{e \in \text{Edge}_D(\Gamma), T_e \cap S_v \neq \emptyset\}.$$

Suppose that  $[D] = k[\omega_Y]$  has degree  $k$ . A domain-dependent almost complex structure  $J_\Gamma : \mathcal{S}_\Gamma \rightarrow \mathcal{J}(X)$  is *stabilizing* if it takes values in the set for which there are no holomorphic spheres contained in the Donaldson divisor, that is,

$$J_\Gamma(z) \in \mathcal{J}^{n_D/k}(X), \quad \forall z \in \mathcal{S}_\Gamma$$

with notation from Proposition 4.2.

The set of perturbations satisfying the stabilizing condition contains a non-empty open subset of domain-dependent almost complex structures, by definition.

We will require that the perturbations for each type agree with those already chosen on the boundary. The following morphisms of graphs  $\Gamma_2 \rightarrow \Gamma_1$  correspond to inclusions of  $\overline{\mathcal{M}}_{\Gamma_1}^{<E}(\Lambda)$  in the formal boundary of  $\overline{\mathcal{M}}_{\Gamma_2}^{<E}(\Lambda)$ .

**Definition 4.15.** An *elementary morphism* of domain types  $\Gamma_1 \rightarrow \Gamma_2$  is one of the following:

- (a) (Collapsing or breaking edges) An isomorphism  $\Gamma_1 \rightarrow \Gamma_2$  where  $\Gamma_2$  is obtained from  $\Gamma_1$  by removing an edge  $e$  from a set  $\text{Edge}_0(\Gamma)$  or  $\text{Edge}_\infty(\Gamma)$  and placing it in  $\text{Edge}(\Gamma)$ ; corresponding to a degeneration of  $C$  in which the length  $\ell(e)$  of  $e \in \text{Edge}(\Gamma)$  becomes infinite or zero.
- (b) (Cutting edges) A morphism  $\Gamma_1 \rightarrow \Gamma_2$  where  $\Gamma_2$  is obtained from  $\Gamma_1$  by identifying two vertices  $v_-, v_+$  joined by an edge  $e$  which is then collapsed, corresponding to a degeneration of  $C$  in which a component  $S_v \subset S$  develops a node, either in the interior or boundary depending on the collapsed edge. Thus  $S_v$  becomes the union of two components  $S_{v_1}, S_{v_2}$ .

We apply a Sard-Smale argument to show that the moduli space of expected dimension at most one is cut out transversally for generic perturbations. For this, it is convenient to adopt a space of perturbations introduced by Floer.

**Definition 4.16.** Let  $\underline{\epsilon} = (\epsilon_0, \epsilon_1, \epsilon_2, \dots)$  be a sequence of positive real numbers. For a smooth function  $f : \mathbb{R} \rightarrow \mathbb{R}$  define the  $\underline{\epsilon}$ -norm

$$\|f\|_{\underline{\epsilon}} = \sum_{j \geq 0} \epsilon_j \|f\|_{C^j}$$

denote the (possibly infinite) *Floer norm* of  $f$ , as in [48, Appendix B]. For sequences  $\underline{\epsilon}$  which converge to zero sufficiently fast, the space  $C^{\underline{\epsilon}}$  of functions with finite Floer norm contains bump functions supported in arbitrarily small neighborhoods of any point. Similar spaces are defined for functions on any manifolds, sections of vector bundles and so on. Let

$$\mathcal{P}_{\Gamma}^{\underline{\epsilon}} = \{P_{\Gamma} \in C^{\underline{\epsilon}}\}$$

denote the space of perturbations for type  $\Gamma$  of finite  $\underline{\epsilon}$ -norm. Let

$$\mathcal{B}_{k,p,\underline{\epsilon},\lambda}^{\text{univ}} = \mathcal{M}_{\Gamma} \times \text{Map}(C, X)_{k,p,\lambda} \times \mathcal{P}^{\underline{\epsilon}}$$

the space of triples  $(C, u, P)$  where  $P = (J, \zeta)$  is a perturbation, and the subscript  $k, p, \lambda$  indicates that the restriction of  $u$  to each component  $T_e, S_v$  is in the weighted Sobolev class  $W^{k,p,\lambda}$ . Denote by

$$\begin{aligned} \mathcal{E}^{\text{univ},i} &:= \mathcal{E}_{\Gamma,k-1,p,\underline{\epsilon},\lambda}^{\text{univ},i} := \bigcup_{(C,u,P) \in \mathcal{B}_{\Gamma,k,p,\underline{\epsilon},\lambda}^{\text{univ},i}} \Omega^{0,1}(C, (u|S)^*TX)_{k-1,p,\lambda} \\ &:= \bigcup_{(C,u,P) \in \mathcal{B}^{\text{univ}}} \Omega^{0,1}(S, u^*TX)_{k-1,p,\lambda} \oplus \Omega^1(T_L, (u|T_L)^*TL)_{k-1,p} \\ &\quad \oplus \Omega^1(T_{\circ,-}, (u|T_{\circ,-})^*T\mathcal{R}(\Lambda_-))_{k-1,p} \\ &\quad \oplus \Omega^1(T_{\circ,+}, (u|T_{\circ,+})^*T\mathcal{R}(\Lambda_+))_{k-1,p} \\ &\quad \oplus \Omega^1(T_D, (u|T_D)^*TX)_{k-1,p}. \end{aligned}$$

Define a section

$$(34) \quad \mathcal{F}_{\Gamma}^{\text{univ},i} : \mathcal{B}_{\Gamma}^{\text{univ},i} \rightarrow \mathcal{E}_{\Gamma}^{\text{univ},i} \times X^{\# \text{Edge}_D(\Gamma)}, \quad (C', u, P_{\Gamma}) \mapsto \left( \bar{\partial}_{j(C'), J_{\Gamma}, H_{\Gamma}} u_S, \left( \frac{d}{ds} + \zeta_{C', \Gamma, \cdot}(u_{T, \cdot}) \right), (\text{ev}_e(u))_{e \in \text{Edge}_D(\Gamma)} \right)$$

where  $\bar{\partial}_{j(C'), J_{\Gamma}}$  is the Cauchy-Riemann operator defined by the choice of curve  $C'$  in the domain, and  $\zeta_{C', \Gamma, \cdot}$  is the perturbed vector field determined by the choice of domain  $C'$  and  $\cdot$  ranges over types of edges. Embed

$$\underline{0} : \mathcal{B}_{\Gamma}^{\text{univ},i} \times D^{\# \text{Edge}_D(\Gamma)} \rightarrow \mathcal{E}_{\Gamma}^{\text{univ},i} \times X^{\# \text{Edge}_D(\Gamma)}$$

by the product of the zero section and a number of copies of the Donaldson hyper-surface  $D$ . This ends the Definition.

**Theorem 4.17.** *The section  $\mathcal{F}_{\Gamma}^{\text{univ},i}$  is transverse to the zero section  $\underline{0}$  at a map  $u : C \rightarrow X$  as long as there are no constant spheres  $S_v, v \in \text{Vert}(\Gamma)$  with more than one edge  $e \in \text{Edge}_D(\Gamma)$  attached.*

*Proof.* We treat transversality for constant, non-constant, and trivial strip components separately. Let  $\eta$  be an element of the cokernel  $\text{coker}(\tilde{D}_u)$  of the linearized operator of (29). We write the restriction of  $\eta$  to the sphere or disk components resp. edges as

$$\eta_v \in \Omega^{0,1}(S_v, u_v^*TX), v \in \text{Vert}(\Gamma)$$

and

$$\eta_e \in \Omega^1(T_e, u_e^*TL), e \in \text{Edge}_\bullet(\Gamma), \quad \eta_e \in \Omega^1(T_e, u_e^*T\Lambda), e \in \text{Edge}_\circ(\Gamma).$$

We wish to show that all of the components  $\eta_v$  and  $\eta_e$  vanish.

First we show that the one-form vanishes on components on which the map is non-constant. Suppose  $v \in \text{Vert}(\Gamma)$  with map  $u_v := u|_{S_v}$  mapping to  $X$  such that  $u_v$  is non-constant. Then  $D_{u_v}^* \eta_v = 0$  weakly and  $\eta_v$  is perpendicular to variations  $K_\Gamma \in T_{J_\Gamma} \mathcal{J}_\Gamma(X)$  which are domain-dependent. By [41, Lemma 3.2.2],  $\eta_v$  vanishes in an open neighborhood of any point  $z \in S_v$  on which  $du_v$  is non-zero. On the other hand, we may identify  $\eta_v$  with an element in the kernel  $\ker(\tilde{D}_{u_v}^*)$  of the adjoint, and so  $\eta_v$  vanishes by the unique continuation principle in [41, Section 2.3]. This holds even after the sections  $\xi$  are constrained to vanish on the intersections  $S \cap T$ , as in [41, Section 3.4].

Next, we show that the one-form vanishes on a component mapping to the neck on which the projection to the base is non-constant. Suppose  $v \in \text{Vert}(\Gamma)$  with map  $u_v := u|_{S_v}$  such that

$$\bar{u}_v := p \circ u_v : S_v \rightarrow Y$$

is non-constant. The restriction  $\eta_v$  is a weak solution to  $D_{u_v}^* \eta_v = 0$  and  $\eta_v$  is perpendicular to one-forms produced by variations  $K_\Gamma$  in  $J_\Gamma$ . Such variations satisfy  $J_\Gamma K_\Gamma = -K_\Gamma J_\Gamma$  and vanish on the first factor in the decomposition

$$u^*TX \cong u^* \ker(Dp) \oplus \bar{u}^*TY.$$

Suppose  $\eta_v(z) \neq 0$ . Choose a  $(J_\Gamma, j)$ -antilinear map from  $T_z S$  to  $T_{u(z)} X$  pairing non-trivially with  $\eta_v(z)$ , which we may take to be of the form  $K_\Gamma(z) d_z u j(z)$  for some  $K_\Gamma(z)$  since the second component of  $d_z u$  is non-vanishing. Using a cutoff function, we obtain a one-form  $K_\Gamma du j$  that pairs non-trivially with  $\eta$ , which is a contradiction. Thus  $\eta_v$  vanishes.

The case of components mapping to fibers in the neck piece, that is, to the images of Reeb chords or orbits, requires special consideration and is typically the hardest part of the transversality argument in symplectic field theory. Any disk  $u_v : S_v \rightarrow X$  contained in a fiber

$$p^{-1}(y) \cong \mathbb{CP}^1, y \in Y$$

is by assumption a disk bounding

$$L \cap p^{-1}(y) \cong e^{2\pi i \theta_1} \mathbb{R}P^1 \cup \dots \cup e^{2\pi i \theta_k} \mathbb{R}P^1.$$

The linearized operator  $D_{u_v}$  is a direct sum

$$D_{u_v} \cong D_{u_v}^\vee \oplus D_{u_v}^h$$

of a trivial horizontal part  $D_{u_v}^h \cong \bar{\partial}$  taking values in  $T_{\bar{u}_v} Y$  and a vertical part  $D_{u_v}^\vee$  taking values in  $\mathbb{C}$  which has boundary condition  $\mathbb{R}$ . We examine the kernel and the

cokernel separately. The operator  $D_{u_v}$  splits into a sum of the standard operator  $\bar{\partial}$  on the one-dimensional factor, c.f. Ekholm [24, Lemma 6.8]. The kernel of  $D_{u_v}^h \cong \bar{\partial}$  on  $S_v$  consists of bounded holomorphic functions  $\xi : S_v \rightarrow \mathbb{C}$  with real boundary values. Therefore  $\ker(D_{u_v}^h)$  is the one-dimensional real vector space of constant functions with values in the reals. On the other hand, the cokernel  $\text{coker}(D_{u_v}^h)$  may be identified with holomorphic 0, 1-forms  $\eta$  of Sobolev class  $-k, p^*, \lambda$  enforcing exponential decay (since constant functions near infinity lie in the domain). In the local coordinate on  $\bar{S}_v$  near each puncture such a holomorphic 0, 1-form is vanishing at the puncture. Since there are no such holomorphic 0, 1-forms  $\eta$  on a disk  $S_v$ , the cokernel of  $D_{u_v}^h$  vanishes. As in Oh [42], any rank one Riemann-Hilbert problem on  $\bar{S}$  has either vanishing kernel or vanishing cokernel.

Transversality of the gradient flow equation on both types of edges follows from the fundamental theorem of ordinary differential equations. For any edge  $e \in \text{Edge}_o(\Gamma)$  the cokernel of the operator  $D_{u_e}$  on zero-forms on  $T_e$  is identified, via Hodge star, with the space of solutions to the adjoint equation

$$D_{u_e}^* \xi_e = \nabla \xi_e + \nabla \text{grad}(\zeta(u_e)) ds$$

with vanishing at the endpoints; there are no such solutions. In the case without endpoints  $T_e \cong \mathbb{R}$  that may occur in a broken trajectory, transversality follows from the Morse-Smale assumption on the Morse function  $f_o$ . On the other hand,  $D_{u_e}$  has kernel isomorphic to  $T_{u_e(z)}L$  for any point  $z \in T_e$ , via evaluation.

It remains to show that the matching conditions are transversally cut out. Consider a tree  $\Gamma_0 \subset \Gamma$  such that for each vertex  $v \in \text{Vert}(\Gamma_0)$ , the component  $u_v := u|_{S_v}$  is vertical. The argument of the previous section shows that such configurations are transversally cut out if each  $S_v$  is a disk or sphere and  $\Gamma_0$  is a tree: Let  $u_0$  denote the restriction of  $u$  to  $S_0$  and  $\ker(D_{u_0})'$  the kernel of the linearized operator acting on sections without matching conditions enforced at the nodes of  $S_0$ . By the previous paragraph  $\ker(D_{u_0})'$  is a sum of factors of  $\mathbb{C} \oplus T_{p \circ u(C)}Y$  for each sphere component and  $\mathbb{R} \oplus T_{p \circ u(C)}\Pi$  for each disk component. Transversality is equivalent to the condition that the map

$$\ker(D_{u'})' \rightarrow \bigoplus_{e \in \text{Edge}_{\bullet, \text{fin}}(\Gamma)} T_{p \circ u(C)}Y \oplus \bigoplus_{e \in \text{Edge}_{o, \text{fin}}(\Gamma)} T_{p \circ u(C)}\Pi$$

is surjective. This follows from an induction (for each connected component of the subset of the domain on which the horizontal projection of the map is constant). For each such component, the induction starts with the nodes contained in terminal edges of  $\Gamma_0$ , so that the adjacent disk has a single node represented by an edge in  $\Gamma_0$ . Surjectivity of the linearized evaluation map for this component implies that the matching condition is cut out transversally at this edge; the inductive hypothesis then implies that the remaining matching conditions are transversally cut out.

Putting everything together, we obtain that the operator  $D'_u$  whose domain consists of sections without matching conditions is surjective, and the kernel of  $D'_{u_v}$  surjects onto any intersection point  $TL|_{S_v} \cap T$  or  $T\Pi|_{S_v} \cap T$ . Similarly the kernel of  $D'_{u_e}$  surjects onto any intersection point  $TL|_{T_e} \cap S$ . It follows that the matching conditions at the nodes are cut out transversally.  $\square$

Perturbations are constructed by an induction on the dimension of the source moduli space. Suppose that single-valued perturbations  $P_{\Gamma'}$  have been chosen for all types  $\Gamma' \prec \Gamma$ . A natural gluing procedure gives perturbations  $P_{\Gamma}$  in an open neighborhood  $U_{\Gamma}$  of the boundary of  $\overline{\mathcal{M}}_{\Gamma}$ . The gluing theorem for linearized operators (see, for example, Wehrheim-Woodward [47, Theorem 2.4.5] for the closed case) implies that for  $U_{\Gamma}$  sufficiently small, the perturbation  $P_{\Gamma}$  is regular over  $U_{\Gamma}$  if all the perturbations  $P_{\Gamma'}$  are regular. The case of multi-valued perturbations is the same, with the following caveat: It could be the boundary of  $\mathcal{M}_{\Gamma}$  is disconnected, so that there are for example two types  $\Gamma_1, \Gamma_2 \preceq \Gamma$  so that  $P_{\Gamma_1}$  and  $P_{\Gamma_2}$  have different number of sheets, say  $k_1$  and  $k_2$  respectively. In this case there is an obvious replication process which  $P_{\Gamma_1}, P_{\Gamma_2}$  are each replaced by multi-valued perturbations with  $k_1 k_2$  sheets, with each of the original sheets in  $P_{\Gamma_1}$  repeated  $k_2$  times, and similarly for  $P_{\Gamma_2}$ . The gluing construction then proceeds as before.

Similarly, given a type  $\Gamma$  obtained from  $\Gamma_1, \Gamma_2$  by gluing along an edge, any multivalued perturbations  $P_{\Gamma_1}, P_{\Gamma_2}$  for  $\Gamma_1, \Gamma_2$  with  $k_1, k_2$  sheets respectively induces a multivalued perturbation for type  $\Gamma$  with  $k_1 k_2$  sheets.

**Theorem 4.18.** *Suppose that regular perturbations  $P_{\Gamma'}$  have been chosen for all types  $\Gamma' \prec \Gamma$ . Then there exists a comeager set  $\mathcal{P}_{\Gamma}^{\text{reg}} \subset \mathcal{P}_{\Gamma}$  of perturbations  $P_{\Gamma}$  agreeing with the glued perturbations arising from  $P_{\Gamma'}$  in an open neighborhood  $U_{\Gamma}$  of the boundary of  $\overline{\mathcal{M}}_{\Gamma}$  such that any  $P_{\Gamma} \in \mathcal{P}_{\Gamma}^{\text{reg}}$  is regular.*

*Proof.* The statement of the Theorem is an application of Sard-Smale. By Theorem 4.17 the universal moduli space (locally over the moduli space) is transversally cut out. Consider the forgetful map

$$f_{\mathbb{F}}^i : \mathcal{M}_{\mathbb{F}, k, p, \epsilon, \lambda}^{\text{univ}, i}(L) \rightarrow \mathcal{P}^{\epsilon} \quad (u : C \rightarrow X, P_{\Gamma}) \mapsto P_{\Gamma}.$$

For map types  $\mathbb{F}$  of expected dimension at most one, the index of  $f_{\mathbb{F}}^i$  is at most one and by Sard-Smale the set of regular values of  $f_{\mathbb{F}}^i$  is comeager. Since the space of map types  $\mathbb{F}$  with underlying domain type  $\Gamma$  is countable, there exists a comeager set  $\mathcal{P}_{\Gamma}^{\text{reg}} \subset \mathcal{P}_{\Gamma}$  of perturbations  $P_{\Gamma}$  extending the given perturbations on the boundary so that  $\mathcal{M}_{\mathbb{F}, k, p, \epsilon, \lambda}^{\text{univ}, i}(L)$  is regular for every map type  $\mathbb{F}$  of expected dimension at most one.  $\square$

**Theorem 4.19.** *For any regular perturbation system  $\underline{P} = (P_{\Gamma})$ , for any map type  $\mathbb{F}$  decorating  $\Gamma$  of expected dimension at most one the moduli space  $\mathcal{M}_{\mathbb{F}}(L)$  is transversally cut out and for any pair  $\mathbb{F}_1 \prec \mathbb{F}_2$  with  $\mathbb{F}_1$  resp  $\mathbb{F}_2$  having expected dimension zero resp. one. there is a natural gluing map  $\mathcal{M}_{\mathbb{F}_1}(L) \times [0, \epsilon) \rightarrow \mathcal{M}_{\mathbb{F}_2}$ .*

The Theorem, for which we do not give a proof, is a consequence of gluing results similar to those in Ekholm-Etnyre-Sullivan [25, Chapter 10], who prove gluing theorems in the non-degenerate case, and Palmer-Woodward [43], which deals with a similar Morse-Bott case. Given a regular stratum-wise rigid treed holomorphic building  $u : C \rightarrow \mathbb{X}$  of type  $\Gamma_1$  representing a codimension one stratum  $\mathcal{M}_{\Gamma_1}$  in the boundary of a top-dimensional stratum  $\mathcal{M}_{\Gamma_2}$  there exists a unique family  $u_{\rho} : C_{\rho} \rightarrow X_{\rho}$  of treed holomorphic curves of type  $\Gamma_2$  converging to  $u$ . In the case of collapsing edges or making edges finite, each such sequence  $u_{\rho}$  is defined by

first constructing an *approximate solution*: One removes small balls around a node and uses cutoff functions and geodesic exponential to construct the approximate solutions while in the case of trajectories one removes a small ball around the intermediate critical point and patches together using cutoff function. On the other hand, the case of making zero-length edges positive length follows immediately from the implicit function theorem.

**4.4. Intersection multiplicities.** We develop a notion of fractional intersection multiplicity with the divisors at infinity, related to the sum of actions of Reeb chords. Fractional intersection multiplicities with the divisors at infinity are defined by integrating the Thom class. Let  $u : (S, \partial S) \rightarrow (\bar{X}, \bar{L})$  be a punctured surface bounding. The *intersection number* of  $u$  with  $Y_\pm$  is the pairing of  $[u]$  with  $[\text{Thom}_{Y_\pm}]$ . As usual, the intersection number may be computed as a sum of (now fractional) local intersection numbers, proportional to the sum of actions at the intersection points:

**Lemma 4.20.** *Suppose that  $u : (S, \partial S) \rightarrow (X, L)$  is a punctured disk asymptotic to a collection of Reeb orbits and chords  $\gamma_e, e \in \mathcal{E}(S)$ . The intersection number  $[\bar{u}] \cdot [Y_\pm]$  is equal to the sum of the actions*

$$[\bar{u}] \cdot [Y_\pm] = \sum_{e \in \mathcal{E}_\pm(S)} \theta_e$$

of the Reeb chords and orbits limiting to  $Y_\pm$ .

*Proof.* The proof is an argument using Stokes' formula. The cutoff-function  $\rho$  is equal to 1 near the zero section. Hence

$$\begin{aligned} \lim_{s \rightarrow \infty} \int_{S - \cup_{e \in \mathcal{E}_\pm(S)} \kappa_e^{-1}(\pm(s, \infty) \times Z_\pm)} d(\rho \alpha_\pm) &= \lim_{s \rightarrow \infty} \int_{\partial(S - \cup_{e \in \mathcal{E}_\pm(S)} \kappa_e^{-1}(\pm(s, \infty) \times Z_\pm))} \alpha_\pm \\ &= \sum_{e \in \mathcal{E}_\pm(S)} \theta_e \end{aligned}$$

as claimed.  $\square$

**4.5. Compactness.** We will need a version of Gromov compactness for buildings in cobordisms with Lagrangian boundary conditions, adapted to the case of domain-dependent perturbations. In order for compactness to hold we must assume that the perturbations satisfy coherence properties.

We first note the following construction. Given a type  $\Gamma$  denote by  $\Gamma_\circ$  the type obtained by collapsing all sphere components  $S_v \subset S$  to interior leaves, and for any vertex  $v \in \text{Vert}(\Gamma)$  let

$$\Gamma(v) = (\{v\}, \{e \in \text{Edge}(\Gamma) | e \ni v\})$$

denote the union of edges meeting  $v$ . Let

$$\mathcal{U}_{\Gamma, v} = \{(C, z) \in \mathcal{U}_\Gamma | z \in S_v\}$$

denote the union of points lying on the components  $S_v \subset S$ .



**Definition 4.21** (Multivalued pull-back). Let  $\Gamma$  be a combinatorial type of treed disc which has an edge  $e$  of infinite length  $\ell(e) = \infty$ , and thus obtained from gluing two types  $\Gamma_1$  and  $\Gamma_2$  along edges  $e_1, e_2$  respectively. Let

$$\tilde{P}_{\Gamma_i} = \sum_{j=1}^{k_i} c_{k,i} P_{\Gamma_i, k}$$

be multivalued perturbations for  $i \in \{1, 2\}$ . The multivalued pull-back,  $\tilde{P}_\Gamma$ , of  $\tilde{P}_{\Gamma_1}$  and  $\tilde{P}_{\Gamma_2}$  is defined as

$$\tilde{P}_\Gamma = \sum_{j_1=1}^{k_1} \sum_{j_2=1}^{k_2} c_{j_1,1} c_{j_2,2} P_{\Gamma, j_1, j_2},$$

where  $P_{\Gamma, j_1, j_2}$  is defined as the pull-back of  $P_{\Gamma_1, j_1}$  and  $P_{\Gamma_2, j_2}$  under the isomorphism

$$\bar{\mathcal{U}}_\Gamma \rightarrow \pi_1^* \bar{\mathcal{U}}_{\Gamma_1} \times \pi_2^* \bar{\mathcal{U}}_{\Gamma_2}.$$

Here

$$\pi_b : \bar{\mathcal{M}}_\Gamma \rightarrow \bar{\mathcal{M}}_{\Gamma_b}, b \in \{1, 2\}$$

are the natural projections.

**Definition 4.22.** A collection of multi-valued perturbations

$$\underline{P} = (\tilde{P}_\Gamma \in \mathcal{P}_\Gamma)$$

for all types of domain  $\Gamma$  is *coherent* if the following holds:

- (a) (Collapsing Edges) Whenever  $\Gamma_1 \rightarrow \Gamma_2$ , so that  $\Gamma_2$  is obtained from  $\Gamma_1$  by collapsing edges, the number of sheets of  $\tilde{P}_{\Gamma_1}$  is equal to the number of sheets of  $\tilde{P}_{\Gamma_2}$  and each sheet  $P_{\Gamma_2, i}$  restricts to  $P_{\Gamma_1, i}$  on  $\mathcal{U}_{\Gamma_2}$  and their corresponding weights match  $c_i^{\Gamma_1} = c_i^{\Gamma_2}$ . That is,

$$J_{\Gamma_2, i} : \mathcal{S}_{\Gamma_2} \rightarrow \mathcal{J}(X)$$

restricts to  $J_{\Gamma_1, i}$  on  $\mathcal{S}_{\Gamma_1} \subset \bar{\mathcal{S}}_{\Gamma_2}$ , the vector field

$$\zeta_{\Gamma_2, \bullet, i} : \mathcal{T}_{\Gamma_2, \bullet, i} \rightarrow \text{Vect}(X)$$

restricts to  $\zeta_{\Gamma_1, \bullet, i}$  on  $\mathcal{T}_{\Gamma_1, \bullet, i}$  and similarly for the vector fields  $\zeta_{\Gamma_2, \circ, i}, \zeta_{\Gamma_1, \circ, i}$ .

- (b) (Breaking Edges) Whenever  $\Gamma$  has an edge of infinite length, and so obtained from gluing  $\Gamma_1$  and  $\Gamma_2$ , then  $\tilde{P}_\Gamma$  restricts to the multivalued pull-backs of  $\tilde{P}_{\Gamma_1}$  and  $\tilde{P}_{\Gamma_2}$  under the isomorphism

$$\bar{\mathcal{U}}_\Gamma \rightarrow \pi_1^* \bar{\mathcal{U}}_{\Gamma_1} \times \pi_2^* \bar{\mathcal{U}}_{\Gamma_2}$$

where  $\pi_b : \bar{\mathcal{M}}_\Gamma \rightarrow \bar{\mathcal{M}}_{\Gamma_b}, b \in \{1, 2\}$  are the natural projections.

- (c) (Locality) Let  $C_\circ \subset C$  be the subset consisting only of disk components and boundary edges. We have a natural map

$$f_{\Gamma, v} : \mathcal{U}_{\Gamma, v} \rightarrow \mathcal{U}_{\Gamma(v)} \times \mathcal{M}_{C_\circ}$$

which sends the pair  $(C, z \in S_v)$  to the pair  $((S_v, z \in S_v), C_\circ)$ . We require that the restriction of the perturbation  $J_\Gamma$  to  $S_v$  is pulled back under  $f_{\Gamma, v}$ . In particular, if  $\Gamma'$  is another component obtained from  $\Gamma$  by removing an

interior leaf  $T_e$  on some sphere component  $v$  then  $P_\Gamma$  induces a perturbation datum  $P_{\Gamma'}$  for the type  $\Gamma'$  by taking  $J_{\Gamma'}$  to be constant on  $S'_v$  and equal to  $J_\Gamma$  on the other components. Any  $P_\Gamma$  holomorphic map from a curve  $C$  of type  $\Gamma$  to  $X$  induces a  $P_{\Gamma'}$ -holomorphic map from the curve  $C'$  to  $X$ , where  $C'$  is obtained from  $C$  by forgetting  $T_e$ .

We have the following version of sequential Gromov compactness.

**Theorem 4.23.** *Any sequence of stable buildings  $u_\nu : C_\nu \rightarrow \mathbb{X}$  with boundary in  $\mathbb{R} \times \Lambda$  with bounded area  $A(u_\nu)$  and bounded number of leaves  $n(u_\nu)$  has a subsequence Gromov-converging to a stable building  $u : C \rightarrow \mathbb{X}$ .*

*Proof.* The limits of the surface parts and tree parts of the sequence may be constructed separately. For convergence on the curve parts, see Chanda [10] or, for the stable Hamiltonian case, Venugopalan-Woodward [46] (the latter without Lagrangian boundary condition). For convergence on the tree parts, let  $u_\nu : C_\nu \rightarrow \mathbb{X}$  be a sequence as in the statement of the Theorem. Each edge  $T_e, e \in \text{Edge}(\Gamma)$  in a level  $C_i$  mapping to  $\mathbb{R} \times \Lambda$  gives rise to a sequence of trajectories by restriction

$$u_{e,\nu} : [-T_\nu, T_\nu] \rightarrow \mathbb{R} \times \Lambda.$$

By compactness of gradient trajectories up to breaking, the projections

$$\bar{u}_{e,\nu} := p \circ u_\nu : [0, T_\nu] \rightarrow \Pi$$

converge after passing to a subsequence to a (possibly broken) trajectory. That is, that the exist sequences  $t_\nu$  so that the trajectories  $\bar{u}_{e,\nu}(t + t_\nu)$  converge in all derivatives on compact sets to some limit

$$\bar{u}_{e,i} : [-T_{e,i}, T_{e,i}] \rightarrow \Lambda$$

which is a trajectory of  $\text{grad}(f_\circ)$ , for some  $T_{e,i} \in [0, \infty]$ ; see for example Audin-Damian [5]. Choose a sequence  $T_\nu \in \mathbb{R}$  so that the point  $T_\nu u_{e,\nu}(0)$  (that is, the translation of  $u_{e,\nu}(0)$  in the  $\mathbb{R}$ -direction by  $T_\nu$ ) converges to some limiting point  $u_e(0)$ ; such a sequence  $T_\nu$  exists since  $\Lambda$  is compact. By the fundamental theorem of ordinary differential equations, the sequence  $u_{e,\nu}$  converges in all derivatives on compact sets to a trajectory of  $\zeta$ . Since the projections  $\bar{u}_{e,i}$  form a broken trajectory, the lifts  $u_{e,i}$  form a broken trajectory.  $\square$

We analyze the boundary of the one-dimensional moduli spaces  $\overline{\mathcal{M}}(\Lambda)_1$  resp.  $\overline{\mathcal{M}}(L)_1$  from (30), assuming that every map is regular.

**Definition 4.24.** (a) Strata  $\mathcal{M}_\Gamma(\Lambda)$  of configurations  $u : C \rightarrow \mathbb{X}$  with a single edge  $T_e \subset C$  of length zero and of expected dimension  $\text{vdim } \mathcal{M}_\Gamma(\Lambda) = 0$  will be called *fake boundary components* of  $\overline{\mathcal{M}}(\Lambda)_1$ . If  $\mathcal{M}_\Gamma(\Lambda)$  is regular then there are exactly two one-dimensional strata  $\mathcal{M}_{\Gamma'}(\Lambda), \mathcal{M}_{\Gamma''}(\Lambda)$  containing  $\mathcal{M}_\Gamma(\Lambda)$  in their closure. These strata consist of configurations  $u : C \rightarrow \mathbb{X}$  with an edge  $e$  of positive length  $\ell(e)$ , resp. the two adjacent disk components  $S_{v_-}, S_{v_+}$  have been glued to form a single disk component.

- (b) Strata with an edge of infinite length  $\ell(e) = \infty$  and of expected dimension  $\text{vdim } \mathcal{M}_{\mathbb{F}}(\Lambda) = 0$  are *true boundary components* of  $\overline{\mathcal{M}}(\Lambda)_1$ . There are two types of such edges  $e$ , depending on what type of edge has acquired infinite length:

- (i) edges  $e \in \text{Edge}_o(\Gamma)$ , which if length  $\ell(e)$  zero corresponds to nodes  $w \in \partial S$  separating components  $S_{v_+}, S_{v_-}$  and
- (ii) edges  $e \in \text{Edge}_o(\Gamma)$  joining different levels via Reeb chords  $\gamma \in \mathcal{R}(\Lambda)$ .

The definitions of fake and true boundary strata of  $\overline{\mathcal{M}}(L)_1$  are similar.

We have the following description of the boundary strata of the one-dimensional moduli spaces of buildings in cobordisms, stated in two separate theorems:

**Theorem 4.25.** *Suppose that  $\overline{P} = (P_{\Gamma})$  is a regular, coherent collection of perturbation data for a Legendrian  $\Lambda$ . The boundary of the one-dimensional stratum  $\overline{\mathcal{M}}(\Lambda)_1$  is a union of strata  $\mathcal{M}_{\mathbb{F}}(\Lambda)$  where  $\mathbb{F} = (\mathbb{F}_1, \mathbb{F}_2)$  is a type of building  $u : C \rightarrow \mathbb{R} \times Z$  with exactly two levels  $u_1 : C_1 \rightarrow \mathbb{R} \times Z, u_2 : C_2 \rightarrow \mathbb{R} \times Z$ .*

**Theorem 4.26.** *Suppose that  $\overline{P} = (P_{\Gamma})$  is a regular, coherent collection of perturbation data for a tame Lagrangian cobordism  $L$  from  $\Lambda_-$  to  $\Lambda_+$ . The boundary of  $\overline{\mathcal{M}}_{\mathbb{F}}(L)_1$  is a union of strata  $\mathcal{M}_{\mathbb{F}}(L)$  where  $\mathbb{F} = (\mathbb{F}_1, \mathbb{F}_2)$  a type corresponding to a building that is either*

- (a) *a treed building  $u : C \rightarrow X$  with exactly two levels  $u_1 : C_1 \rightarrow X, u_2 : C_2 \rightarrow \mathbb{R} \times Z_{\pm}$ , one of which maps to  $X$  and the other to  $\mathbb{R} \times Z_{\pm}$ , or*
- (b) *a building  $u : C \rightarrow X$  with a single level mapping to  $X$  consisting of two components  $u_- : C_- \rightarrow X, u_+ : C_+ \rightarrow X$  glued together at an infinite length trajectory  $u_e : T_e \rightarrow X$  at a critical point of the Morse function  $f_{\bullet}$ .*

*Proof of Theorems 4.25 and 4.26.* We prove the claim for the moduli space  $\overline{\mathcal{M}}(L)_1$  of buildings in a cobordism  $L$  from  $\Lambda_-$  to  $\Lambda_+$ ; the case of buildings in  $\mathbb{R} \times \Lambda$  is easier.

We start with ruling out the formation of interior nodes. There are two possibilities, depending on whether the node is in a sub-level, and so maps to the interior of a cobordism, or separates levels, in which case the node maps to a Reeb orbit. In the first case, let  $u : C \rightarrow \mathbb{X}$  be a building of type  $\mathbb{F}$  occurring in the boundary of  $\overline{\mathcal{M}}(\Lambda)_1$ . The surface part of the domain  $S$  consists of two components  $S_{v_1}, S_{v_2}$  with intersection  $w \in S_{v_1} \cap S_{v_2}$  lying in the interior of both components and  $S_{v_1}, S_{v_2}$  in the same level is a codimension two phenomenon in the space of domains and does not occur in zero and one-dimensional moduli spaces cut out transversally.

In the second case, the absence of nodes forming at Reeb orbits is a feature of our tameness assumptions in Definition 3.19: Suppose  $S_{v_1}, S_{v_2}$  are connected by a node  $w$  mapping to a Reeb orbit; that is,  $S_{v_1}$  and  $S_{v_2}$  each have a cylindrical end corresponding to an edge  $e \in \text{Edge}_o(\Gamma)$ . The disk components  $S_{v_1}, S_{v_2}$  are parts of different levels in the building. Let  $\mathbb{F} = \mathbb{F}_1 \cup \mathbb{F}_2$  denote the decomposition of  $\mathbb{F}$  into subgraphs obtained by dividing at  $e$ . Since the subgraph  $\Gamma_o$  of disk components is connected, either  $\Gamma_1$  or  $\Gamma_2$  consists entirely of spherical vertices  $v \in \text{Vert}_{\bullet}(\Gamma)$ . Suppose  $C = C_1 \cup C_2$  is the corresponding decomposition of domains, so that  $C_2$  is a treed sphere and let  $u_2 = u|_{C_2}$ . Since the combinatorial type is a tree, we may assume that  $C_2$  has incoming punctures but no outgoing punctures or vice versa,

and in particular,  $u_2$  is not a cover of a trivial cylinder. If  $u_2$  is a level mapping to  $X$  with a single incoming puncture and no outgoing punctures then as in 3.25, with  $e(\bullet)$  the number of interior punctures,

$$\begin{aligned} \dim \mathcal{M}_{\Gamma_2}(L) &= \dim(X) + 2c_1(\overline{X}) \cdot [u_2] + 2e(\bullet) - 2([Y_-] \cdot [u_2] - 1) - 6 \\ &\geq \dim(X) + 2(1 + \lambda_-)[\overline{\omega}] \cdot [u_2] - 2 > \dim(X). \end{aligned}$$

Therefore, the moduli space cannot be made rigid by adding constraints at the puncture. On the other hand, if  $u_2$  is a level with a single outgoing puncture and no incoming punctures then from (P3) we have,

$$0 > -[Y_+] \cdot [u_2] = \lambda_+[\omega] \cdot [u_2],$$

which is a contradiction. Similarly, if  $u_2$  is a level mapping to  $\mathbb{R} \times Z_+$  with one incoming puncture and no outgoing punctures then using  $\dim(X) = \dim(Z) + 1$

$$\begin{aligned} \dim \mathcal{M}_{\Gamma_2}(\Lambda) &= \dim(X) + 2c_1(\mathbb{P}(N_+ \oplus \mathbb{C})) \cdot [u_2] + 2e(\bullet) - 2([Y_+] \cdot [u_2] - 1) - 6 - 1 \\ &\geq \dim(X) + 2c_1(\overline{Y_+}) \cdot [\overline{u}_2] - 5 \geq \dim(X) - 1 \geq \dim(Y_+) + 1. \end{aligned}$$

The last inequality uses the fact that  $\overline{Y}$  has minimal Chern number at least two, by the monotonicity assumption in (3.19). Thus, the moduli space cannot be rigid by the addition of a constraint. The possibility that  $u_2$  has one outgoing puncture and no incoming punctures violates the angle equality in Lemma 4.20.

We now describe the boundary configurations for the one-dimensional moduli spaces. Let  $u : C \rightarrow \mathbb{X}$  be a configuration in the boundary of a one-dimensional component of  $\mathcal{M}(\mathcal{L})$ . By the previous paragraphs,  $C$  has a single boundary edge  $T_e$  with zero or infinite length. In the case of zero length,  $u$  is contained in a fake boundary component in the sense of Definition 4.24, and is not a topological boundary point. Thus the length of  $T_e$  is infinite, and either  $T_e$  separates two levels  $C_i, C_{i+1}$  or two components  $C', C''$  of the same level  $C_i$ . These are the possibilities listed in the statement of the Theorem.  $\square$

**4.6. Orientations.** Orientations of the moduli spaces in the case that the Legendrians resp. Lagrangians are equipped with relative spin structures are constructed by Ekholm-Etnyre-Sullivan [26], based on early work of Fukaya-Oh-Ohta-Ono [33]. See also Wehrheim-Woodward [47]. A *spin structure* on an oriented manifold  $L$  of dimension  $n \geq 3$  is an equivalence class of pairs consisting of a principal  $\text{Spin}(n)$ -bundle  $\text{Spin}(L) \rightarrow L$  and an isomorphism of the associated vector bundle

$$\text{Spin}(L) \times_{\text{Spin}(n)} \mathbb{R}^n \rightarrow TL$$

for the standard representation  $\mathbb{R}^n$  with the tangent bundle  $L$ .

Spin structures are used to distinguish trivializations of the pull-back of the tangent bundle to any curve. Since  $SO(n)$  is connected, the pull-back  $\gamma^*TL$  of  $TL$  to any curve  $\gamma : S^1 \rightarrow L$  is trivializable in two non-homotopic ways for  $n > 2$  related by a loop  $\epsilon : S^1 \rightarrow SO(n)$  with non-trivial homotopy class. Since  $\text{Spin}(n)$  is simply




FIGURE 8. Three types of boundary configurations (a) two levels with one incoming puncture on each component (b) one level with a broken trajectory (c) two levels with zero incoming puncture on some component

connected for  $n \geq 3$ , a spin structure gives in particular a choice of such trivialization up to homotopy. For  $n = 2$  a spin structure gives a homotopy class of such trivializations after stabilization.

In our study of fillings we will be particularly interested in spin structures on the boundary of a filling that extend over the interior. Let  $L$  be a manifold with boundary. Any spin structure on  $L$  induces a spin structure on the boundary  $\partial L$  using the trivialization  $TL_{\partial L} \cong T\partial L \oplus \mathbb{R}$ . For example, consider the unit circle  $\partial L = S^1$  in the unit disk  $L = B^2$ . On the one hand  $S^1$  has a unique *trivial spin structure* corresponding to the unique trivialization of the tangent bundle. On the other hand, the unique spin structure on  $B^2$  induces a *non-trivial spin structure* on the boundary  $S^1$ , since the two corresponding (stable) trivializations are related by a map  $S^1 \rightarrow SO(2)$  which generates  $\pi_1(SO(2))$ . A *relative spin structure* is defined similarly, but allowing the transition maps for  $TL$  to lift to a collection of transition maps taking values in  $\text{Spin}(n)$  that satisfy the cocycle condition up to the pull-back of a Čech two-chain on  $M$  with values in  $\mathbb{Z}_2$ .

Orientations on the moduli spaces are constructed by the following deformation argument. Each tangent space at a regular element  $u \in \mathcal{M}(L)$  is isomorphic to the kernel  $\tilde{D}_u$  of the linearized operator from (29):

$$T_u \mathcal{M}(L) := \ker(\tilde{D}_u).$$

The determinant line of  $\tilde{D}_u$  is denoted

$$\mathbb{D}_u = \Lambda^{\text{top}}(\ker(\tilde{D}_u)) \cong \Lambda^{\text{top}}(\ker(\tilde{D}_u)) \otimes \Lambda^{\text{top}}(\text{coker}(\tilde{D}_u)^*).$$

An orientation of  $\mathcal{M}(L)$  is a non-zero element  $o_u$  of the determinant line, modulo multiplication by a positive scalar.

Any path of Fredholm operators induces an identification of determinant lines up to isotopy, and orientations are induced by a particular equivalence class of deformations induced by the relative spin structure. In particular, the operator  $\tilde{D}_u$

is isotopic to a Fredholm operator  $\tilde{D}_u^\vee$  obtained by gluing together Cauchy-Riemann operators on collection of disks  $\bar{S}_v$  without punctures with the operators  $\tilde{D}_u^{e,\pm}$  over punctured surfaces  $S_e$  obtained whose boundary condition is given by a path  $\kappa$  in  $T_{\gamma(t)}(\mathbb{R} \times Z)$  connecting  $T_{\gamma(0)}(\mathbb{R} \times \Lambda)$  with  $T_{\gamma(1)}(\mathbb{R} \times \Lambda)$  in the family of isomorphic  $T_{\gamma(t)}\mathbb{R} \times Z$ ; see for example [47, Section 2.4]. Let  $\Sigma_{\text{ev}_e(u)}^\pm$  denote the stable resp. unstable manifolds associated to the critical point  $\text{ev}_e(u)$  at the end of the edge  $e$ . Define

$$\mathbb{D}_e^\pm := \det(\Sigma_{\text{ev}_e(u)}^\pm) \otimes \mathbb{D}_{\kappa(e)}$$

where  $\mathbb{D}_{\kappa(e)}$  is the determinant of the index of the Cauchy-Riemann operator  $\tilde{D}_u^{e,\pm}$  associated to the capping path  $\kappa$ , tensored with the determinant line  $\det(T_{\gamma(0)}\Lambda \cap T_{\gamma(1)}\Lambda)$  (using the identifications of  $T_{\gamma(t)}(\mathbb{R} \times Z)$ ). For each surface component  $S_v$  we have an isomorphism of  $\mathbb{D}_u$  with a product of determinant lines

$$(35) \quad \mathbb{D}_u \cong \det(T_{[S_v]}\mathcal{M}_\Gamma) \otimes \bigotimes_{e \in \mathcal{E}_-(\Gamma_v)} \mathbb{D}_e \otimes \mathbb{D}_u^\vee \otimes \bigotimes_{e \in \mathcal{E}_+(\Gamma_v)} \mathbb{D}_e^+.$$

Now the relative spin structure on  $L$  gives a deformation of  $\mathbb{D}_u^\vee$  to a combination of Cauchy-Riemann operators on spheres and operators on disks with trivial bundles and boundary conditions, as in [33] and [47, Section 4]. For the subset  $\mathcal{M}(L)_0$  of rigid maps denote by

$$\epsilon : \mathcal{M}(L)_0 \rightarrow \{\pm 1\}$$

the sign obtained by comparing the given orientation to the orientation of a point. In the case  $L = \mathbb{R} \times \Lambda$ , we obtain orientations on  $\mathcal{M}(\Lambda) = \mathcal{M}(L)/\mathbb{R}$  as well, and in particular a sign map  $\epsilon : \mathcal{M}(\Lambda)_0 \rightarrow \{\pm 1\}$ .

## REFERENCES

- [1] Casim Abbas. *An introduction to compactness results in symplectic field theory*. Springer, Heidelberg, 2014.
- [2] Mina Aganagic, Tobias Ekholm, Lenhard Ng, and Cumrun Vafa. Topological strings, D-model, and knot contact homology. *Adv. Theor. Math. Phys.*, 18(4):827–956, 2014.
- [3] Manabu Akaho and Dominic Joyce. Immersed Lagrangian Floer theory. *J. Differential Geom.*, 86(3):381–500, 2010.
- [4] Johan Asplund. Contact homology of Legendrian knots in five-dimensional circle bundles. <https://www.diva-portal.org/smash/get/diva2:940045/FULLTEXT01.pdf>.
- [5] Michèle Audin and Mihai Damian. *Morse theory and Floer homology*. Universitext. Springer, London; EDP Sciences, Les Ulis, 2014. Translated from the 2010 French original by Reinie Ern .
- [6] Denis Auroux, Damien Gayet, and Jean-Paul Mohsen. Symplectic hypersurfaces in the complement of an isotropic submanifold. *Mathematische Annalen*, 321(4):739–754, 2001.
- [7] Kenneth Blakey, Soham Chanda, Yuhan Sun, and Chris Woodward. Augmentation varieties and disk potentials II, 2024. <https://arxiv.org/abs/2401.13021>.

- [8] Kenneth Blakey, Soham Chanda, Yuhan Sun, and Chris Woodward. Augmentation varieties and disk potentials III, 2024. <https://arxiv.org/abs/2401.13024>.
- [9] Frédéric Bourgeois, Yakov Eliashberg, Helmut Hofer, Krzysztof Wysocki, and Eduard Zehnder. Compactness results in symplectic field theory. *Geom. Topol.*, 7:799–888, 2003.
- [10] Soham Chanda. Floer cohomology and higher mutations, 2023. <https://arxiv.org/abs/2301.08311>.
- [11] Baptiste Chantraine. Lagrangian concordance of Legendrian knots. *Algebr. Geom. Topol.*, 10(1):63–85, 2010.
- [12] Baptiste Chantraine. A note on exact Lagrangian cobordisms with disconnected Legendrian ends. *Proc. Amer. Math. Soc.*, 143(3):1325–1331, 2015.
- [13] François Charest and Chris Woodward. Floer trajectories and stabilizing divisors. *J. Fixed Point Theory Appl.*, 19(2):1165–1236, 2017.
- [14] François Charest and Chris T. Woodward. Floer cohomology and flips. *Mem. Amer. Math. Soc.*, 279(1372):v+166, 2022.
- [15] Yuri Chekanov. New invariants of Legendrian knots. In *European Congress of Mathematics, Vol. II (Barcelona, 2000)*, volume 202 of *Progr. Math.*, pages 525–534. Birkhäuser, Basel, 2001.
- [16] Cheol-Hyun Cho and Yong-Geun Oh. Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds. *Asian J. Math.*, 10(4):773–814, 2006.
- [17] Kai Cieliebak and Klaus Mohnke. Symplectic hypersurfaces and transversality in Gromov-Witten theory. *J. Symplectic Geom.*, 5(3):281–356, 2007.
- [18] Kai Cieliebak and Evgeny Volkov. First steps in stable Hamiltonian topology. *J. Eur. Math. Soc. (JEMS)*, 17(2):321–404, 2015.
- [19] Herbert Clemens. Curves on generic hypersurfaces. *Ann. Sci. École Norm. Sup. (4)*, 19(4):629–636, 1986.
- [20] Octav Cornea and François Lalonde. Cluster homology: an overview of the construction and results. *Electron. Res. Announc. Amer. Math. Soc.*, 12:1–12, 2006.
- [21] Thomas Delzant. Hamiltoniens périodiques et images convexes de l’application moment. *Bull. Soc. Math. France*, 116(3):315–339, 1988.
- [22] Georgios Dimitroglou Rizell and Roman Golovko. Legendrian submanifolds from Bohr–Sommerfeld covers of monotone Lagrangian tori. *Comm. Anal. Geom.*, 31(4):905–978, 2023.
- [23] Simon K. Donaldson. Symplectic submanifolds and almost-complex geometry. *Journal of Differential Geometry*, 44(4):666–705, 1996.
- [24] Tobias Ekholm. Morse flow trees and Legendrian contact homology in 1-jet spaces. *Geom. Topol.*, 11:1083–1224, 2007.
- [25] Tobias Ekholm, John Etnyre, and Michael Sullivan. The contact homology of Legendrian submanifolds in  $\mathbb{R}^{2n+1}$ . *J. Differential Geom.*, 71(2):177–305, 2005.
- [26] Tobias Ekholm, John Etnyre, and Michael Sullivan. Orientations in Legendrian contact homology and exact Lagrangian immersions. *Internat. J. Math.*, 16(5):453–532, 2005.



- [27] Tobias Ekholm, John Etnyre, and Michael Sullivan. Legendrian contact homology in  $P \times \mathbb{R}$ . *Trans. Amer. Math. Soc.*, 359(7):3301–3335, 2007.
- [28] Tobias Ekholm, John B. Etnyre, Lenhard Ng, and Michael G. Sullivan. Knot contact homology. *Geom. Topol.*, 17(2):975–1112, 2013.
- [29] Tobias Ekholm and Lenhard Ng. Higher genus knot contact homology and recursion for colored HOMFLY-PT polynomials. *Adv. Theor. Math. Phys.*, 24(8):2067–2145, 2020.
- [30] Yakov Eliashberg. Invariants in contact topology. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, number Extra Vol. II, pages 327–338, 1998.
- [31] Yakov Eliashberg, Alexander Givental, and Helmut Hofer. Introduction to symplectic field theory. *Geom. Funct. Anal.*, (Special Volume, Part II):560–673, 2000. GAFA 2000 (Tel Aviv, 1999).
- [32] Jonathan David Evans. Lectures on Lagrangian torus fibrations, 2022. <https://arxiv.org/abs/2110.08643>.
- [33] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono. *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*, volume 46 of *AMS/IP Studies in Advanced Mathematics*. American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2009.
- [34] Hansjörg Geiges. *An introduction to contact topology*, volume 109 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2008.
- [35] Roman Golovko. A note on Lagrangian cobordisms between Legendrian submanifolds of  $\mathbb{R}^{2n+1}$ . *Pacific J. Math.*, 261(1):101–116, 2013.
- [36] Victor Guillemin and Shlomo Sternberg. *Symplectic techniques in physics*. Cambridge University Press, Cambridge, second edition, 1990.
- [37] Reese Harvey and H. Blaine Lawson, Jr. Calibrated geometries. *Acta Math.*, 148:47–157, 1982.
- [38] Dominic Joyce. Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications. *J. Differential Geom.*, 63(2):279–347, 2003.
- [39] Eugene Lerman. Symplectic cuts. *Math. Res. Lett.*, 2(3):247–258, 1995.
- [40] Charles-Michel Marle. Sous-variétés de rang constant et sous-variétés symplectiquement régulières d’une variété symplectique. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(2):119–122, 1982.
- [41] Dusa McDuff and Dietmar Salamon. *J-holomorphic curves and symplectic topology*, volume 52 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, second edition, 2012.
- [42] Yong-Geun Oh. Riemann-Hilbert problem and application to the perturbation theory of analytic discs. *Kyungpook Math. J.*, 35(1):39–75, 1995.
- [43] Joseph Palmer and Chris Woodward. Invariance of immersed Floer cohomology under Lagrangian surgery, 2023. <https://arxiv.org/abs/1903.01943>.
- [44] James Pascaleff and Dmitry Tonkonog. The wall-crossing formula and Lagrangian mutations. *Adv. Math.*, 361:106850, 67, 2020.
- [45] Joshua M. Sabloff. Invariants of Legendrian knots in circle bundles. *Commun. Contemp. Math.*, 5(4):569–627, 2003.



- [46] Sushmita Venugopalan and Chris Woodward. Tropical Fukaya algebras, 2022. <https://arxiv.org/abs/2004.14314>.
- [47] Katrin Wehrheim and Chris Woodward. Orientations for pseudoholomorphic quilts, 2015. <https://arxiv.org/abs/1503.07803>.
- [48] Chris Wendl. Lectures on symplectic field theory. [https://www.mathematik.hu-berlin.de/~wendl/pub/SFTlectures\\_book3.pdf](https://www.mathematik.hu-berlin.de/~wendl/pub/SFTlectures_book3.pdf).

DEPARTMENT OF MATHEMATICS, MIT, 182 MEMORIAL DRIVE, CAMBRIDGE, MA 02139, U.S.A.  
*Email address:* `kblakey@mit.edu`

MATHEMATICS-HILL CENTER, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019, U.S.A.  
*Email address:* `sc1929@math.rutgers.edu`

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, SOUTH KENSINGTON, LONDON, SW7 2AZ, U.K.  
*Email address:* `yuhan.sun@imperial.ac.uk`

MATHEMATICS-HILL CENTER, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019, U.S.A.  
*Email address:* `ctw@math.rutgers.edu`