

An Asymptotic Local-Global Principle for Integral Kleinian Sphere Packings


Edna Jones

Rutgers University


March 2021

Apollonian Circle Packings: The Construction


Given three mutually tangent circles with disjoint points of tangency, there are exactly two circles tangent to the given ones. (Proved by Apollonius of Perga.)




Apollonian Circle Packings: The Construction



Apollonian Circle Packings: The Construction



Apollonian Circle Packings: The Construction



Apollonian Circle Packings: The Construction






Figure: An Apollonian circle packing.

Apollonian Circle Packings



Apollonian Circle Packings




Figure: An Apollonian circle packing.

Label on circle:
 $\text{bend} = 1/\text{radius}$

All of the bends of this Apollonian circle packing are integers.

Which integers appear as bends?

Are there any congruence or local obstructions?

Definition (Admissible integers for Apollonian circle packings)

Let \mathcal{P} be an integral Apollonian circle packing.

An integer m is **admissible (or locally represented)** if for every $q \geq 1$

$$m \equiv \text{bend of some circle in } \mathcal{P} \pmod{q}.$$

Equivalently, m is admissible if m has no local obstructions.


Admissible Integers

Theorem (Fuchs, 2011)

m is admissible if and only if m is in certain congruence classes modulo 24.

(The congruence classes depend on the packing.)

Example



m is admissible \iff
 $m \equiv 0, 4, 12, 13, 16, \text{ or } 21 \pmod{24}.$


Local-Global Conjecture

Conjecture (Graham–Lagarias–Mallows–Wilks–Yan, 2003)

The bends of a fixed primitive integral Apollonian circle packing \mathcal{P} satisfy an asymptotic local-global principle.

That is, there is an $N_0 = N_0(\mathcal{P})$ so that, if $m > N_0$ and m is admissible, then m is the bend of a circle in the packing.

Example



Why Do We Have a Local-Global Conjecture?

Theorem (Kontorovich–Oh, 2011)

The number of circles in an Apollonian circle packing \mathcal{P} with bend at most N (counted with multiplicity) is asymptotically equal to a constant times N^δ , where $\delta =$ the Hausdorff dimension of the closure of \mathcal{P} .

For Apollonian circle packings, we have

$$\delta \approx 1.30568\dots$$

Thus, we would expect that the multiplicity of a given admissible bend up to N is roughly $N^{\delta-1} \approx N^{0.30568} \geq 1$, so we should expect that every sufficiently large admissible number to be represented.

The Best We Can Do Right Now

Theorem (Bourgain–Kontorovich, 2014)

Almost every admissible number is the bend of a circle in a fixed primitive integral Apollonian circle packing \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$, where $\eta > 0$ is effectively computable.

The Best We Can Do Right Now

Theorem (Bourgain–Kontorovich, 2014)

Almost every admissible number is the bend of a circle in a fixed primitive integral Apollonian circle packing \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$, where $\eta > 0$ is effectively computable.

Proof outline:

- 1 Show that the automorphism group of the Apollonian circle packing contains the congruence subgroup $\Gamma(2)$ of $\mathrm{PSL}_2(\mathbb{Z})$, and $\Gamma(2)$ is the stabilizer of a particular circle. This implies that the set of bends contains primitive values of a shifted **binary** quadratic form. (Sarnak, 2007)

The Best We Can Do Right Now

Theorem (Bourgain–Kontorovich, 2014)

Almost every admissible number is the bend of a circle in a fixed primitive integral Apollonian circle packing \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$, where $\eta > 0$ is effectively computable.

Proof outline:

- 1 Show that the automorphism group of the Apollonian circle packing contains the congruence subgroup $\Gamma(2)$ of $\mathrm{PSL}_2(\mathbb{Z})$, and $\Gamma(2)$ is the stabilizer of a particular circle. This implies that the set of bends contains primitive values of a shifted **binary** quadratic form. (Sarnak, 2007)
- 2 The shifted **binary** quadratic form gives you enough to work with so that you can apply the circle method (with some other tools, including spectral theory, used in the major arc and minor arc analyses) to obtain an “almost all” statement.

Soddy Sphere Packings: The Construction

Given four mutually tangent spheres with disjoint points of tangency, there are exactly two spheres tangent to the given ones.




Figure: Four tangent spheres.




Figure: Four tangent spheres with two additional tangent spheres.

Soddy Sphere Packings: The Construction




Figure: Four tangent spheres.




Figure: Four tangent spheres with two additional tangent spheres.

Soddy Sphere Packings: The Construction




Figure: Four tangent spheres.




Figure: Four tangent spheres with two additional tangent spheres.




Figure: More tangent spheres.

Soddy Sphere Packings: The Construction




Figure: Four tangent spheres.




Figure: Four tangent spheres with two additional tangent spheres.




Figure: More tangent spheres.




Figure: A Soddy sphere packing.

Soddy Sphere Packings




Figure: A Soddy sphere packing made by Nicolas Hannachi.

Label on sphere:
 $\text{bend} = 1/\text{radius}$

Soddy Sphere Packings




Figure: A Soddy sphere packing made by Nicolas Hannachi.

Label on sphere:
 $\text{bend} = 1/\text{radius}$

All of the bends of this Soddy sphere packing are integers.

Soddy Sphere Packings




Figure: A Soddy sphere packing made by Nicolas Hannachi.

Label on sphere:
 $\text{bend} = 1/\text{radius}$

All of the bends of this Soddy sphere packing are integers.

Which integers appear as bends?

Soddy Sphere Packings




Figure: A Soddy sphere packing made by Nicolas Hannachi.

Label on sphere:
 $\text{bend} = 1/\text{radius}$

All of the bends of this Soddy sphere packing are integers.

Which integers appear as bends?

Are there any congruence or local obstructions?

Definition (Admissible integers for Soddy sphere packings)

Let \mathcal{P} be an integral Soddy sphere packing.

An integer m is **admissible (or locally represented)** if for every $q \geq 1$

$$m \equiv \text{bend of some sphere in } \mathcal{P} \pmod{q}.$$

Equivalently, m is admissible if m has no local obstructions.

Admissible Integers


Theorem (Kontorovich, 2019)

m is admissible in a primitive integral Soddy sphere packing \mathcal{P} if and only if

$$m \equiv 0 \text{ or } \varepsilon(\mathcal{P}) \pmod{3},$$

where $\varepsilon(\mathcal{P}) \in \{\pm 1\}$ depends only on the packing.

Example



m is admissible \iff
 $m \equiv 0 \text{ or } 1 \pmod{3}.$


An Asymptotic Local-Global Theorem

Theorem (Kontorovich, 2019)

The bends of a fixed primitive integral Soddy sphere packing \mathcal{P} satisfy an asymptotic local-global principle.

That is, there is an $N_0 = N_0(\mathcal{P})$ so that, if $m > N_0$ and m is admissible, then m is the bend of a sphere in the packing.

Example



If $m \equiv 0$ or $1 \pmod{3}$ and m is sufficiently large, then m is the bend of a sphere in the packing.

Proof Outline for Soddy Sphere Packing Result

- 1 Show that the automorphism group of the Soddy sphere packing contains a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z}[e^{\pi i/3}])$, and this congruence subgroup maps a particular sphere to itself. This implies that the set of bends contains “primitive” values of a shifted **quaternary** quadratic form.

Proof Outline for Soddy Sphere Packing Result

- 1 Show that the automorphism group of the Soddy sphere packing contains a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z}[e^{\pi i/3}])$, and this congruence subgroup maps a particular sphere to itself. This implies that the set of bends contains “primitive” values of a shifted **quaternary** quadratic form.
- 2 The shifted **quaternary** quadratic form gives you enough to work with so that you can quote the circle method to show that every sufficiently large admissible number is represented by the quadratic form without the primitivity restriction.

Proof Outline for Soddy Sphere Packing Result

- 1 Show that the automorphism group of the Soddy sphere packing contains a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z}[e^{\pi i/3}])$, and this congruence subgroup maps a particular sphere to itself. This implies that the set of bends contains “primitive” values of a shifted **quaternary** quadratic form.
- 2 The shifted **quaternary** quadratic form gives you enough to work with so that you can quote the circle method to show that every sufficiently large admissible number is represented by the quadratic form without the primitivity restriction.
- 3 Show that the singular series (with the primitivity restriction) is bounded away from zero when m is admissible.

Congruence Subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$

Definition (Principal congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$)

For an imaginary quadratic field K , a **principal congruence subgroup** of $\mathrm{PSL}_2(\mathcal{O}_K)$ is a subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ of the form

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathcal{O}_K) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{\varrho} \right\}$$

for a fixed ideal ϱ of \mathcal{O}_K .

Congruence Subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$

Definition (Principal congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$)

For an imaginary quadratic field K , a **principal congruence subgroup** of $\mathrm{PSL}_2(\mathcal{O}_K)$ is a subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ of the form

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathcal{O}_K) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{\varrho} \right\}$$

for a fixed ideal ϱ of \mathcal{O}_K .

Example (Soddy sphere packing, Kontorovich, 2019)

There exists a sphere $S_0 \in \mathcal{P}$ such that the stabilizer of S_0 in Γ contains (up to conjugacy) the congruence subgroup

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathcal{O}) : b, c \equiv 0 \pmod{\varrho} \right\},$$

where $\mathcal{O} = \mathbb{Z}[e^{\pi i/3}]$ and $\varrho = (1 + e^{\pi i/3})$.

Kleinian Sphere Packings

Definition (Kleinian sphere packing)

An $(n - 1)$ -sphere packing \mathcal{P} is **Kleinian** if its limit set is that of a geometrically finite group $\Gamma < \text{Isom}(\mathcal{H}^{n+1})$.




Figure: Apollonian circle packing as the limit set of Γ .

Definition (Kleinian sphere packing)

An $(n - 1)$ -sphere packing \mathcal{P} is **Kleinian** if its limit set is that of a geometrically finite group $\Gamma < \text{Isom}(\mathcal{H}^{n+1})$.






- Action of $\text{Isom}(\mathcal{H}^{n+1})$ extends continuously to $\widehat{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$, the boundary of \mathcal{H}^{n+1} .

Definition (Kleinian sphere packing)

An $(n - 1)$ -sphere packing \mathcal{P} is **Kleinian** if its limit set is that of a geometrically finite group $\Gamma < \text{Isom}(\mathcal{H}^{n+1})$.

- Action of $\text{Isom}(\mathcal{H}^{n+1})$ extends continuously to $\widehat{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$, the boundary of \mathcal{H}^{n+1} .
- Γ stabilizes \mathcal{P} (i.e., Γ maps \mathcal{P} to itself).

Examples of Kleinian Sphere Packings



Asymptotic Local-Global Principles

We want to prove asymptotic local-global principles for certain integral Kleinian sphere packings, that is, we want to prove the following:

If m is admissible and sufficiently large, then m is a bend of an $(n - 1)$ -sphere in the packing.

Definition (Admissible integers)

Let \mathcal{P} be an integral Kleinian sphere packing.

An integer m is **admissible (or locally represented)** if for every $q \geq 1$

$$m \equiv \text{bend of some } (n - 1)\text{-sphere in } \mathcal{P} \pmod{q}.$$

Group of Möbius Transformations

Group of orientation-preserving Möbius transformations

$$\text{Möb}(\mathcal{H}^{n+1}) < \text{Isom}(\mathcal{H}^{n+1})$$

$$\text{Möb}(\mathcal{H}^{n+1}) : \mathcal{H}^{n+1} \rightarrow \mathcal{H}^{n+1}$$

$$z \mapsto g(z) = (a \cdot z + b)(c \cdot z + d)^{-1},$$

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Möb}(\mathcal{H}^{n+1})$$

a, b, c, d in a Clifford algebra C_{n-1} with some restrictions.
Action extends to $\widehat{\mathbb{R}^n}$.

Group of Möbius Transformations

Group of orientation-preserving Möbius transformations
 $\text{Möb}(\mathcal{H}^{n+1}) < \text{Isom}(\mathcal{H}^{n+1})$

$$\text{Möb}(\mathcal{H}^{n+1}) : \mathcal{H}^{n+1} \rightarrow \mathcal{H}^{n+1}$$

$$z \mapsto g(z) = (a \cdot z + b)(c \cdot z + d)^{-1},$$

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Möb}(\mathcal{H}^{n+1})$$

a, b, c, d in a Clifford algebra C_{n-1} with some restrictions.
Action extends to $\widehat{\mathbb{R}^n}$.

Clifford algebras for small n :

- $C_1 = \mathbb{C}$
- $C_2 = \mathbb{H}$, the set of quaternions

Example ($n = 2$)

$\text{Möb}(\mathcal{H}^3) \cong \text{PSL}_2(\mathbb{C})$ acts on $\widehat{\mathbb{R}^2} \cong \widehat{\mathbb{C}}$ via

$$z \mapsto g(z) = (az + b)(cz + d)^{-1},$$

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2(\mathbb{C})$$

Example ($n = 2$)

$\text{Möb}(\mathcal{H}^3) \cong \text{PSL}_2(\mathbb{C})$ acts on $\widehat{\mathbb{R}^2} \cong \widehat{\mathbb{C}}$ via

$$z \mapsto g(z) = (az + b)(cz + d)^{-1},$$

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2(\mathbb{C})$$

Restrictions make defining what $\text{Möb}(\mathcal{H}^{n+1})$ is isomorphic to trickier for $n > 2$.

(For example, $\text{Möb}(\mathcal{H}^4) \not\cong \text{PSL}_2(\mathbb{H})$, even though $C_2 = \mathbb{H}$.)

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

Almost every admissible number is the bend of a circle in \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$.

Generalizes Bourgain–Kontorovich result for Apollonian circle packings to other Kleinian 1-sphere packings.

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

Suppose that

- \mathcal{P} is a primitive integral Kleinian 1-sphere packing in $\widehat{\mathbb{R}^2}$ with an automorphism group $\Gamma < \mathrm{PSL}_2(\mathfrak{a})$, where \mathfrak{a} is a fractional ideal of an imaginary quadratic field K ,

Almost every admissible number is the bend of a circle in \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$.

Generalizes Bourgain–Kontorovich result for Apollonian circle packings to other Kleinian 1-sphere packings.

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

Suppose that

- \mathcal{P} is a primitive integral Kleinian 1-sphere packing in $\widehat{\mathbb{R}^2}$ with an automorphism group $\Gamma < \mathrm{PSL}_2(\mathfrak{a})$, where \mathfrak{a} is a fractional ideal of an imaginary quadratic field K ,
- there exist circles S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and

Almost every admissible number is the bend of a circle in \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$.

Generalizes Bourgain–Kontorovich result for Apollonian circle packings to other Kleinian 1-sphere packings.

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

Suppose that

- \mathcal{P} is a primitive integral Kleinian 1-sphere packing in $\widehat{\mathbb{R}^2}$ with an automorphism group $\Gamma < \mathrm{PSL}_2(\mathfrak{a})$, where \mathfrak{a} is a fractional ideal of an imaginary quadratic field K ,
- there exist circles S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and
- the stabilizer of S_0 in Γ (up to conjugacy) contains a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z})$.

Almost every admissible number is the bend of a circle in \mathcal{P} . Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$.

Generalizes Bourgain–Kontorovich result for Apollonian circle packings to other Kleinian 1-sphere packings.

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

Suppose that

- \mathcal{P} is a primitive integral Kleinian 1-sphere packing in $\widehat{\mathbb{R}^2}$ with an automorphism group $\Gamma < \mathrm{PSL}_2(\mathfrak{a})$, where \mathfrak{a} is a fractional ideal of an imaginary quadratic field K ,
- there exist circles S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and
- the stabilizer of S_0 in Γ (up to conjugacy) contains a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z})$.

Almost every admissible number is the bend of a circle in $\Gamma \cdot S_1 \subseteq \mathcal{P}$. Quantitatively, the number of exceptions up to N is bounded by $O(N^{1-\eta})$.

Generalizes Bourgain–Kontorovich result for Apollonian circle packings to other Kleinian 1-sphere packings.

Result for Kleinian 1-Sphere (Circle) Packings

Theorem (Fuchs–Stange–Zhang, 2019)

If \mathcal{P} is a primitive integral Kleinian 1-sphere packing in $\widehat{\mathbb{R}^2}$ satisfying certain conditions, almost every admissible number is the bend of a circle in \mathcal{P} .




Figure: An integral Kleinian (more specifically, cuboctahedral) 1-sphere packing that satisfies the conditions of the theorem. Figure taken from “Local-Global Principles in Circle Packings” by Fuchs, Stange, and Zhang.

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z})$ implies that the set of bends of \mathcal{P} contains primitive values of a shifted **binary** quadratic form.

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathbb{Z})$ implies that the set of bends of \mathcal{P} contains primitive values of a shifted **binary** quadratic form.
- 2 The shifted **binary** quadratic form gives you enough to work with so that you can apply the circle method (with some other tools, including spectral theory, used in the major arc and minor arc analyses) to obtain an “almost all” statement.

Goal: Prove an asymptotic local-global principle for bends of certain integral Kleinian sphere packings.




Figure: An integral Kleinian (more specifically, an orthoplicial) sphere packing made by Kei Nakamura.




Figure: A fundamental domain of an integral Kleinian sphere packing made by Arseniy (Senia) Sheydvasser.

Suppose that

- $n \geq 3$,
- \mathcal{P} is a primitive integral Kleinian $(n - 1)$ -sphere packing in $\widehat{\mathbb{R}^n}$ with an automorphism group Γ of Möbius transformations,

Suppose that

- $n \geq 3$,
- \mathcal{P} is a primitive integral Kleinian $(n - 1)$ -sphere packing in $\widehat{\mathbb{R}^n}$ with an automorphism group Γ of Möbius transformations,
- there exist $(n - 1)$ -spheres S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and

Suppose that

- $n \geq 3$,
- \mathcal{P} is a primitive integral Kleinian $(n - 1)$ -sphere packing in $\widehat{\mathbb{R}^n}$ with an automorphism group Γ of Möbius transformations,
- there exist $(n - 1)$ -spheres S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and
- the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$, where K is an imaginary quadratic field.

Suppose that

- $n \geq 3$,
- \mathcal{P} is a primitive integral Kleinian $(n - 1)$ -sphere packing in $\widehat{\mathbb{R}^n}$ with an automorphism group Γ of Möbius transformations,
- there exist $(n - 1)$ -spheres S_0 and S_1 such that S_1 is in \mathcal{P} and is tangent to S_0 , and
- the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$, where K is an imaginary quadratic field.

Then every sufficiently large admissible integer is a bend of an $(n - 1)$ -sphere in $\Gamma \cdot S_1 \subseteq \mathcal{P}$.

Proof Methods Outline

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ implies that the set of bends of \mathcal{P} contains “primitive” values of a shifted **quaternary** quadratic form.

Proof Methods Outline

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ implies that the set of bends of \mathcal{P} contains “primitive” values of a shifted **quaternary** quadratic form.
- 2 This shifted **quaternary** quadratic form should give you enough to work with so that you can apply the circle method to show that every sufficiently large admissible number is represented as a bend.

Proof Methods Outline

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ implies that the set of bends of \mathcal{P} contains “primitive” values of a shifted **quaternary** quadratic form.
- 2 This shifted **quaternary** quadratic form should give you enough to work with so that you can apply the circle method to show that every sufficiently large admissible number is represented as a bend.
 - Major arcs (for main term that contains local information): use spectral theory and expander graphs

Proof Methods Outline

- 1 The assumption that the stabilizer of S_0 in Γ contains (up to conjugacy) a congruence subgroup of $\mathrm{PSL}_2(\mathcal{O}_K)$ implies that the set of bends of \mathcal{P} contains “primitive” values of a shifted **quaternary** quadratic form.
- 2 This shifted **quaternary** quadratic form should give you enough to work with so that you can apply the circle method to show that every sufficiently large admissible number is represented as a bend.
 - Major arcs (for main term that contains local information): use spectral theory and expander graphs
 - Minor arcs (for error term): use Kloosterman circle method

Quadratic Form for Soddy Sphere Packings

Example (Quadratic form for Soddy sphere packings)

Shifted quaternary quadratic form in $a_0, a_1, c_0, c_1 \in \mathbb{Z}$:

$$\hat{\beta} |C(a_0 + a_1\omega) + D\rho(c_0 + c_1\omega)|^2 - Dj\bar{C} + Cj\bar{D},$$

$$\omega = e^{\pi i/3}$$

$$\rho = 1 + \omega$$

$$\gcd(a_0 + a_1\omega, \rho(c_0 + c_1\omega)) = 1$$

$\hat{\beta} \in \mathbb{R}$ and $C, D \in \mathbb{H}$ only depend on the packing.

(Scale appropriately to obtain a primitive integral quadratic form.)

Obtained by examining how Möbius transformations affect bends of $(n - 1)$ -spheres

Besides the illustrations previously credited and a few Apollonian circle packing construction illustrations created by the presenter, the illustrations for this talk came from the following papers:

- Jean Bourgain and Alex Kontorovich, “On the local-global conjecture for integral Apollonian gaskets,” *Inventiones mathematicae*, volume 196, pp. 589–650, 2014.
- Alex Kontorovich, “From Apollonius to Zaremba: Local-global phenomena in thin orbits,” *Bulletin of the American Mathematical Society*, volume 50, number 2, pp. 187-228, 2013, <https://www.ams.org/journals/bull/2013-50-02/S0273-0979-2013-01402-2/>.
- Alex Kontorovich, “The Local-Global Principle for Integral Soddy Sphere Packings,” *Journal of Modern Dynamics*, volume 15, pp. 209-236, 2019, <https://www.aims sciences.org/article/doi/10.3934/jmd.2019019>.

Thank you for listening!