

## MATH 575 ASSIGNMENT 6

1. Consider the approximation of the one dimensional heat equation  $u_t = \sigma u_{xx}$  by the generalized Crank-Nicholson method:

$$\frac{u_j^{n+1} - u_j^n}{k} = \frac{\sigma}{h^2} \{ (1 - \theta)[u_{j+1}^n - 2u_j^n + u_{j-1}^n] + \theta[u_{j+1}^{n+1} - 2u_j^{n+1} + u_{j-1}^{n+1}] \}, \quad 0 \leq \theta \leq 1.$$

a) Show the the local truncation error of this method is  $O(k + h^2)$ .

b) If  $u$  satisfies the heat equation and is sufficiently smooth and  $\theta = 1/2$ , show that the error in part (a) is  $O(k^2 + h^2)$ .

2. Write a Matlab program to approximate the initial boundary value problem:

$$\begin{aligned} u_t &= u_{xx}, & 0 < x < 1, & \quad t > 0, \\ u(0, t) &= u(1, t) = 0, & u(x, 0) &= x(1 - x) \end{aligned}$$

by the explicit method

$$[u_j^{n+1} - u_j^n]/k = [u_{j+1}^n - 2u_j^n + u_{j-1}^n]/h^2$$

for the three mesh sizes (i)  $k = 1/256$ ,  $h = 1/16$ , (ii)  $k = 1/512$ ,  $h = 1/16$ , and (iii)  $k = 1/1024$ ,  $h = 1/16$ . Plot the approximate solution and true solution on the same plot at  $t = 1$ . Explain your results. Note that the true solution

$$u(x, t) = \frac{8}{\pi^3} \sum_{n=1,3,5,\dots} \frac{1}{n^3} e^{-n^2 \pi^2 t} \sin(n\pi x)$$

and the first two terms of the series give a good approximation. See the file *heatsoln.m* for some help in defining and plotting the exact solution.

3. Consider the approximation of the transport equation  $u_t + au_x = 0$  by the difference scheme:

$$L_{h,k} U_j^n \equiv \left[ U_j^{n+1} - \frac{1}{2}(U_{j+1}^n + U_{j-1}^n) \right] / k + a[U_{j+1}^n - U_{j-1}^n]/(2h) = 0.$$

a) Show that if  $h = O(k)$ , the local truncation error of the method is  $O(k)$ .

b) Show that if  $\lambda \equiv ak/h$  satisfies  $|\lambda| \leq 1$ , then

$$\max_{j, 0 \leq n \leq N} |W_j^n| \leq \max_j |W_j^0| + T \max_{j, 0 \leq n \leq N-1} |L_{h,k} W_j^n|$$

for all mesh functions  $W_j^n$ , where  $T = Nk$ .

c) Show that for the pure initial value problem

$$u_t + au_x = 0, \quad u(x, 0) = \phi(x),$$

the error estimate,  $\max_j |u - U| \leq O(k)$  is valid for  $0 \leq n \leq N$  under the hypothesis of part (b).