

CONDENSATION AND LEFT-ORDERABLE GROUPS

FILIPPO CALDERONI AND ADAM CLAY

ABSTRACT. We discuss condensed left-orderings and develop new techniques to show that the conjugacy relation on the space of left-orderings is not smooth. These techniques apply to the solvable Baumslag Solitar groups $BS(1, n)$ and to Thompson's group F .

1. INTRODUCTION

The study of definable quotients of Polish spaces is one of the main themes in modern descriptive set theory, with the primary goal being to understand the Borel structure of Polish spaces modulo analytic equivalence relations. A fundamental question is whether the quotient space, equipped with the quotient Borel structure, is standard. The first trace of such an analysis dates back to the work of Glimm [Gli61] and Effros [Eff65].

If G is a countable group acting continuously on the Polish space X , we denote by X/G the space of orbits, endowed with the quotient Borel structure. In this case X/G is standard if and only if the orbit equivalence relation on X induced by the G -action is *smooth*. That is, if and only if there is a Borel map $\theta: X \rightarrow \mathbb{R}$ such that x_1, x_2 lie in the same G -orbit if and only if $\theta(x_1) = \theta(x_2)$. It is owing to this definition that for the remainder of the manuscript, we assume that all groups are countable unless otherwise indicated.

Following this trend, a question posed by Derooin, Navas, and Rivas [DNR16] raised the problem of whether the space of left-orderings $LO(G)$ of a left-orderable group G modulo the conjugacy G -action is always standard, or equivalently, whether or not the orbit equivalence relation is always smooth. Using descriptive set theory to demonstrate non-smoothness, the authors of this manuscript showed that $LO(G)/G$ is not a standard Borel space in many cases; for example, when G is a non-abelian free group or a free product of left-orderable groups [CC22, CC23].

Denote the equivalence relation induced by the conjugacy action of a group G on its space of left-orderings by $E_{lo}(G)$, and let $BS(1, n)$ denote the Baumslag-Solitar group. In this manuscript, we show:

2020 *Mathematics Subject Classification*. Primary: 03E15, 06F15, 20F60.

The first author would like to thank Justin Moore for asking about the Borel complexity of the conjugacy action of Thompson's group F on its space of left-orderings. This prompted the investigation of Section 3.5.

Theorem 1.1. (1) *For all $n \geq 2$, $E_{\text{lo}}(\text{BS}(1, n))$ is not smooth.*
 (2) *The conjugacy equivalence relation $E_{\text{lo}}(\text{BS}(1, 2))$ is Borel bi-reducible with E_0 .*

The novelty of Theorem 1.1 is twofold. Theorem 1.1(1) provides the first examples of left-orderable solvable groups G with non-standard quotient $\text{LO}(G)/G$. Moreover, Theorem 1.1(2) shows the first example of a finitely generated group G for which $E_{\text{lo}}(G)$ is not smooth, yet hyperfinite.

Using similar techniques, we also show how work of Navas implies that $E_{\text{lo}}(G)$ is not smooth whenever $\text{LO}(G)$ contains isolated points, and are able to tackle Thompson's group F in a similar manner.

Theorem 1.2. *For Thompson's group F , the conjugacy relation $E_{\text{lo}}(F)$ is not smooth.*

Central to our analysis is the idea of condensed left-orderings in $\text{LO}(G)$, which are orderings that can be approximated by their conjugates. Their existence turns out to be equivalent to non-smoothness of $E_{\text{lo}}(G)$ (Proposition 2.2), moreover, they can be detected by analyzing $\text{LO}((G))$, the free part of the conjugacy G -action on $\text{LO}(G)$ (See Proposition 3.7).

2. CONDENSED POINTS

A Polish space is a separable and completely metrizable topological space. For a Polish space X we denote by $F(X)$ the Effros standard Borel space of closed subsets of X . The standard Borel structure on $F(X)$ is generated by the sets

$$F_U = \{F \in F(X) \mid F \cap U \neq \emptyset\}$$

for all open $U \subseteq X$.

An equivalence relation E on the Polish space X is Borel if $E \subseteq X \times X$ is a Borel subset of $X \times X$. Most of the Borel equivalence relations that we will consider in this paper arise from group actions as follows. Let G be a countable group. Then a Polish G -space is a Polish space X equipped with a continuous action $(g, x) \mapsto g \cdot x$ of G on X . The corresponding G -orbit equivalence relation on X , which we will denote by E_G^X , is a Borel equivalence relation with countable classes. Let G be a countable discrete group and let X be a fixed Polish G -space. For a subgroup $H \leq G$, denote $\text{Orb}_H(x)$ the orbit of x under the induced H -action. Whenever $G = H$, we let $\text{Orb}(x) = \text{Orb}_G(x)$.

Recall that a Borel equivalence relation E is *smooth* if there exist a standard Borel space Y and a Borel map $\theta: X \rightarrow Y$ such that

$$x_1 E x_2 \iff \theta(x_1) = \theta(x_2).$$

An equivalence relation on a Polish space is *generically ergodic* if every invariant set with the Baire property is meager or comeager. Whenever X is a Polish G -space, the following are equivalent:

- (i) E_G^X is generically ergodic.

(ii) There is $x \in X$ such that $\text{Orb}(x)$ is dense in X .

Generic ergodicity is an obstruction to smoothness in many cases. In this manuscript, we will use the following fact:

Proposition 2.1. *Suppose that X is a Polish G -space with no isolated points. If E_G^X is generically ergodic, then E_G^X is not smooth.*

Following the terminology of Osin [Osi21b, Osi21a], we say that a point $x \in X$ is *condensed* if it is an accumulation point of $\text{Orb}(x)$.

The next proposition is essentially due to Osin [Osi21b, Proposition 2.7], who analyzed condensation in the Polish space of finitely generated marked groups. Since we could not find the proof in the literature, we give the proof of this general fact below.

Proposition 2.2. *Let G act on X continuously. Then the following are equivalent:*

- (1) E_G^X is smooth.
- (2) There are no condensed points in X .

Proof. Suppose that E_G^X is smooth and let x be any element of X . Consider the closed G -invariant set $Y = \overline{\text{Orb}(x)}$. If E_G^X is smooth, then E_G^Y is also smooth. As $\text{Orb}(x)$ is a dense G -orbit in Y , the action $G \curvearrowright Y$ is generically ergodic, so there must be an isolated point in $x_0 \in Y$. The point x_0 cannot be an element of $Y \setminus \text{Orb}(x)$ since these points are non-isolated by definition, and so $x_0 \in \text{Orb}(x)$. Now as the G -action is continuous, every point of $\text{Orb}(x)$, and in particular x itself, must be isolated in the subspace topology. It follows that x cannot be a condensed point.

On the other hand, suppose that no $x \in X$ is a condensed point. Then for every $x \in X$, the subspace topology on $\text{Orb}(x)$ is discrete, and since $\text{Orb}(x)$ is countable it is therefore Polish. By Alexandrov's theorem, $\text{Orb}(x)$ must be a G_δ set for all $x \in X$. (See [Kec95, Theorem 3.11].) Further note that the saturation of an arbitrary open set $U \subset X$ is itself open, since the saturation can be written as a union of the sets $g \cdot U$ where $g \in G$, each of which is open since G acts continuously. This implies that the map $X \rightarrow F(X), x \mapsto \overline{\text{Orb}(x)}$ is Borel showing that E_G^X is Borel reducible to $=_{F(X)}$. (E.g., see [Gao09, Exercise 5.4.8].) \square

3. THE CONJUGACY RELATION ON THE SPACES OF LEFT-ORDERINGS

A group G is *left-orderable* if it admits a strict total ordering $<$ such that $g < h$ implies $fg < fh$ for all $f, g, h \in G$.

Proposition 3.1. *The following are equivalent:*

- (1) G is left-orderable.
- (2) There is $P \subseteq G$ such that
 - (a) $P \cdot P \subseteq G$;
 - (b) $P \sqcup P^{-1} = G \setminus \{id\}$.

(3) *There is a totally ordered set $(\Omega, <)$ such that $G \hookrightarrow \text{Aut}(\Omega, <)$.*

The subset P in (2) above is referred to as a *positive cone*. Every left-ordering $<$ of G determines a positive cone $P_< = \{g \in G : g > id\}$. The identification of left-orderings with the corresponding positive cones allows us to define the space of left-orderings as follows. Equip $\{0, 1\}$ with the discrete topology, $\{0, 1\}^G$ with the product topology, and set

$$\text{LO}(G) = \{P \subset G : P \text{ is a positive cone}\} \subset \{0, 1\}^G,$$

equipped with the subspace topology. Note that the subbasic open sets in $\text{LO}(G)$ are the sets of the form $U_g = \{P : g \in P\}$, where $g \in G \setminus \{id\}$. One can easily check that $\text{LO}(G)$ is a closed subset of $\{0, 1\}^G$, hence a compact Polish space. We regard $\text{LO}(G)$ as a Polish G -space in the following precise sense. There is a G -action by homeomorphisms on $\text{LO}(G)$, given by $g \cdot P = gPg^{-1}$. As mentioned in the introduction, we denote by $E_{\text{lo}}(G)$ the orbit equivalence relation on $\text{LO}(G)$ induced by the conjugacy G -action.

3.1. Smoothness of $E_{\text{lo}}(G)$ and relatively convex subgroups. Let G be a group equipped with a fixed left-ordering $<$. A subgroup C of G is *convex relative to $<$* if whenever $g, h \in C$ and $f \in G$ with $g < f < h$, then $f \in C$. A subgroup $C \subseteq G$ is *left-relatively convex* in G if C is convex relative to some left ordering of G .

First we leverage the dichotomy established in Proposition 2.2 to prove that the non-smoothness of $E_{\text{lo}}(G)$ is detected by relatively convex subgroups.

Theorem 3.2. *For a left-orderable group G , the following are equivalent:*

- (1) $E_{\text{lo}}(G)$ is smooth.
- (2) *For every relatively convex $C \leq G$ the conjugacy orbit equivalence relation $E_{\text{lo}}(C)$ is smooth.*

Proof. The only non-trivial implication is (1) \implies (2). Fix a relatively convex $C \leq G$ such that $E_{\text{lo}}(C)$ is not smooth. It follows from Proposition 2.2 that there is a positive cone $Q \in \text{LO}(C)$ that is an accumulation point in $\text{Orb}_C(Q)$. Since C is relatively convex in G we can find some positive cone $P \in \text{LO}(G)$ such that $Q = P \cap C$. We are going to show that P is an accumulation point of $\text{Orb}_G(P)$. We will need the following:

Claim. If $c \in C$ and $P \in \bigcap_{i=1}^n U_{g_i}$ for $g_i \in G \setminus C$, then $cPc^{-1} \in \bigcap_{i=1}^n U_{g_i}$.

Proof of the Claim. Assume that $c \in P$. Then since C is convex with respect to $<_P$, we have $c <_P g_i$ for all $i = 1, \dots, n$. So, for $i = 1, \dots, n$ we have $c^{-1}g_i \in P$ and therefore $c^{-1}g_i c \in P$. We obtain $g_i \in cPc^{-1}$ for all $i \leq n$, therefore $cPc^{-1} \in \bigcap_{i=1}^n U_{g_i}$.

Next suppose that $c \in P^{-1}$. Then $g_i^{-1} <_P c$ for all $i = 1, \dots, n$ since C is convex. Therefore $g_i c \in P$ and thus $c^{-1}g_i c \in P$ and we conclude as in the previous case, completing the proof of the claim. \square

Now let $P \in \bigcap_{i=1}^n U_{c_i} \cap \bigcap_{j=1}^m U_{g_j}$ for some $c_1, \dots, c_n \in C$ and $g_1, \dots, g_m \in G \setminus C$. Since $P \cap C \in \text{Orb}_C(P \cap C)$ there exists $c \in C$ such that

$$c(P \cap C)c^{-1} \neq P \cap C \quad \text{and} \quad c(P \cap C)c^{-1} \in \bigcap_{i=1}^n U_{c_i}.$$

Then $cPc^{-1} = c(P \cap C)c^{-1} \cup c(P \setminus C)c^{-1} \neq P$, and $cPc^{-1} \in \bigcap_{i=1}^n U_{c_i} \cap \bigcap_{j=1}^m U_{g_j}$ by the previous claim. \square

Remark 3.3. Note that the condition on relatively convex subgroups in Theorem 3.2(2) cannot be replaced with a condition on *proper* relatively convex subgroups, as the example below shows (see also [CC22]).

Consider the infinitely generated group ¹

$$H_\infty = \langle x_1, x_2, \dots \mid x_i x_{i-1} x_i^{-1} = x_{i-1}^{-1} \text{ for } 1 < i \text{ and } x_i x_j = x_j x_i \text{ for } |i - j| > 1 \rangle.$$

Then for every left-ordering of H_∞ , one can show the convex subgroups are precisely the finitely generated subgroups of the form $H_n = \langle x_1, x_2, \dots, x_n \rangle$ where $j \geq 1$. This follows from first observing that every element H_j can be represented by a word of the form

$$x_1^{a_1} x_2^{a_2} \dots x_j^{a_j}$$

where $a_i \in \mathbb{Z}$, by using repeated applications of $x_j x_{j-1} = x_{j-1}^{-1} x_j$ and $x_j x_i = x_i x_j$ for all $i < j-1$ to shuffle all occurrences of x_j to the right hand side of any representative word. By writing every element of H_∞ in this form, it is straightforward to check that H_j is convex relative to every left-ordering of H_∞ . Moreover, there are no other relatively convex subgroups aside from the subgroups H_j . For if C were such a subgroup, there would exist j such that $H_j \leq C \leq H_{j+1}$. But then C should descend to a convex subgroup of $H_{j+1}/H_j \cong \mathbb{Z}$ under the quotient map, which is only possible if $C = H_j$ or $C = H_{j+1}$ since there are no proper, nontrivial convex subgroups in \mathbb{Z} .

Now one observes that the left-orders of H_∞ are in bijective correspondence with sequences $(\epsilon_i) \in \{0, 1\}^{\mathbb{N}}$ that encode the signs of the generators: for example we can set $x_i > id$ if and only if $\epsilon_i = 1$. It is not hard to see that the conjugacy action of H_∞ on the set $\text{LO}(H_\infty)$ yields an action of H_∞ on $\{0, 1\}^{\mathbb{N}}$ given by $x_j \cdot (\epsilon_i)$ is the same as (ϵ_i) in every entry except the $(j-1)$ -th position, which has been changed. Two left-orderings of H_∞ are in the same orbit if and only if their corresponding sequences in $\{0, 1\}^{\mathbb{N}}$ are eventually equal.

Thus every relatively convex proper subgroup $C \leq H_\infty$ is a Tararin group, so $\text{LO}(C)$ is finite, and yet $E_{\text{lo}}(H_\infty)$ is not smooth.

¹This example also appears in [CC22, Example 2.10], where there is a typo in the group presentation which is corrected here. We acknowledge Meng Che “Turbo” Ho for finding the typo and suggesting how to fix it.

3.2. Non-smoothness and isolated points. Recall that a positive cone P determines a *Conradian* left-ordering of G if $g, h \in P$ implies $g^{-1}hg^2 \in P$ for all $g, h \in G$ [Nav10]. Given a positive cone $P \in \text{LO}(G)$, a subgroup $C \leq G$ that is maximal with respect to the conditions:

- (1) C is convex relative to the ordering $<_P$ of G , and
- (2) $P \cap C$ determines a Conradian left-ordering of C

is called the *Conradian soul* of $<_P$.

We recall the following theorem proved via different techniques in both [Cla10] and [Nav10].

Theorem 3.4. *If the Conradian soul of $<_P$ is trivial, then P is condensed.*

Thus if G admits a positive cone P having trivial Conradian soul, then $E_{\text{lo}}(G)$ is not smooth.

As a consequence of Theorem 3.4, every isolated point in $\text{LO}(G)$ (that is, $P \in \text{LO}(G)$ such that there exist $g_1, \dots, g_n \in G \setminus \{id\}$ with $\{P\} = \bigcup_{i=1}^n U_{g_i}$) must have nontrivial Conradian soul, as isolated points cannot be condensed. In fact, Navas shows much more:

Theorem 3.5. [Nav10, Proposition 4.9] *Suppose that P is an isolated point and let $C \leq G$ its Conradian soul. Then C is a Tararin group, so $\text{LO}(C) = \{Q_1, \dots, Q_{2^k}\}$ for some $k > 0$, moreover, if G is not a Tararin group then there exists $i \in \{1, \dots, 2^k\}$ such that $P \setminus C \cup Q_i$ is a condensed point of $\text{LO}(G)$.*

As an immediate consequence, we apply Proposition 2.2 and observe:

Corollary 3.6. *Suppose that G is not a Tararin group. If $\text{LO}(G)$ contains an isolated point, then $E_{\text{lo}}(G)$ is not smooth.*

3.3. The free part of $\text{LO}(G)/G$. For a left-orderable group G denote by $\text{LO}((G))$ the free part of its conjugacy action. That is, we set

$$\text{LO}((G)) = \{P \in \text{LO}(G) : \forall g \neq 1 (g^{-1}Pg \neq P)\}.$$

Note that for any $P \in \text{LO}((G))$, the orbit $\text{Orb}(P)$ is infinite. For if $g^{-1}Pg = h^{-1}Ph$, then we get $hg^{-1}Ph^{-1} = P$ which implies $hg^{-1} = gh^{-1} = 1$ because P has trivial stabilizer.

Proposition 3.7. *If $\text{LO}((G)) \neq \emptyset$, then $E_{\text{lo}}(G)$ is not smooth.*

Proof. Suppose $P \in \text{LO}((G))$. By Proposition 2.2 it suffices to show that P is condensed. Let $P \in \bigcap_{i=1}^n U_{g_i}$, which is a basic open neighborhood of P . And assume that $g_1 <_P \dots <_P g_n$ without loss of generality. Then, we claim that

$$g_1^{-1}g_i g_1 \in P$$

for $i = 1, \dots, n$. For $i = 1$, it follows from the assumption that $P \in U_{g_1}$. For $i \geq 2$, $g_1 <_P g_i$ implies that $1 <_P g_1^{-1} g_i$, whence $1 <_P g_1^{-1} g_i g_1$ because P is a semigroup. Therefore, for $i = 1, \dots, n$, we have

$$g_i \in g_1 P g_1^{-1}.$$

This shows that $g_1 P g_1^{-1} \in \text{Orb}(P) \cap \bigcap_{i=1}^n U_{g_i}$. Since $P \in \text{LO}((G))$, we conclude that $g_1^{-1} P g_1 \neq P$, therefore P is condensed. \square

3.4. Baumslag-Solitar groups. Fix an integer n . The Baumslag-Solitar group $\text{BS}(1, n)$ is given by the presentation $\langle a, b \mid bab^{-1} = a^n \rangle$. There is an injective homomorphism $\rho: \text{BS}(1, n) \rightarrow \text{Homeo}_+(\mathbb{R})$ defined by setting

$$\begin{aligned} \rho(a)(x) &= x + 1, \\ \rho(b)(x) &= nx. \end{aligned}$$

The following construction of left-orderings on $\text{BS}(1, n)$ is due to Smirnov [Smi66]. For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ we can define a corresponding $P_\alpha \in \text{LO}(\text{BS}(1, n))$ by declaring

$$g \in P_\alpha \iff \rho(g)(\alpha) > \alpha.$$

Note that the map $\mathbb{R} \setminus \mathbb{Q} \rightarrow \text{LO}(\text{BS}(1, n))$ is injective. In fact, for different irrational numbers $\alpha < \beta$, we can find some $g \in \text{BS}(1, n)$ such that $g \in P_\beta \setminus P_\alpha$ or $g \in P_\alpha \setminus P_\beta$. To see this, we argue by cases:

Case 1. $\beta > 0$: Let $q \in (\alpha, \beta) \cap \mathbb{Q}^+$ and let $g \in \text{BS}(1, n)$ such that $\rho(g)(x) = \frac{q+1}{q}x - 1$. It is immediate to check that $\rho(g)(\beta) > \beta$ and $\rho(g)(\alpha) < \alpha$, therefore $g \in P_\beta$ and $g \notin P_\alpha$.

Case 2. $\beta < 0$: Let $q \in (\alpha, \beta) \cap \mathbb{Q}$ and let $g \in \text{BS}(1, n)$ such that $\rho(g)(x) = \frac{q+1}{q}x - 1$. Similarly to the previous cases, one can check that $\rho(g)(\alpha) < \alpha$ and $\rho(g)(\beta) < \beta$, therefore $g \in P_\alpha$ and $g \notin P_\beta$.

It is well known that the conjugacy action $\text{BS}(1, n) \curvearrowright \text{LO}(\text{BS}(1, n))$ is not generically ergodic, however with our new technique we can easily prove the following:

Corollary 3.8. *For $n > 1$ and $G = \text{BS}(1, n)$, the conjugacy relation $E_{\text{lo}}(G)$ is not smooth.*

Proof. By Proposition 3.7 it suffices to prove that for any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, the positive cone P_α belongs to the free part of the conjugacy action. To see this, assume that $hP_\alpha h^{-1} = P_\alpha$. One checks that $hP_\alpha h^{-1} = P_{\rho(h)(\alpha)}$, and therefore $P_{\rho(h)(\alpha)} = P_\alpha$, which gives $\rho(h)(\alpha) = \alpha$. However, for every $h \neq 1$, the order-preserving homeomorphism $\rho(h)$ has only rational fixed points. Therefore, it must hold that $h = 1$ as desired. \square

It is worth pointing out that Corollary 3.8 also follows from Proposition 2.2 and the work of Rivas and Tessera [RT16, Proposition 2.12]. However, our analysis of the free part of $\text{BS}(1, 2) \curvearrowright \text{LO}(\text{BS}(1, 2))$ allows us to further prove the following:

Corollary 3.9. *$E_{\text{lo}}(\text{BS}(1, 2))$ is hyperfinite.*

Proof. Let $\Gamma = \text{BS}(1, 2)$. The set $Y = \{P_\alpha \mid \alpha \in \mathbb{R} \setminus \mathbb{Q}\}$ is Borel because it is the image of a Borel set through a one-to-one Borel map (see [Kec95, Theorem 15.1]), and moreover Y is invariant under conjugacy. Therefore Y with the standard Borel structure induced by $\text{LO}(\Gamma)$ is a free standard Borel Γ -space. Moreover, Rivas [Riv10, Theorem 4.2] establishes that $\text{LO}(\Gamma) \setminus Y$ is countable. It follows that $E_{\text{lo}}(\Gamma) \sim_B E_\Gamma^Y$, and the latter equivalence relation is hyperfinite by [CJM⁺ar, Corollary 7.4]. \square

To the best of our knowledge Corollary 3.9 provides the first example of finitely generated left-orderable group, whose conjugacy equivalence relation is not smooth, yet hyperfinite.

3.5. Thompson's group F . Thompson's group F may be defined by the presentation

$$F = \langle a, b \mid [ab^{-1}, a^{-1}ba], [ab^{-1}, a^{-2}ba^2] \rangle.$$

There is an injective homomorphism $\rho: F \rightarrow \text{PL}_+([0, 1])$ whose image consists of all piecewise linear homeomorphisms of $[0, 1]$ having dyadic rational breakpoints, and whose linear segments have slopes that are integral powers of two.

Given an interval $I = [\frac{p}{2^q}, \frac{p+1}{2^q}] \subset [0, 1]$, we can define functions $b_I^+, b_I^-: [0, 1] \rightarrow [0, 1]$ that lie in the image of ρ and whose support is equal to I , as follows. First, the function b_I^+ is given by

$$b_I^+(t) = \begin{cases} t & \text{if } 0 \leq t \leq \frac{p}{2^q} \\ 2t - \frac{p}{2^q} & \text{if } \frac{p}{2^q} \leq t \leq \frac{p}{2^q} + \frac{1}{2^{q+2}} \\ t + \frac{1}{2^{q+2}} & \text{if } \frac{p}{2^q} + \frac{1}{2^{q+2}} \leq t \leq \frac{p}{2^q} + \frac{1}{2^{q+1}} \\ \frac{1}{2}t + \frac{p+1}{2^{q+1}} & \text{if } \frac{p}{2^q} + \frac{1}{2^{q+1}} \leq t \leq \frac{p+1}{2^q} \\ t & \text{if } \frac{p+1}{2^q} \leq t \leq 1 \end{cases}$$

It is clear from this description that b_I^+ lies in the image of ρ . On the interval I , the graph of b_I^+ appears as in Figure 1(A). We can analogously define b_I^- , which is the identity outside of I and whose graph appears as in Figure 1(B).

Proposition 3.10. *Let $S \subset [0, 1]$ be finite, and choose $x, y \in [0, 1] \setminus S$ with $x \neq y$. Then there exists $g \in F$ such that $\rho(g)(s) = s$ for all $s \in S$, $\rho(g)(x) > x$, and $\rho(g)(y) < y$.*

Proof. As the dyadic rational numbers are dense, we may choose disjoint intervals I, J with dyadic rational endpoints, satisfying $I \cap S = J \cap S = \emptyset$, with $x \in I$ and $y \in J$. Now set $f = b_I^+ \circ b_J^-$, then f satisfies $f(s) = s$ for all $s \in S$, and $f(x) > x$ while $f(y) < y$. Moreover, f is in the image of ρ , so the proposition follows. \square

Fix an enumeration $e: \mathbb{N} \rightarrow \mathbb{Q} \cap (0, 1)$, writing $e(i) = r_i$. Every enumeration of \mathbb{Q} can be used to define a positive cone $P_e \subset F$ in the usual way: Given $g \in F$, let r_i




FIGURE 1. The graphs of the functions b_I^+ and b_I^- on the interval I .

denote the first rational number in the enumeration satisfying $\rho(g)(r_i) \neq r_i$. Then declare $g \in P_e$ if and only if $\rho(g)(r_i) > r_i$.

Theorem 1.2 now follows from the following:

Proposition 3.11. *For every enumeration $e : \mathbb{N} \rightarrow \mathbb{Q} \cap (0, 1)$, we have $P_e \in \text{LO}((F))$.*

Proof. Let $h \in F \setminus \{id\}$ and suppose that $\rho(h)(r_i) = r_i$ for all $i < N$, while $\rho(h)(r_N) \neq r_N$. Set $S = \{r_0, \dots, r_{N-1}\}$, $x = r_N$ and $y = \rho(h)(r_N)$. Apply Proposition 3.10 to arrive at $g \in F$ with $\rho(g)(r_i) = r_i$ for all $i < N$ and $\rho(g)(r_N) > r_N$, so that $g \in P_e$. On the other hand, $\rho(h^{-1}gh)(r_i) = r_i$ for all $i < N$, while $\rho(g)(\rho(h)(r_N)) < \rho(h)(r_N)$ holds by our choice of g , which is equivalent to $\rho(h^{-1}gh)(r_N) < r_N$. Thus $h^{-1}gh \notin P_e$, meaning $g \notin hP_e h^{-1}$. Thus $P_e \neq hP_e h^{-1}$. \square

From this, we conclude:

Corollary 3.12. *$E_{\text{lo}}(F)$ is not smooth.*

4. OPEN PROBLEMS

Theorem 3.2 shows that for a left-orderable group G , the property of $E_{\text{lo}}(G)$ being smooth transfers to relatively convex subgroups. It is natural to ask if the analogue is true for hyperfiniteness.

Question 4.1. *Is it true that if there is a relatively convex $C \leq G$ such that $E_{\text{lo}}(C)$ is not hyperfinite, then $E_{\text{lo}}(G)$ is not hyperfinite?*

The results of this paper and our previous work produce several techniques to handle the problem whether $E_{\text{lo}}(G)$ is smooth. The next degree of Borel complexity is that of hyperfinite equivalence relations that are not smooth. Since every Borel equivalence relation with countable classes is hyperfinite on a comeager set, it is generally very difficult to show that a given equivalence relation is hyperfinite. So we pose the following open questions, which are likely to require new techniques.

Question 4.2. Is $E_{\text{lo}}(\text{BS}(1, n))$ hyperfinite for $n > 2$?

Question 4.3. Is $E_{\text{lo}}(F)$ hyperfinite?

REFERENCES

- [CC22] Filippo Calderoni and Adam Clay. Borel structures on the space of left-orderings. *Bull. Lond. Math. Soc.*, 54:83–94, 2022.
- [CC23] Filippo Calderoni and Adam Clay. The Borel complexity of the space of left-orderings, low-dimensional topology, and dynamics. Preprint, available on ArXiv at <https://arxiv.org/pdf/2305.03927.pdf>, 2023.
- [CJM⁺ar] Clinton T. Conley, Steve Jackson, Andrew S. Marks, Brandon Seward, and Robin D. Tucker-Drob. Borel asymptotic dimension and hyperfinite equivalence relations. *Duke Mathematical Journal*, to appear.
- [Cla10] Adam Clay. Isolated points in the space of left orderings of a group. *Groups Geom. Dyn.*, 4(3):517–532, 2010.
- [DNR16] Bertrand Deroin, Andrés Navas, and Cristóbal Rivas. Groups, orders, and dynamics. Preprint, available on ArXiv at <https://arxiv.org/abs/1408.5805>, 2016.
- [Eff65] Edward G. Effros. Transformation groups and C^* -algebras. *Ann. of Math. (2)*, 81:38–55, 1965.
- [Gao09] Su Gao. *Invariant descriptive set theory*, volume 293 of *Pure and Applied Mathematics (Boca Raton)*. CRC Press, Boca Raton, FL, 2009.
- [Gli61] James Glimm. Locally compact transformation groups. *Transactions of the American Mathematical Society*, 101(1):124–138, 1961.
- [Kec95] Alexander S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Nav10] Andrés Navas. On the dynamics of (left) orderable groups. *Ann. Inst. Fourier (Grenoble)*, 60(5):1685–1740, 2010.
- [Osi21a] Denis Osin. Condensed groups in product varieties. *Journal of Group Theory*, 24(4):753–763, 2021.
- [Osi21b] Denis Osin. A topological zero-one law and elementary equivalence of finitely generated groups. *Annals of Pure and Applied Logic*, 172(3):102915, 2021.
- [Riv10] Cristóbal Rivas. On spaces of conradian group orderings. *Journal of Group Theory*, 13(3):337–353, 2010.
- [RT16] Cristóbal Rivas and Romain Tessera. On the space of left-orderings of virtually solvable groups. *Groups Geom. Dyn.*, 10(1):65–90, 2016.
- [Smi66] D. M. Smirnov. Right-ordered groups. *Algebra i Logika Sem.*, 5(6):41–59, 1966.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, HILL CENTER FOR THE MATHEMATICAL SCIENCES, 110 FRELINGHUYSEN RD., PISCATAWAY, NJ 08854-8019
 Email address: filippo.calderoni@rutgers.edu

DEPARTMENT OF MATHEMATICS, 420 MACHRAY HALL, UNIVERSITY OF MANITOBA, WINNIPEG, MB, R3T 2N2, CANADA
 Email address: Adam.Clay@umanitoba.ca