

(14) 1. Use the method of partial fractions to verify that  $\int_0^1 \frac{1}{(x+1)(x^2+1)} dx = \frac{1}{4} \ln 2 + \frac{1}{8} \pi$ .

**Answer** Write  $\frac{1}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+1} = \frac{A(x^2+1)+(Bx+C)(x+1)}{(x+1)(x^2+1)}$ , so  $1 = A(x^2+1) + (Bx+C)(x+1)$ . When  $x = -1$  we get  $A = \frac{1}{2}$ . Comparing  $x^2$  coefficients of both sides, we see  $B = -\frac{1}{2}$ . Finally, comparing constant coefficients on both sides, we see that  $1 = A + C$  so  $C = \frac{1}{2}$ . Compute:  $\int_0^1 \frac{1}{(x+1)(x^2+1)} dx = \int_0^1 \frac{\frac{1}{2}}{x+1} + \frac{-\frac{1}{2}x + \frac{1}{2}}{x^2+1} dx = \int_0^1 \frac{\frac{1}{2}}{x+1} + \frac{-\frac{1}{2}x}{x^2+1} + \frac{\frac{1}{2}}{x^2+1} dx = \frac{1}{2} \ln(x+1) - \frac{1}{4} \ln(x^2+1) + \frac{1}{2} \arctan(x) \Big|_0^1 = \frac{1}{2} \ln(2) - \frac{1}{4} \ln(2) + \frac{1}{2} \arctan(1) - \left( \frac{1}{2} \ln(1) - \frac{1}{4} \ln(1) + \frac{1}{2} \arctan(0) \right) = \frac{1}{4} \ln(2) + \frac{1}{2} \frac{\pi}{4} = \frac{1}{4} \ln 2 + \frac{\pi}{8}$ . Whew! All done.

(8) 2. Suppose  $\mathcal{R}$  is the region bounded by  $y = \frac{1}{x}$ ,  $x = 1$ ,  $x = 2$ , and  $y = 0$ .

a) Find the volume of the solid that results from rotating  $\mathcal{R}$  around the  $x$ -axis.      b) Find the volume of the solid that results from rotating  $\mathcal{R}$  around the  $y$ -axis.

**Answer**  $\pi \int_1^2 \pi \left(\frac{1}{x}\right)^2 dx = \pi \cdot \left(\frac{-1}{x}\right) \Big|_1^2 = \frac{\pi}{2}$ .


**Answer**  $2\pi \int_1^2 x \left(\frac{1}{x}\right) dx = 2\pi$ .

(12) 3. a) Carefully sketch  $y = \sin x$  and  $y = \sin 2x$  on the axes given for  $0 \leq x \leq \pi$ . Be sure to label each of the curves.

**Answer** The dashed curve is  $y = \sin 2x$  and the solid curve is  $y = \sin x$ .

b) Find **all** points of intersection of the curves sketched in part a).

**Answer** The sketch has three intersection points. Since  $\sin 2x = 2 \sin x \cos x$ , these occur when  $2 \sin x \cos x = \sin x$ : when  $\sin x = 0$ , which occurs  $x = 0$  or  $x = \pi$  in the sketch, or when  $\cos x = \frac{1}{2}$  which happens when  $x = \frac{\pi}{3}$ .



c) Find the area enclosed between the curves sketched in part a). Give an exact answer in terms of mathematical constants such as  $\pi$  and  $e$ .

**Answer** In  $[0, \frac{\pi}{3}]$ , the top curve is  $\sin 2x$ . The enclosed area there is  $\int_0^{\frac{\pi}{3}} \sin 2x - \sin x dx = -\frac{1}{2} \cos 2x + \cos x \Big|_0^{\frac{\pi}{3}} = \frac{1}{4}$ . In  $[\frac{\pi}{3}, \pi]$ , the top curve is  $\sin x$ . The enclosed area there is  $\int_{\frac{\pi}{3}}^{\pi} \sin x - \sin 2x dx = -\cos x + \frac{1}{2} \cos 2x \Big|_{\frac{\pi}{3}}^{\pi} = \frac{9}{4}$ . So the total area is  $\frac{1}{4} + \frac{9}{4} = \frac{5}{2}$ .

(10) 4. a) Suppose  $w$  is a positive number. Define  $A(w)$  to be the average value of  $(\cos x)^2$  on the interval  $0 \leq x \leq w$ . Compute  $A(w)$ , and show how the integral is calculated. **Answer**  $\int_0^w (\cos x)^2 dx = \int_0^w \frac{1}{2}(1 + \cos 2x) dx = \frac{1}{2}(x + \frac{1}{2} \sin 2x) \Big|_0^w = \frac{1}{2}(w + \frac{1}{2} \sin 2w) - \frac{1}{2}(0 + \sin 0)$ . The average value is the integral divided by the length of the interval, which is  $w$ , so  $A(w) = \frac{1}{2} + \frac{1}{4} \frac{\sin 2w}{w}$ .

b) What is the limit of  $A(w)$  as  $w \rightarrow 0^+$ ? **Answer** The limit of  $\frac{\sin 2w}{w}$  can be computed using L'Hospital's rule. It is  $\lim_{w \rightarrow 0^+} \frac{2 \cos 2w}{1} = 2$ . So the limit requested is  $\frac{1}{2} + \frac{1}{4} \cdot 2 = 1$ . The limit can also be computed by noticing that  $\sin 2w = 2 \sin w \cos w$ , and  $\lim_{w \rightarrow 0} \frac{\sin w}{w} = 1$ , familiar from earlier calculus study.

(12) 5. a) Explain why  $\int_1^{\infty} \frac{1}{x^2 + e^{2x}} dx$  converges. **Answer**  $\frac{1}{x^2 + e^{2x}}$  is positive on the interval  $[1, \infty)$ , and is certainly less than either  $\frac{1}{x^2}$  or  $\frac{1}{e^{2x}}$ . Either of those integrals converges on that interval, so by comparison, the integral given converges.

b) Explain why the value of the integral in a) is less than  $\frac{1}{2}$ . **Answer** In a) either integral is "good enough" to show convergence. Here, however, since  $\int_1^{\infty} \frac{1}{x^2} dx = \lim_{A \rightarrow \infty} -\frac{1}{x} \Big|_1^A = 1$  and  $\int_1^{\infty} \frac{1}{e^{2x}} dx = \lim_{A \rightarrow \infty} -\frac{1}{2} e^{-2x} \Big|_1^A = \frac{1}{2e^2}$  we'd better use the second, exponential integral (decaying exponentials  $\rightarrow 0$  much more rapidly than *any* inverse power of  $x$ ). So the value of the integral given is less than  $\frac{1}{2e^2}$  which is certainly less than  $\frac{1}{2}$ .

(12) 6. Calculate the following integrals, showing your work. a)  $\int \frac{dx}{\sqrt{1+x^2}}$  **Answer** If  $x = \tan \theta$ , then  $dx = (\sec \theta)^2 d\theta$  and  $\sqrt{1+x^2} = \sec \theta$ . So  $\int \frac{dx}{\sqrt{1+x^2}} = \int \sec \theta d\theta = \ln(\sec \theta + \tan \theta) + C = \ln(\sqrt{1+x^2} + x) + C$ .

b)  $\int \frac{dx}{2e^x + 1}$  **Answer** If  $u = e^x$ , then  $du = e^x dx$  so  $dx$  is  $\frac{du}{u}$ . The integral given becomes  $\int \frac{1}{u(2u+1)} du$ . Rewrite the integrand using partial fractions:  $\frac{1}{u(2u+1)} = \frac{A}{u} + \frac{B}{2u+1}$  giving  $1 = A(2u+1) + Bu$ . When  $u = 0$ , we get  $A = 1$ . When  $u = -\frac{1}{2}$  we get  $B = -2$ . Then antidifferentiating:  $\int \frac{1}{u} + \frac{-2}{2u+1} du = \ln u - \ln(2u+1) + C$ . Converting back to  $x$ 's, the answer becomes  $\ln(e^x) - \ln(2e^x + 1) + C$ . (Yes,  $\ln(e^x)$  is  $x$ .)

(10) 7. Use a substitution followed by integration by parts to verify that  $\int_0^1 e^{\sqrt{x}} dx = 2$ .

**Answer** Try  $w = \sqrt{x}$  and therefore  $w^2 = x$  with  $2w dw = dx$ . The integral changes:  $\int e^{\sqrt{x}} dx = \int e^w 2w dw$ . This integral is a well-known candidate for integration by parts. Let's do it (I'll save the 2 until later):

$$\int e^w w dw = we^w - \int e^w dw = we^w - e^w + C \quad \text{with} \quad \left. \begin{array}{l} u = w \\ dv = e^w dw \end{array} \right\} \left\{ \begin{array}{l} du = dw \\ v = e^w \end{array} \right. \quad \text{We can reverse the substitution so the indefinite integral becomes (with the 2!)} \quad 2 \left( \sqrt{x} e^{\sqrt{x}} - e^{\sqrt{x}} \right) + C.$$

Therefore,  $\int_0^1 e^{\sqrt{x}} dx = 2 \left( \sqrt{x} e^{\sqrt{x}} - e^{\sqrt{x}} \right) \Big|_0^1 = (2(e - e)) - (2(0 - 1)) = 2$ .

(12) 8. a) Suppose  $m$  and  $n$  are positive integers. Find a reduction formula for  $\int x^m (\ln x)^n dx$ . (Here the aim is to have an equation with the integral given here on one side and a similar integral with reduced  $n$  on the other side, since if we can push  $n$  to 0 we'll just have a polynomial to integrate, which is easy.)

**Answer** Again, integrate by parts.  $\int x^m (\ln x)^n dx = \left( \frac{x^{m+1}}{m+1} \right) \cdot (\ln x)^n - \int n (\ln x)^{n-1} \cdot \left( \frac{1}{x} \right) \left( \frac{x^{m+1}}{m+1} \right) dx + C$

$$\int u dv = uv - \int v du$$

with  $\left. \begin{array}{l} u = (\ln x)^n \\ dv = x^m dx \end{array} \right\} \left\{ \begin{array}{l} du = n(\ln x)^{n-1} dx \\ v = \frac{x^{m+1}}{m+1} \end{array} \right. \quad \text{I'll clean this up a bit since the formula will be used in part b):}$ 

$$\int x^m (\ln x)^n dx = \left( \frac{x^{m+1}}{m+1} \right) (\ln x)^n - \frac{n}{m+1} \int x^m (\ln x)^{n-1} dx + C.$$

b) Use the formula obtained in a) to compute  $\int x^{20} (\ln x)^2 dx$ .

**Answer** Since  $n = 2$  we need the formula above twice.

$$\int x^{20} (\ln x)^2 dx = \frac{x^{21}}{21} (\ln x)^2 - \frac{2}{21} \int x^{20} \ln x dx = \frac{x^{21}}{21} (\ln x)^2 - \frac{2}{21} \left( \frac{1}{21} x^{21} \ln x - \frac{1}{21} \int x^{20} dx \right) =$$

$$\frac{x^{21}}{21} (\ln x)^2 - \frac{2}{21} \left( \frac{1}{21} x^{21} \ln x - \frac{1}{21} \int x^{20} dx \right) = \frac{x^{21}}{21} (\ln x)^2 - \frac{2}{21} \left( \frac{1}{21} x^{21} \ln x - \frac{1}{21} \cdot \frac{x^{21}}{21} \right) + C$$

I doubt whether I would "simplify" anything here except under duress.

(10) 9. The integral  $\int_0^2 (x^3 + 1)^{7/2} dx$  is approximated using the Trapezoidal Rule by dividing  $[0, 2]$  into  $n$  segments of equal length. How large should  $n$  be in order to guarantee that the error is at most  $10^{-6}$ ?

**Note** You must give some reason explaining why any overestimates of derivatives you make are valid on the entire interval.

**Answer** We will need to overestimate  $|f''(x)|$  for all  $x$ 's in  $[0, 2]$  and for  $f(x) = (x^3 + 1)^{7/2}$ . So  $f'(x) = \frac{7}{2} (x^3 + 1)^{5/2} (3x^2)$  and  $f''(x) = \frac{7}{2} \cdot \frac{5}{2} \cdot (x^3 + 1)^{3/2} (3x^2)^2 + \frac{7}{2} (x^3 + 1)^{5/2} 6x$ . Notice that  $x^3 + 1$  is increasing in  $[0, 2]$  since its derivative,  $3x^2$ , is positive there. So the value of  $x^3 + 1$  for any  $x$  in  $[0, 2]$  is less than the value of  $x^3 + 1$  at the right-hand endpoint, which is 9. The other pieces of  $f''$  are also increasing and positive, so  $f''(2)$  will be an appropriate overestimate of  $|f''(x)|$  for all  $x$  in  $[0, 2]$ .

Now  $f''(2) = \frac{7}{2} \cdot \frac{5}{2} \cdot (2^3 + 1)^{3/2} (3 \cdot 2^2)^2 + \frac{7}{2} (2^3 + 1)^{5/2} (6 \cdot 2)$ . This is 44,226, which I will "plug into"  $M_2(b - a)^3 / 12n^2$  in place of  $M_2$ . Of course  $a = 0$  and  $b = 2$ , so the Trapezoid Rule error will be less than  $44,226 \cdot 8 / 12n^2$ . This is less than  $10^{-6}$  when  $n \geq 172,000$  (the approximate result my calculator reports). The number is large and, to me, rather unimpressive.

Detailed partial credit information is available on the web: look at the **Exam material** page, please.