

## Problems on Vector Geometry

This material corresponds roughly to sections 12.1, 12.2, 12.3 and 12.4 in the book, as well as the study guide Vector Geometry.

**Problem 1.** Find the vector  $\mathbf{v}$  of norm 3 that makes an angle of  $\frac{3\pi}{4}$  radians with the positive  $x$  axis.

The vector is

$$\mathbf{v} = 3 \left( \cos \left( \frac{3\pi}{4} \right), \sin \left( \frac{3\pi}{4} \right) \right) = 3 \left( -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right) \quad (1)$$

**Problem 2.** Determine whether the vectors  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$  are parallel, where  $A = (0, 2, 5)$ ,  $B = (3, 8, 2)$ ,  $C = (3, 5, -7)$  and  $D = (8, 15, -12)$

$$\begin{cases} \overrightarrow{AB} = B - A = (3, 6, -3) \\ \overrightarrow{CD} = D - C = (5, 10, -5) \end{cases} \quad (2)$$

If the vectors were parallel, one can find a real number such that

$$\overrightarrow{CD} = t\overrightarrow{AB} \quad (3)$$

In other words, we need to solve

$$\begin{cases} 5 = 3t \\ 10 = 6t \\ -5 = -3t \end{cases} \quad (4)$$

The solution of this system of equations is  $t = \frac{5}{3}$  so the vectors are parallel. Alternatively, one can observe that

$$\overrightarrow{CD} \times \overrightarrow{AB} = (0, 0, 0) \quad (5)$$

which as I mentioned in class is another way of verifying whether they are parallel or not.

**Problem 3.** Consider  $P = (1, 3)$ ,  $Q = (5, -1)$ ,  $R = (2, 3)$ ,  $S = (x, 2)$  where  $x$  is unknown.

1) Find  $\overrightarrow{PQ}$  and  $\overrightarrow{RS}$

2) Find the value of  $x$  so that  $\overrightarrow{PQ}$  and  $\overrightarrow{RS}$  become parallel.

3) Find a unit vector in the direction of  $\overrightarrow{RS}$

1)

$$\begin{cases} \overrightarrow{PQ} = Q - P = (4, -4) \\ \overrightarrow{RS} = S - R = (x - 2, -1) \end{cases} \quad (6)$$

2) we want to find a number  $t$  so that

$$(4, -4) = t(x - 2, -1) \quad (7)$$

This gives the equations

$$\begin{cases} 4 = t(x - 2) \\ -4 = -t \end{cases} \quad (8)$$

which has solution

$$x = 3 \quad (9)$$

3) the vector is

$$\hat{v} = \frac{\overrightarrow{RS}}{\|\overrightarrow{RS}\|} = \frac{(1, -1)}{\sqrt{2}} \quad (10)$$

**Problem 4. 1) Write the equation of a sphere with center  $P = (1, -1, 3)$  and radius 2.**

$$(x - 1)^2 + (y + 1)^2 + (z - 3)^2 = 4 \quad (11)$$

**2) Find the value (or values) of  $c$  such that  $Q = (2, -1, c)$  is on the sphere.**

We need  $Q$  to satisfy the previous equation, that is,

$$(2 - 1)^2 + (-1 + 1)^2 + (c - 3)^2 = 4 \quad (12)$$

which gives

$$c = 3 \pm \sqrt{3} \quad (13)$$

**Problem 5. Take  $P = (1, 1)$  and  $\overrightarrow{PQ} = \langle -2, 3 \rangle$ .**

1. Find point  $Q$

2. What is the length of  $\overrightarrow{PQ}$  ?

3. Find a unit vector parallel to  $\overrightarrow{PQ}$  .

1) We need

$$Q - P = \overrightarrow{PQ} = (-2, 3) \quad (14)$$

from which we see that

$$Q = P + (-2, 3) = (-1, 4) \quad (15)$$

2)  $\|\overrightarrow{PQ}\| = \sqrt{(-2)^2 + 3^2} = \sqrt{4 + 9} = \sqrt{13}$

3)  $\hat{v} = \frac{\overrightarrow{PQ}}{\|\overrightarrow{PQ}\|} = \frac{(-2, 3)}{\sqrt{13}}$

**Problem 6.** Suppose that  $\|\mathbf{v}\| = \sqrt{2}$  and  $\mathbf{u}$  is a vector such that the angle between  $\mathbf{u}$  and  $\mathbf{v}$  is  $\frac{\pi}{4}$  radians.

- 1) What must the value of  $\|\mathbf{u}\|$  so that  $\|\mathbf{u} + \mathbf{v}\| = \sqrt{5}$
- 2) Find the length of the orthogonal projection of  $\mathbf{u}$  onto the line going through  $\mathbf{v}$

1) Square both sides of the equation to obtain

$$\|\mathbf{u} + \mathbf{v}\|^2 = 5 \quad (16)$$

Now, notice that for any vector  $\mathbf{a}$   $\mathbf{a} \cdot \mathbf{a} = \|\mathbf{a}\|^2$  so taking  $\mathbf{a} = \mathbf{u} + \mathbf{v}$  the left hand side can be rewritten as

$$(\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} + 2\mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{v} = \|\mathbf{u}\|^2 + 2\|\mathbf{u}\|\|\mathbf{v}\|\cos\frac{\pi}{4} + \|\mathbf{v}\|^2 \quad (17)$$

By assumption  $\|\mathbf{v}\| = \sqrt{2}$  so we must solve the equation

$$\|\mathbf{u}\|^2 + 2\|\mathbf{u}\| + 2 = 5 \quad (18)$$

or

$$\|\mathbf{u}\|^2 + 2\|\mathbf{u}\| - 3 = 0 \quad (19)$$

whose only positive solution is

$$\|\mathbf{u}\| = 1 \quad (20)$$

2) Using the formula in the PDF (page 11) we need to find

$$\mathbf{u}_{\parallel} = \frac{(\mathbf{u} \cdot \mathbf{v})}{\|\mathbf{v}\|^2} \mathbf{v} = \frac{\|\mathbf{u}\|\|\mathbf{v}\|\cos\pi/4}{2} \mathbf{v} = \frac{1}{2} \mathbf{v} \quad (21)$$

so

$$\|\mathbf{u}_{\parallel}\| = \frac{1}{2}\|\mathbf{v}\| = \frac{\sqrt{2}}{2} \quad (22)$$

**Problem 7.** Consider the points  $P = (1, 2, 0)$ ,  $Q = (-1, 1, 1)$  and  $R = (2, 0, 1)$ .

a) Find the vectors  $\overrightarrow{PQ}$  and  $\overrightarrow{PR}$

$$\overrightarrow{PQ} = Q - P = (-2, -1, 1) \text{ and } \overrightarrow{PR} = R - P = (1, -2, 1)$$

b) Find the cross product  $\overrightarrow{PQ} \times (\overrightarrow{PR} + 3\overrightarrow{PQ})$

$$\overrightarrow{PQ} \times (\overrightarrow{PR} + 3\overrightarrow{PQ}) = \overrightarrow{PQ} \times \overrightarrow{PR} + 3\overrightarrow{PQ} \times \overrightarrow{PQ} = \overrightarrow{PQ} \times \overrightarrow{PR} = (1, 3, 5) \quad (23)$$

c) Find the area of the parallelogram whose sides are the vectors  $2\overrightarrow{PQ}$  and  $-3\overrightarrow{PR}$

The area is simply

$$\|2\overrightarrow{PQ} \times (-3\overrightarrow{PR})\| = 6\|\overrightarrow{PQ} \times \overrightarrow{PR}\| = 6\|(1, 3, 5)\| = 6 \cdot \sqrt{35} \quad (24)$$

**Problem 8.** a) Consider the points  $P = (1, 2, 1)$  and  $Q = (x, 4, -x)$ . Find the value (or values) of  $x$  such that  $\|\overrightarrow{PQ}\| = \sqrt{14}$ .

Squaring both sides we need to solve  $|\vec{PQ}|^2 = 14$  which is the same as

$$(x - 1)^2 + (4 - 2)^2 + (-x - 1)^2 = 14 \quad (25)$$

and this gives the solution  $x = \pm 2$ .

**b) Using the value (or values) found in a), find the vector projection of  $\vec{PQ}$  along the vector  $\mathbf{v} = (3, 0, 4)$ , that is, find  $\text{proj}_{\mathbf{v}} \vec{PQ}$**

We need to compute (page 11 PDF notes)

$$\frac{\vec{PQ} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} = \frac{(x - 1)3 + 0(2) + (-x - 1)4}{25} (3, 0, 4) = \frac{-x - 7}{25} (3, 0, 4) \quad (26)$$

depending on the value of  $x$ , we obtain  $\frac{-9}{25}(3, 0, 4)$  or  $\frac{-5}{25}(3, 0, 4)$ .