


Combinatorial Atlas for Log-concave Inequalities

Swee Hong Chan

joint with Igor Pak




What is log-concavity?

A sequence $a_1, \dots, a_n \in \mathbb{R}_{\geq 0}$ is **log-concave** if

$$a_k^2 \geq a_{k+1} a_{k-1} \quad (1 < k < n).$$

Log-concavity (and positivity) implies **unimodality**:

$a_1 \leq \dots \leq a_m \geq \dots \geq a_n$ for some $1 \leq m \leq n$.



Example: binomial coefficients

$$a_k = \binom{n}{k} \quad k = 0, 1, \dots, n.$$

This sequence is **log-concave** because

$$\frac{a_k^2}{a_{k+1} a_{k-1}} = \frac{\binom{n}{k}^2}{\binom{n}{k+1} \binom{n}{k-1}} = \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right),$$

which is greater than 1.

Example: permutations with k inversions


a_k = number of $\pi \in S_n$ with k inversions,

where **inversion** of π is pair $i < j$ s.t. $\pi_i > \pi_j$.

This sequence is **log-concave** because

$$\sum_{0 \leq k \leq \binom{n}{2}} a_k q^k = [n]_q! = \prod_{i=1}^{n-1} (1 + q + q^2 + \dots + q^i)$$

is a product of log-concave polynomials.



Log-concavity appears in different objects
for different reasons.

Today we focus on reason for **matroids**.

Warmup: graphs and forests

Let $G = (V, E)$ be a graph.

A (spanning) forest $F = (V, E')$ with $E' \subseteq E$ is a subset of edges without cycles.



G



forest



not forest



spanning tree

Log-concavity for forests

Theorem (Huh '15)

For every graph and $k \geq 1$,

$$I_k^2 \geq I_{k+1} I_{k-1},$$

where I_k is the number of forests with k edges.

Proof used [Hodge theory](#) from algebraic geometry.

In fact, [stronger](#) inequalities for [more general](#) objects are true.

Object: Matroids

Matroid $\mathcal{M} = (X, \mathcal{I})$ is ground set X with collection of independent sets $\mathcal{I} \subseteq 2^X$.

Graphical matroids


- X = edges of a graph G ,
- \mathcal{I} = forests in G .

Realizable matroids


- X = finite set of vectors over field \mathbb{F} ,
- \mathcal{I} = sets of linearly independent vectors.

Matroids: Conditions

- $S \subseteq T$ and $T \in \mathcal{I}$ implies $S \in \mathcal{I}$.



- If $S, T \in \mathcal{I}$ and $|S| < |T|$, then there is $x \in T \setminus S$ such that $S \cup \{x\} \in \mathcal{I}$.



Note: These are natural properties of sets of linearly independent vectors.

Mason's Conjecture (1972)

For every matroid and $k \geq 1$,

$$(1) \quad I_k^2 \geq I_{k+1} I_{k-1};$$

$$(2) \quad I_k^2 \geq \left(1 + \frac{1}{k}\right) I_{k+1} I_{k-1};$$

$$(3) \quad I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1}.$$

I_k is number of ind. sets of size k , and $n = |X|$.

Note: (3) \Rightarrow (2) \Rightarrow (1).

Why $(1 + \frac{1}{k}) (1 + \frac{1}{n-k})$?

Mason (3) is equivalent to **ultra/binomial log-concavity**,

$$\frac{I_k^2}{\binom{n}{k}^2} \geq \frac{I_{k+1}}{\binom{n}{k+1}} \frac{I_{k-1}}{\binom{n}{k-1}}.$$

Equality occurs **if** every $(k + 1)$ -subset is independent.

Solution to Mason (1)

Theorem (Adiprasito-Huh-Katz '18)

For every matroid and $k \geq 1$,

$$I_k^2 \geq I_{k+1} I_{k-1}.$$

Proof used [combinatorial Hodge theory](#) for
matroids.

Solution to Mason (2)

Theorem (Huh-Schröter-Wang '18)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) I_{k+1} I_{k-1}.$$

Proof used [combinatorial Hodge theory](#) for [correlation inequality](#) on matroids.

Solution to Mason (3)

Theorem

(Anari-Liu-Oveis Gharan-Vinzant, Brändén-Huh '20)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1}.$$

Proof used theory of strong log-concave polynomials /
Lorentzian polynomials.

Solution to Mason (3)

Theorem

(Anari-Liu-Oveis Gharan-Vinzant, Brändén-Huh '20)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1}.$$

Theorem (Murai-Nagaoka-Yazawa '21)

Equality occurs if and only if every $(k+1)$ -subset is independent.

Our contribution

Method: Combinatorial atlas

Results: Log-concave inequalities, and
if and only if conditions for equality

- Matroids (refined);
- Morphism of matroids (refined);
- Discrete polymatroids;
- Stanley's poset inequality (refined);
- Poset antimatroids;
- Branching greedoid (log-convex);
- Interval greedoids.

Method: Combinatorial atlas

Results: Log-concave inequalities, and
if and only if conditions for equality

- **Matroids (refined);**
- Morphism of matroids (refined);
- Discrete polymatroids;
- Stanley's poset inequality (refined);
- Poset antimatroids;
- Branching greedoid (log-convex);
- Interval greedoids.

**Combinatorial atlas application:
Matroids**

Warmup: graphical matroids refinement

Corollary (C.-Pak)

For graphical matroid of simple connected graph

$G = (V, E)$, and $k = |V| - 2$,

$$(I_k)^2 \geq \frac{3}{2} \left(1 + \frac{1}{k}\right) I_{k+1} I_{k-1},$$

with equality *if and only if* G is cycle graph.

Numerically *better* than Mason (3), because

$$\frac{3}{2} \geq 1 + \frac{1}{n-k} = 1 + \frac{1}{|E| - |V| + 2}$$

for G that is not tree.

Comparison with Mason (3)

Our bound gives

$$\frac{(I_k)^2}{I_{k+1} I_{k-1}} \geq \frac{3}{2} \quad \text{when } |E| - |V| \rightarrow \infty,$$

Meanwhile, Mason (3) bound only gives

$$\frac{(I_k)^2}{I_{k+1} I_{k-1}} \geq 1 \quad \text{when } |E| - |V| \rightarrow \infty.$$

Our bound is **better** numerically and asymptotically.

Refinement for Mason (3)

Theorem 1 (C.-Pak)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{\text{prl}_{\mathcal{M}}(k-1) - 1}\right) I_{k+1} I_{k-1}.$$

This refines Mason (3),

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1},$$

since

$$\text{prl}_{\mathcal{M}}(k-1) \leq n - k + 1.$$

Refinement for different matroids

- For all matroids,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1}.$$

- Graphical matroids and $k = |V| - 2$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \frac{3}{2} I_{k+1} I_{k-1}.$$

- Realizable matroids over \mathbb{F}_q ,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{q^{m-k+1}-2}\right) I_{k+1} I_{k-1}.$$

- (k, m, n) -Steiner system matroid,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \frac{n-k+1}{n-m} I_{k+1} I_{k-1}.$$

Refinement for Mason (3)

Theorem 2 (C.-Pak)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{\text{prl}_{\mathcal{M}}(k-1) - 1}\right) I_{k+1} I_{k-1}.$$

This refines Mason (3),

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1},$$

since

$$\text{prl}_{\mathcal{M}}(k-1) \leq n - k + 1.$$


Parallel classes of matroid \mathcal{M}

Loop is $x \in X$ such that $\{x\} \notin \mathcal{I}$.

Non-loops x, y are **parallel** if $\{x, y\} \notin \mathcal{I}$.

Parallelship equiv. relation: $x \sim y$ if $\{x, y\} \notin \mathcal{I}$.



Parallel class = equivalence class of \sim .



Matroid contraction

Contraction of $S \in \mathcal{I}$ is matroid \mathcal{M}_S with

$$X_S = X \setminus S, \quad \mathcal{I}_S = \{T \setminus S : S \subseteq T\}.$$




$\text{prl}(S) :=$ number of parallel classes of \mathcal{M}_S


Parallel number

The k -parallel number is


$$\text{prl}_{\mathcal{M}}(k) := \max\{\text{prl}(S) \mid S \in \mathcal{I} \text{ with } |S| = k\}.$$




$$\text{prl}(S) = 2$$



$$\text{prl}(S) = 2$$



$$\text{prl}(S) = 2$$



$$\text{prl}(S) = 3$$

$$\text{prl}_{\mathcal{M}}(1) = 3$$

Refinement for Mason (3)

Theorem 3 (C.-Pak)

For every matroid and $k \geq 1$,

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{\text{prl}_{\mathcal{M}}(k-1) - 1}\right) I_{k+1} I_{k-1}.$$

This refines Mason (3),

$$I_k^2 \geq \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right) I_{k+1} I_{k-1},$$

since

$$\text{prl}_{\mathcal{M}}(k-1) \leq n - k + 1.$$

When is equality achieved?

- When every $(k + 1)$ -subset is independent,
 $\text{prl}_{\mathcal{M}}(k - 1) = n - k + 1$.
- Graphical matroid when G is a cycle,
 $\text{prl}_{\mathcal{M}}(k - 1) = 3$.
- Realizable matroids of every m -vectors over \mathbb{F}_q ,
 $\text{prl}_{\mathcal{M}}(k - 1) = q^{m-k+1} - 1$.
- (k, m, n) -Steiner system matroid,
 $\text{prl}_{\mathcal{M}}(k - 1) = \frac{n - k + 1}{m - k + 1}$.

Equality conditions

Theorem 4 (C.-Pak)

For every matroid and $k \geq 1$,

$$I_k^2 = \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{\text{prl}_{\mathcal{M}}(k-1) - 1}\right) I_{k+1} I_{k-1}$$

if and only if

for every $S \in \mathcal{I}$ with $|S| = k - 1$,

- \mathcal{M}_S has $\text{prl}_{\mathcal{M}}(k - 1)$ parallel classes; and
- Every parallel class of \mathcal{M}_S has same size.


Combinatorial atlas: the method

Combinatorial atlas



Input: Acyclic digraph \mathcal{A} , where each vertex v is associated with

- Symmetric matrix M with nonnegative entries;
- Vector g, h with nonnegative entries.

Atlas: example




Atlas: example (zoomed in)



Atlas example: matroid (simplified)

For matroid with $X = \{a, b, c\}$, the atlas for $k = 2$ is



Atlas example: matroid (simplified)

The matrix for the top vertex is

$$M_{a,b} = (k + 1)! \times \text{number of independent sets} \\ \text{of size } k + 1 \text{ containing } a, b$$

$$M_{a,*} = k! \times \text{number of independent sets} \\ \text{of size } k \text{ containing } a$$

$$M_{*,*} = (k - 1)! \times \text{number of independent sets} \\ \text{of size } k - 1$$

Combinatorial atlas

Input: Acyclic digraph \mathcal{A} , where each vertex v is associated with

- Symmetric matrix M with nonnegative entries;
- Vector g, h with nonnegative entries.

Goal: Show every M has [hyperbolic inequality](#).

Hyperbolic inequality

M has **hyperbolic inequality** property if

$$\langle \mathbf{x}, M\mathbf{y} \rangle^2 \geq \langle \mathbf{x}, M\mathbf{x} \rangle \langle \mathbf{y}, M\mathbf{y} \rangle,$$

for every $\mathbf{x} \in \mathbb{R}^r$, $\mathbf{y} \in \mathbb{R}_{\geq 0}^r$.

This condition is equivalent to

M has at most one positive eigenvalue.

Note: Already known to be important in **Lorentzian polynomials** and **Bochner's method proof of Aleksandrov-Fenchel inequality**.

How to get log-concave inequalities?

Assume a_{k-1}, a_k, a_{k+1} can be computed by

$$a_k = \langle \mathbf{g}, \mathbf{M}\mathbf{h} \rangle, \quad a_{k+1} = \langle \mathbf{g}, \mathbf{M}\mathbf{g} \rangle, \quad a_{k-1} = \langle \mathbf{h}, \mathbf{M}\mathbf{h} \rangle,$$

for $\mathbf{M}, \mathbf{g}, \mathbf{h}$ from a top vertex of the atlas.

$$\langle \mathbf{g}, \mathbf{M}\mathbf{h} \rangle^2 \geq \langle \mathbf{g}, \mathbf{M}\mathbf{g} \rangle \langle \mathbf{h}, \mathbf{M}\mathbf{h} \rangle \quad (\text{hyperbolic ineq.})$$

then implies

$$a_k^2 \geq a_{k+1} a_{k-1} \quad (\text{log-concave ineq.})$$

Combinatorial atlas

Input: Acyclic digraph \mathcal{A} , where each vertex v is associated with

- Symmetric matrix M with nonnegative entries;
- Vector g, h with nonnegative entries.

Goal: Show every M has hyperbolic inequality.

Method: Verify three conditions:

- **Irreducibility condition;**
- **Inheritance condition;**
- **Subdivergence condition.**

Irreducibility condition

- Matrix M associated to v is **irreducible** when restricted to its support;
- Vector h is associated to v is a **positive** vector.

For **matroids**, this means that the **base exchange graph** is connected.

This is a consequence of the **exchange property**.

Inheritance condition

Edge $e = (v, v_i)$ of v is associated with linear map $T_i : \mathbb{R}^r \rightarrow \mathbb{R}^r$ such that, for every $\mathbf{x} \in \mathbb{R}^r$,

$$i\text{-th coordinate of } \mathbf{M}\mathbf{x} = \langle T_i\mathbf{x}, \mathbf{M}_i T_i\mathbf{h} \rangle,$$

where \mathbf{M} and \mathbf{h} are associated to v , and \mathbf{M}_i is associated to v_i .

For **matroids** with $X = \{e_1, \dots, e_n\}$, this means

$$\begin{aligned} & k \times \text{number of independent } k\text{-sets} \\ &= \sum_{i=1}^n \text{number of independent } k\text{-sets containing } e_i. \end{aligned}$$

Subdivergence condition

For every $\mathbf{x} \in \mathbb{R}^r$,

$$\sum_{i=1}^r h_i \langle T_i \mathbf{x}, \mathbf{M}_i T_i \mathbf{x} \rangle \geq \langle \mathbf{x}, \mathbf{M} \mathbf{x} \rangle,$$

where $h_i = i$ -th coordinate of \mathbf{h} .

Note: Equality occurs for Lorentzian polynomials
and for matroids.

For matroids, this is consequence of hereditary property.

Bottom-to-top principle for hyperbolic inequalities


Proposition

Assume *irreducibility, inheritance, subdivergence*.


If every child vertex has *hyperbolic inequality property*, then so does the parent vertex.

Bottom-to-top principle **reduces Goal** to checking hyperbolic inequality only for **sink vertices**.


Bottom-to-top principle




Bottom-to-top principle



Bottom-to-top principle



Bottom-to-top principle



How about equalities?

Combinatorial atlas equality

Input:

- An acyclic digraph $\mathcal{A} := (\mathcal{V}, \mathcal{E})$ satisfying **previous** conditions;
- Vectors $\mathbf{g}, \mathbf{h} \in \mathbb{R}_{\geq 0}$;

Goal: Show “every” M has **hyperbolic equality**,

$$\langle \mathbf{g}, M\mathbf{h} \rangle^2 = \langle \mathbf{g}, M\mathbf{g} \rangle \langle \mathbf{h}, M\mathbf{h} \rangle.$$


Top-to-bottom principle for equalities

Proposition


*Assume **regularity** condition. If parent vertex has hyperbolic equality property, then so do children vertices.*

Top-to-bottom principle **expands** hyperbolic equality to **sink vertices**, and gives **combinatorial characterizations**.


Top-to-bottom principle




Top-to-bottom principle



Top-to-bottom principle



Top-to-bottom principle



Other applications

Full version: 2110.10740 (71 pages)

Expository version: 2203.01533 (28 pages)

Results: Log-concave inequalities and equalities for

- [Matroids \(refined\)](#);
- Morphism of matroids (refined);
- Discrete polymatroids;
- Stanley's poset inequality (refined);
- Poset antimatroids;
- Branching greedoid (log-convex);
- Interval greedoids.

THANK YOU!

Preprint: www.arxiv.org/abs/2110.10740

www.arxiv.org/abs/2203.01533

Webpage: www.math.rutgers.edu/~sc2518/

Email: sc2518@rutgers.edu

What is next?

Log-concavity for chromatic polynomials

Theorem (Huh '12)

For every *graph* G and $k \geq 1$,

$$C_k^2 \geq C_{k+1} C_{k-1},$$

where C_0, C_1, \dots are absolute coefficients of the *chromatic polynomial* of G .

Comparison to Mason (1):

- $(I_k)_{k \geq 0}$ is f-vector of *independence complex*;
- $(C_k)_{k \geq 0}$ is f-vector of *broken circuit complex*.

Stronger log-concavity for chromatic polynomials

Conjecture (Brylawski '82)


For every connected graph $G = (V, E)$ and $k \geq 1$,

$$C_k^2 \geq \left(1 + \frac{1}{|V| - k}\right) \left(1 + \frac{1}{|E| - |V| + k}\right) C_{k+1} C_{k-1},$$


Note: Brylawski conjectured the inequality for
characteristic polynomial of all matroids.

Atlas example: matroid (simplified)

Consider the graphical matroid for



The corresponding combinatorial atlas is



Atlas example: matroid (simplified)

	a	b	c	d	null	
	0	$\frac{3}{2} \times 1$	$\frac{3}{2} \times 1$	$\frac{3}{2} \times 2$	3	a
$\frac{3}{2} \times 1$	0	$\frac{3}{2} \times 1$	$\frac{3}{2} \times 1$	$\frac{3}{2} \times 2$	3	b
$\frac{3}{2} \times 1$	$\frac{3}{2} \times 1$	$\frac{3}{2} \times 1$	0	$\frac{3}{2} \times 2$	3	c
$\frac{3}{2} \times 2$	$\frac{3}{2} \times 2$	$\frac{3}{2} \times 2$	$\frac{3}{2} \times 2$	0	3	d
3	3	3	3	3	4	null

$$M_{a,b} = \frac{3}{2} \times \text{numbers of 3-forests containing } a, b$$

$$M_{a,\text{null}} = \text{number of 2-forests containing } a$$


$$M_{\text{null},\text{null}} = \text{number of 1-forests}$$

Here $\frac{3}{2}$ is the contribution from $1 + \frac{1}{\text{prl}_{\mathcal{M}}(k-1)-1}$.

**Combinatorial atlas application:
Stanley's poset inequality**



Partially ordered sets

A poset P is a set X with a partial order \prec on X .



Linear extension

A linear extension L is a complete order of \prec .



We write $L(x) = k$ if x is k -th smallest in L .

Stanley's inequality

Fix $z \in P$.

N_k is number of linear extensions with $L(z) = k$.

Theorem (Stanley '81)

For every poset and $k \geq 1$,

$$N_k^2 \geq N_{k+1} N_{k-1}.$$

Proof used [Aleksandrov-Fenchel inequality](#) for mixed volumes.

When is equality achieved?

Theorem (Shenfeld-van Handel)

Suppose $N_k > 0$. Then

$$N_k^2 = N_{k+1} N_{k-1}$$

if and only if

$$N_k = N_{k+1} = N_{k-1}.$$

Proof used classifications of extremals of Aleksandrov-Fenchel inequality for convex polytopes.

Our contribution

Open Problem (Folklore)

Give a *combinatorial* proof to Stanley's inequality.

Answer (C.–Pak)

We give new *combinatorial proof* for Stanley's ineq. and extend to *weighted version*.




Order-reversing weight

A weight $w : X \rightarrow \mathbb{R}_{>0}$ is **order-reversing** if

$$w(x) \geq w(y) \quad \text{whenever} \quad x \prec y.$$

Weight of linear extension L is

$$w(L) := \prod_{L(x) < L(z)} w(x).$$



Weighted Stanley's inequality

Fix $z \in P$.

$N_{w,k}$ is w -weight of linear extensions with $L(z) = k$.

Theorem 5 (C. Pak)

For every poset and $k \geq 1$,

$$N_{w,k}^2 \geq N_{w,k+1} N_{w,k-1}.$$

When is equality achieved?

Theorem 6 (C.-Pak)

Suppose $N_{w,k} > 0$. Then

$$N_{w,k}^2 = N_{w,k+1} N_{w,k-1}$$

if and only if

for every linear extension L with $L(z) = k$,

$$w(L^{-1}(k+1)) = w(L^{-1}(k-1)) =: s,$$


and

$$\frac{N_{w,k}}{s^k} = \frac{N_{w,k+1}}{s^{k+1}} = \frac{N_{w,k-1}}{s^{k-1}}.$$

**Combinatorial atlas application:
Poset antimatroids**

Feasible words of a poset

A word $\alpha \in X^*$ is **feasible** if no repeating elements, and y occurs in α and $x \prec y \Rightarrow x$ occurs in α before y .




Feasible: \emptyset , a , ab , ac , abc , acb , $abcd$, $acbd$.

Not feasible: aa , bc , ba .

Chain weight

For $x \in P$, chain weight is

$\omega(x)$ = number of maximal chains that starts with x .





$$\omega(a) = 2$$

$$\omega(b) = 1$$

$$\omega(c) = 1$$

$$\omega(d) = 1$$



Weight of word α is $\omega(\alpha) := \omega(\alpha_1) \dots \omega(\alpha_\ell)$.

Log-concave inequality for poset antimatroids

$F_{\omega,k}$ is sum of ω -weight of feasible words of length k .

Theorem 7 (C.-Pak)

For every poset and $k \geq 1$,

$$F_{\omega,k}^2 \geq F_{\omega,k+1} F_{\omega,k-1}.$$

When is equality achieved?

Theorem 8 (C.-Pak)

*Equality occurs for $k = 1, \dots, \text{height}(P) - 1$
if and only if*

*Hasse diagram of P is a forest where every leaf is of
the same level.*

