


Complexity of Log-concave Inequalities for Matroids

Swee Hong Chan

joint with Igor Pak




What is log-concavity?

A sequence $a_1, \dots, a_n \in \mathbb{R}_{\geq 0}$ is **log-concave** if

$$a_k^2 \geq a_{k+1} a_{k-1} \quad (1 < k < n).$$

Log-concavity (and positivity) implies **unimodality**:

$a_1 \leq \dots \leq a_m \geq \dots \geq a_n$ for some $1 \leq m \leq n$.



Example 1: binomial coefficients

$$a_k = \binom{n}{k} \quad k = 0, 1, \dots, n.$$

This sequence is **log-concave** because

$$\frac{a_k^2}{a_{k+1} a_{k-1}} = \frac{\binom{n}{k}^2}{\binom{n}{k+1} \binom{n}{k-1}} = \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right),$$

which is greater than 1.

Example 2: permutation inversion sequence

Let

$a_k :=$ number of $\pi \in S_n$ with k inversions,

where **inversion** of π is pair $i < j$ s.t. $\pi_i > \pi_j$.

This sequence is **log-concave** because

$$\sum_{0 \leq k \leq \binom{n}{2}} a_k q^k = [n]_q! = \prod_{i=1}^{n-1} (1 + q + q^2 + \dots + q^i)$$

is a product of log-concave polynomials.

Example 3: Mason's conjecture for matroids

Let \mathcal{M} be a **matroid**, and

$a_k :=$ number of **independent sets** with k elements.

Log-concavity was conjectured for all **matroids** (Mason '72), and was proved using **combinatorial Hodge theory** (Adiprasito–Huh–Katz '18).

Motivation

Which log-concave inequality is more “difficult”?

3 is more
difficult!




1,2,3 are
equally easy!



Motivation


Which log-concave inequality is more “difficult”?



We will show that [REDACTED] (3)
is **strictly more** difficult than the rest, using
Complexity Theory.

Motivation

Which log-concave inequality is more “difficult”?



We will show that a slight generalization of (3) is strictly more difficult than the rest, using Complexity Theory.

Stanley–Yan inequality

Stanley–Yan inequality (simple case)

Let \mathcal{M} be a matroid with ground set X and rank r .

Fix a subset S of X . Let

$B(k) :=$ no. of **bases** B such that $|B \cap S| = k$,
multiplied by $r! \times \binom{r}{k}^{-1}$.

Theorem (Stanley '81, Yan '23)

The sequence $B(1), B(2), \dots$ is log-concave,

$$B(k)^2 \geq B(k+1)B(k-1) \quad (k \in \mathbb{N}).$$

Stanley–Yan inequality (simple)

Theorem (Stanley '81, Yan '23)

$$B(k)^2 \geq B(k+1)B(k-1) \quad (k \in \mathbb{N}).$$

Proved for **regular** matroids by (Stanley '81) using **Alexandrov–Fenchel inequality** for mixed volumes.

Proved for **all** matroids by (Yan '23) using theory of **Lorentzian polynomials**.

Proof of Mason's conjecture using SY inequality

Let

$\mathcal{M}' :=$ direct sum of \mathcal{M} with the
free matroid of r elements;

$S :=$ ground set of \mathcal{M} .

Then

$$I(k) \text{ for } \mathcal{M} = \frac{1}{r!} \times B(k) \text{ for } \mathcal{M}',$$

where $I(k)$ is no. of independent sets with k elts.

Thus [Stanley–Yan](#) inequality for \mathcal{M}'
implies [Mason's conjecture](#) for \mathcal{M} .

Stanley–Yan inequality (full version)

Fix $d \geq 0$, disjoint subsets S, S_1, \dots, S_d of X ,
and $\ell_1, \dots, \ell_d \in \mathbb{N}$.

$B_d(k) :=$ number of bases B of \mathcal{M} such that
 $|B \cap S| = k, |B \cap S_i| = \ell_i$ for $i \in [d]$,
multiplied by $r! \times \binom{r}{k, \ell_1, \dots, \ell_d}^{-1}$.

Theorem (Stanley '81, Yan '23)

The sequence $B_d(1), B_d(2), \dots$ is log-concave,

$$B_d(k)^2 \geq B_d(k+1)B_d(k-1) \quad (k \in \mathbb{N}).$$

What we want to do

Theorem (Stanley '81, Yan '23)

The sequence $B_d(1), B_d(2), \dots$ is log-concave,

$$B_d(k)^2 \geq B_d(k+1)B_d(k-1) \quad (k \in \mathbb{N}).$$

Both LHS and RHS of this inequality has
combinatorial interpretations.


But we will show that this inequality has
no combinatorial injective proof.

Combinatorial injective proof

Combinatorial injection

An injection $f : A \rightarrow B$ is **combinatorial** if

- Given $x \in A$, the image $f(x)$ is computable in $\text{poly}(|x|)$ steps;
- Given $y \in B$, it takes $\text{poly}(|y|)$ steps **to decide if y is in image of f** ; and if so, the pre-image $f^{-1}(y)$ is computable in $\text{poly}(|y|)$ steps.




Example: Injective proof of binomial inequality

$$\binom{n}{k}^2 \geq \binom{n}{k+1} \binom{n}{k-1} \quad (1 < k < n).$$

This inequality has a **lattice path interpretation**:

$K(a \rightarrow c, b \rightarrow d) :=$ no. of pairs of north-east lattice paths from a to c and b to d ,

for $a, b, c, d \in \mathbb{Z}^2$.



Example: Injective proof of binomial inequality

Let



$$a = (0, 1), \quad c = (k, n - k + 1),$$

$$b = (1, 0), \quad d = (k + 1, n - k).$$

Then


$$K(a \rightarrow c, b \rightarrow d) = \binom{n}{k},$$

$$K(a \rightarrow d, b \rightarrow c) = \binom{n}{k-1} \binom{n}{k+1}.$$



Example: Injective proof of binomial inequality

$f : K(a \rightarrow d, b \rightarrow c) \rightarrow K(a \rightarrow c, b \rightarrow d)$
is defined by **path-swapping injections**.



Images of f are pairs of lattice paths that **intersects**.

First main result

Theorem 1 (C.–Pak '24+)

There is *no combinatorial injective proof* for Stanley–Yan inequality, assuming $\text{NP}^{\text{NP}} \neq \text{coNP}^{\text{NP}}$.

The *assumption* above is slightly stronger than $\text{P} \neq \text{NP}$, and is widely used in Complexity Theory.

First main result

Theorem 1 (C.–Pak '24+)

There is *no combinatorial injective proof* for Stanley–Yan inequality, assuming $\text{NP}^{\text{NP}} \neq \text{coNP}^{\text{NP}}$.

This result is a consequence of Stanley–Yan inequality being *not in $\#P$* (explained next slide).

Complexity class $\#P$

Complexity class $\#P$

Informal definition for intuition:

Problems of counting the number
 $\#P :=$ of objects satisfying some **property**;
this **property** is simple to **verify**.

Example (Problem in $\#P$)

*Count number of **proper** 3-colorings of graph G .*

Complexity class NP

Problems asking about **existence** of
NP := a solution S for input x , where validity
of S can be verified in polynomial time.

Example (Problem in NP)

Does graph G have a proper 3-coloring?

Complexity class #P: Formal definition

Problems asking for **number** of solutions

$\#P$:= S for input x , where validity of S can be verified in polynomial time.

Example (Problem in #P)

Count the number of proper 3-colorings of graph G .

It might take **exponential time**
to solve a problem in #P.

Second main result

Theorem 2 (C.–Pak '24+)


Let \mathcal{M} be a *binary* matroid. Then the *defect* of Stanley–Yan inequality

$$B_d(k)^2 - B_d(k+1) B_d(k-1)$$

is *not in* $\#P$, assuming $NP^{NP} \neq coNP^{NP}$.

This means LHS and RHS of Stanley–Yan inequality belongs to $\#P$, but their difference *does not*.

Recall our goal




We will now show that **Stanley–Yan inequality** is strictly more difficult than the **binomial inequality** and **permutation inversion inequality**.

Example 1: Binomial inequality

It follows from **path-swapping injections** that

$\binom{n}{k}^2 - \binom{n}{k+1} \binom{n}{k-1} =$ number of **non-intersecting lattice paths** from a to c and b to d .



Thus the defect of this inequality belongs to $\#P$.

Example 2: Permutation inversion inequality

Let a_k = number of $\pi \in S_n$ with k inversions.

$$\text{Then } \sum_{0 \leq k \leq \binom{n}{2}} a_k q^k = \prod_{i=1}^{n-1} (1 + q + \dots + q^i)$$

is computable in $\text{poly}(n)$ time.

Thus $a_k^2 - a_{k+1}a_{k-1}$ is computable in $\text{poly}(n)$ time;

and thus belongs to $\#P$.

Conclusion

We compare three log-concave inequalities:

Binomial inequality: in $\#P$;

Permutation inversion inequality: in $\#P$;

Stanley–Yan inequality: not in $\#P$.

This differentiates Stanley–Yan inequality from binomial inequality and permutation inversion inequality.

감사합니다!

THANK YOU!

Preprint: www.arxiv.org/abs/2407.19608

Webpage: www.math.rutgers.edu/~sc2518/

Email: sweehong.chan@rutgers.edu