


# Complexity of Log-concave Inequalities for Matroids

Swee Hong Chan

joint with Igor Pak




## What is log-concavity?

A sequence  $a_1, \dots, a_n \in \mathbb{N}_{\geq 0}$  is **log-concave** if

$$a_k^2 \geq a_{k+1} a_{k-1} \quad (1 < k < n).$$

Log-concavity (and positivity) implies **unimodality**:

$a_1 \leq \dots \leq a_m \geq \dots \geq a_n$  for some  $1 \leq m \leq n$ .



## Log-concave shaped objects in real life



Esplanade – Theatres on the Bay,  
Singapore (Credit TyLin).

## Example 1: Binomial coefficients

$$a_k = \binom{n}{k} \quad k = 0, 1, \dots, n.$$

This sequence is **log-concave** because

$$\frac{a_k^2}{a_{k+1} a_{k-1}} = \frac{\binom{n}{k}^2}{\binom{n}{k+1} \binom{n}{k-1}} = \left(1 + \frac{1}{k}\right) \left(1 + \frac{1}{n-k}\right),$$

which is greater than 1.

## Example 2: Permutation inversion sequence

Let

$a_k :=$  number of  $\pi \in S_n$  with  $k$  inversions,

where **inversion** of  $\pi$  is pair  $i < j$  s.t.  $\pi_i > \pi_j$ .

This sequence is **log-concave** because

$$\sum_{0 \leq k \leq \binom{n}{2}} a_k q^k = [n]_q! = \prod_{i=1}^{n-1} (1 + q + q^2 + \dots + q^i)$$


is a product of log-concave polynomials.

### Example 3: Forests of a graph


$a_k$  = number of forests with  $k$  edges of graph  $G$ .

**Forest** is a subset of edges of  $G$  that has no cycles.


**Log-concavity** was conjectured for all **matroids** (Mason '72), and was proved through **combinatorial Hodge theory** (Huh '15).




$G$



forest



not forest




spanning tree

## Example 3: Forests of a graph

$a_k$  = number of forests with  $k$  edges of graph  $G$ .

**Forest** is a subset of edges of  $G$  that has no cycles.

**Log-concavity** was conjectured for all **matroids** (Mason '72), and was proved through **combinatorial Hodge theory** (Huh '15).



June Huh



Fields Medal

# Motivation

Which log-concave inequality is more “difficult”?

3 is more  
difficult!

Swee  
Hong




1,2,3 are  
equally easy!

Devil's  
advocate



# Motivation


Which log-concave inequality is more “difficult”?



We will show that [REDACTED] (3)  
is **strictly more** difficult than the rest,  
using **Complexity Theory**.

# Motivation

Which log-concave inequality is more “difficult”?



We will show that a **generalization** of (3) is **strictly more** difficult than the rest, using **Complexity Theory**.

# Matroids

## Object: Matroids

Matroid  $\mathcal{M} = (X, \mathcal{I})$  is ground set  $X$  with collection of independent sets  $\mathcal{I} \subseteq 2^X$ .

### Graphic matroids


- $X$  = edges of a graph  $G$ ,
- $\mathcal{I}$  = forests in  $G$ .

### Binary matroids


- $X$  = set of vectors over finite field  $\mathbb{F}_2$ ,
- $\mathcal{I}$  = sets of linearly independent vectors.

## Matroids: Axioms

- (Hereditary) If  $S \subseteq T$  and  $T \in \mathcal{I}$ , then  $S \in \mathcal{I}$ .




- (Exchange) If  $S, T \in \mathcal{I}$  and  $|S| < |T|$ , then there is  $x \in T \setminus S$  such that  $S \cup \{x\} \in \mathcal{I}$ .




## Matroid: Bases and ranks

A **basis** of  $\mathcal{M}$  is a **maximal** independent set.


**Rank**  $r$  of  $\mathcal{M}$  is the size of the bases.




$G$



Basis 1



Basis 2



Not Basis

---

Matroid generalizes the notion of **vector spaces**.

## **Mason's conjecture**

## First Mason's conjecture

For matroid  $\mathcal{M}$ , let

$I(k) :=$  no. of **independents sets** with  $k$  elements.

For **graphic** matroid,  $I(k)$  is no. of **forest** with  $k$  edges.

### Conjecture (Mason '72)

*The sequence  $I(1), I(2), \dots$  is log-concave,*

$$I(k)^2 \geq I(k+1)I(k-1) \quad (k \in \mathbb{N}),$$

## First Mason's conjecture (continued)

### Conjecture (Mason '72)

$$I(k)^2 \geq I(k+1)I(k-1) \quad (k \in \mathbb{N}).$$

Conjecture was proved for **graphic** matroids  
by (**Huh '15**), and for **all** matroids  
by (**Adiprasito–Huh–Katz '18**).

Both proofs used **combinatorial Hodge theory**.

## First Mason's conjecture (continued)

### Conjecture (Mason '72)

$$I(k)^2 \geq I(k+1)I(k-1) \quad (k \in \mathbb{N}).$$

Conjecture was proved for **graphic** matroids by (Huh '15), and for **all** matroids by (Adiprasito–Huh–Katz '18).

Both proofs used **combinatorial Hodge theory**.

We will show that Mason's conjecture is consequence of a **stronger inequality**.

# Stanley–Yan inequality

## Stanley–Yan inequality (simple case)

Let  $\mathcal{M}$  be a matroid with ground set  $X$  and rank  $r$ .

Fix a subset  $S$  of  $X$ . Let

$B(k) :=$  no. of **bases**  $B$  such that  $|B \cap S| = k$ ,  
multiplied by  $r! \times \binom{r}{k}^{-1}$ .

### Theorem (Stanley '81, Yan '23)


*The sequence  $B(1), B(2), \dots$  is log-concave,*

$$B(k)^2 \geq B(k+1)B(k-1) \quad (k \in \mathbb{N}).$$


# Stanley–Yan inequality (simple)

Theorem (Stanley '81, Yan '23)

$$B(k)^2 \geq B(k+1)B(k-1) \quad (k \in \mathbb{N}).$$



Richard Stanley



Alan Yan

## Stanley–Yan inequality (simple)

Theorem (Stanley '81, Yan '23)

$$B(k)^2 \geq B(k+1)B(k-1) \quad (k \in \mathbb{N}).$$

Proved for **regular** matroids by (Stanley '81) using **Alexandrov–Fenchel inequality** for mixed volumes.

Proved for **all** matroids by (Yan '23) using theory of **Lorentzian polynomials**.

**Proof of Mason's conjecture  
using Stanley–Yan inequality**

## Proof of Mason's conjecture using SY inequality

Let

$\mathcal{M}$  := original matroid in Mason's conjecture;

$\mathcal{F}$  := matroid with  $r$  elements and with every subset being independent;

$\mathcal{M}'$  := direct sum of  $\mathcal{M}$  and  $\mathcal{F}$ ;

$S$  := ground set of  $\mathcal{M}$ .

Then

$$I(k) \text{ for } \mathcal{M} = \frac{1}{r!} \times B(k) \text{ for } \mathcal{M}'.$$

# Proof of Mason's conjecture using SY inequality

Since

$$I(k) \text{ for } \mathcal{M} = \frac{1}{r!} \times B(k) \text{ for } \mathcal{M}',$$

we then conclude that

Stanley–Yan inequality for  $\mathcal{M}'$   
implies Mason's conjecture for  $\mathcal{M}$ .



## Stanley–Yan inequality (full version)

Fix  $d \geq 0$ , disjoint subsets  $S, S_1, \dots, S_d$  of  $X$ ,  
and  $\ell_1, \dots, \ell_d \in \mathbb{N}$ .

$B_d(k) :=$  number of bases  $B$  of  $\mathcal{M}$  such that  
 $|B \cap S| = k, |B \cap S_i| = \ell_i$  for  $i \in [d]$ ,  
multiplied by  $r! \times \binom{r}{k, \ell_1, \dots, \ell_d}^{-1}$ .

### Theorem (Stanley '81, Yan '23)

*The sequence  $B_d(1), B_d(2), \dots$  is log-concave,*

$$B_d(k)^2 \geq B_d(k+1)B_d(k-1) \quad (k \in \mathbb{N}).$$

## What we want to do

Theorem (Stanley '81, Yan '23)

*The sequence  $B_d(1), B_d(2), \dots$  is log-concave,*

$$B_d(k)^2 \geq B_d(k+1)B_d(k-1) \quad (k \in \mathbb{N}).$$

Both LHS and RHS of this inequality has  
combinatorial interpretations.


But we will show that this inequality has  
**no combinatorial injective proof.**

## **Combinatorial injective proof**

## Combinatorial injection

An injection  $f : A \rightarrow B$  is **combinatorial** if

- Given  $x \in A$ , the image  $f(x)$  is computable in  $\text{poly}(|x|)$  steps;
- Given  $y \in B$ , it takes  $\text{poly}(|y|)$  steps **to decide if  $y$  is in image of  $f$** ; and if so, the pre-image  $f^{-1}(y)$  is computable in  $\text{poly}(|y|)$  steps.




## Example: Injective proof of binomial inequality

$$\binom{n}{k}^2 \geq \binom{n}{k+1} \binom{n}{k-1} \quad (1 < k < n).$$

This inequality has a **lattice path interpretation**:

$K(a \rightarrow c, b \rightarrow d) :=$  no. of pairs of north-east lattice paths from  $a$  to  $c$  and  $b$  to  $d$ ,

for  $a, b, c, d \in \mathbb{Z}^2$ .



## Example: Injective proof of binomial inequality

Let



$$a = (0, 1), \quad c = (k, n - k + 1),$$

$$b = (1, 0), \quad d = (k + 1, n - k).$$

Then


$$K(a \rightarrow c, b \rightarrow d) = \binom{n}{k},$$

$$K(a \rightarrow d, b \rightarrow c) = \binom{n}{k-1} \binom{n}{k+1}.$$



## Example: Injective proof of binomial inequality

$f : K(a \rightarrow d, b \rightarrow c) \rightarrow K(a \rightarrow c, b \rightarrow d)$   
is defined by **path-swapping injections**.



Images of  $f$  are pairs of lattice paths that **intersects**.


## First main result

### Theorem 1 (C.–Pak '24+)

There is **no combinatorial injective proof** for the Stanley–Yan inequality, assuming *polynomial hierarchy does not collapse*.

The assumption above is slightly stronger than  $P \neq NP$ , and is widely used in Complexity Theory.

## Recall our goal



We will now show that **Stanley–Yan inequality** is strictly more difficult than the **binomial inequality** and **permutation inversion inequality**.

## Complexity class #P

Problems asking about **existence** of  
**NP** := a solution  $S$  for input  $x$ , where validity of  $S$  can be verified in polynomial time.

Problems asking for **number** of solutions  
**#P** :=  $S$  for input  $x$ , where validity of  $S$  can be verified in polynomial time.

### Example (Problem in #P)

*Count the number of proper 3-colorings of graph  $G$ .*

## Complexity class #P: Equivalent definition

A problem is in #P if, for any input  $x$ ,

$$\text{Output} = \sum_{S \in \{0,1\}^{\text{poly}(|x|)}} V(x, S)$$

where

$$V(x, S) \in \{0, 1\}$$

can be evaluated in  $\text{poly}(|x|)$  time.

Note that the size of the **output** is at most exponential relative to the input  $x$ .

## Second main result

Consider the following computational problem:

**Input:** Binary matroid  $\mathcal{M}$ , subsets  $S, S_1, \dots, S_d$ ,  
integers  $k, \ell_1, \dots, \ell_d$ .

**Output:**  $B_d(k)^2 - B_d(k+1) B_d(k-1)$ .

Theorem 2 (C.–Pak '24+)

*The problem above **does not belong to #P**,  
assuming *polynomial hierarchy does not collapse*.*

## Second main result

### Theorem (C.–Pak '24+)

*The problem of computing*

$$B_d(k)^2 - B_d(k+1) B_d(k-1)$$


is **not in #P**, assuming *polynomial hierarchy does not collapse*.

Both LHS and RHS of Stanley–Yan inequality belongs to **#P**, but their difference **does not**.

## Example 1: Binomial inequality

It follows from **path-swapping injections** that

$\binom{n}{k}^2 - \binom{n}{k+1} \binom{n}{k-1} =$  number of **non-intersecting lattice paths** from  $a$  to  $c$  and  $b$  to  $d$ .



Thus the defect of this inequality belongs to **#P**.

## Example 2: Permutation inversion inequality

Let  $a_k$  = number of  $\pi \in S_n$  with  $k$  inversions.

$$\text{Then } \sum_{0 \leq k \leq \binom{n}{2}} a_k q^k = \prod_{i=1}^{n-1} (1 + q + \dots + q^i)$$

is computable in  $\text{poly}(n)$  time.

Thus  $a_k^2 - a_{k+1}a_{k-1}$  is computable in  $\text{poly}(n)$  time;

and thus belongs to  $\#P$ .

## Conclusion

We compare three log-concave inequalities:

Binomial inequality: **in #P**;

Permutation inversion inequality: **in #P**;

Stanley–Yan inequality: **not in #P**.

This differentiates **Stanley–Yan inequality**  
from **binomial inequality** and **permutation  
inversion inequality**.

# Open Problem

## Conjecture

*Defect of Mason's conjecture*

$$I(k)^2 - I(k+1)I(k-1) \notin \#P.$$

We have shown defect of **Stanley–Yan inequality** does not belong to #P, but not **Mason's conjecture**.

# THANK YOU!

Paper: [www.arxiv.org/abs/2309.05764](http://www.arxiv.org/abs/2309.05764)

[www.arxiv.org/abs/2407.19608](http://www.arxiv.org/abs/2407.19608)

Webpage: [www.math.rutgers.edu/~sc2518/](http://www.math.rutgers.edu/~sc2518/)

Email: [sweehong.chan@rutgers.edu](mailto:sweehong.chan@rutgers.edu)


# Polynomial hierarchy

## Level 0: Complexity class P

$P$  := Decision problems that, given input  $x$ , can be solved in  $\text{poly}(|x|)$  time.

### Example (Problem in P)

*Does a graph  $G$  contain a triangle?*



This complexity class is denoted by  $\Sigma_0^P$ .


## Level 1: Complexity class NP

Problems asking about **existence** of


**NP** := a solution  $S$  for input  $x$ , where **validity** of  $S$  can be **verified** in  $\text{poly}(|x|)$  time.

### Example (Problem in NP)

*Does a graph  $G$  have a proper 3-coloring?*



Proper coloring



Improper coloring

This complexity class is denoted by  $\Sigma_1^P$ .

# Oracle machine

An **oracle machine** is a black box capable of solving problems from a **given class** in a single operation.



## Level $i$ of polynomial hierarchy

The class  $\Sigma_i^P := \text{NP}^{\Sigma_{i-1}^P}$  is

Problems asking about existence of a solution  $S$  for input  $x$ , where validity of  $S$  can be verified in  $\text{poly}(|x|)$  time, augmented by  $\Sigma_{i-1}^P$ -oracle.

---

Note that

$$\Sigma_0^P \subseteq \Sigma_1^P \subseteq \Sigma_2^P \subseteq \Sigma_3^P \subseteq \dots$$

# Polynomial hierarchy (PH)

Polynomial hierarchy is the union of all  $\Sigma_i^P$ 's,

$$\text{PH} := \bigcup_{i=0}^{\infty} \Sigma_i^P.$$

## Conjecture

*Polynomial hierarchy does not collapse,*

$$\Sigma_0^P \subsetneq \Sigma_1^P \subsetneq \Sigma_2^P \subsetneq \Sigma_3^P \subsetneq \dots$$

- $\Sigma_0^P \neq \Sigma_1^P$  is equivalent to  $P \neq NP$ .
- $\Sigma_1^P \neq \Sigma_2^P$  is equivalent to  $NP \neq \text{coNP}$ .