



Sorting probability for Young diagrams

Swee Hong Chan (UCLA)

joint with Igor Pak and Greta Panova

1	2	4	7	8
3	5	6	9	10





<



<



<




<






<



<



<



<




<




Partially ordered set

A poset P is a set X with a partial order \preceq on X .




Linear extension

A linear extension L is a complete order of \preceq .




We write $e(P)$ for number of linear extensions of P .

How many steps needed to complete a partial order?




How many steps needed to complete a partial order?

We first compare c and d , and get $c \preceq d$.




How many steps needed to complete a partial order?

We then compare d and e , and get $d \preceq e$.





How many steps needed to complete a partial order?

We continue with b and e , and get $e \preceq b$.



How many steps needed to complete a partial order?

Completing the partial order took 3 steps.



Strategy to complete the partial order

At each step, compare x and y that satisfies

$$\frac{1}{2} - c \leq P[x \preceq y] \leq \frac{1}{2} + c,$$

where P is uniform on linear extensions of P .

Runtime is $\Theta(\log e(P))$ steps.

$\frac{1}{3} - \frac{2}{3}$ Conjecture

Conjecture (Kislitsyn '68, Fredman '75, Linial '84)

For every finite poset that is not completely ordered, there exists x, y :



$$\frac{1}{3} \leq P[x \preceq y] \leq \frac{2}{3}.$$

(Brightwell-Felsner-Trotter '95)

“This problem remains one of the most intriguing problems in the combinatorial theory of posets.”

Why $\frac{1}{3}$ and $\frac{2}{3}$?

The upper, lower bound are achieved by this poset:



$$P[x \preceq y] = \frac{1}{3}; \quad P[y \preceq x] = \frac{2}{3}.$$

What is known so far

Theorem (Kahn-Saks '84)

For every finite poset, there always exists x, y :

$$\frac{3}{11} \leq \mathbb{P}[x \preceq y] \leq \frac{8}{11},$$

roughly between 0.273 and 0.727.

Proof is by applying **mixed-volume inequalities** to **order polytopes**.

What is known so far

Theorem (Brightwell-Felsner-Trotter '95)

For every finite poset, there always exists x, y :

$$\frac{5 - \sqrt{5}}{10} \leq \mathbb{P}[x \preceq y] \leq \frac{5 + \sqrt{5}}{10},$$


roughly between 0.276 and 0.724.

This bound cannot be improved for **infinite posets**.

Young diagrams

Elements of P_λ are **cells** of Young diagram of shape λ .

$x \preceq y$ if y lies to the Southeast of x .



Young diagram of shape $\lambda = (4, 3, 1)$

We write n for **number of cells** of Young diagram.

Young diagrams

Linear extensions of P_λ correspond to **standard Young tableau** of the Young diagram.

1	2	5	6
3	4	7	
8			


Linear extensions are counted by **hook-length formulas**.

What is known for Young diagrams

Theorem 1 (Olson–Sagan '18)

For *Young diagrams*, there always exists x, y :

$$\frac{1}{3} \leq \mathbb{P}[x \preceq y] \leq \frac{2}{3}.$$



What is known for Young diagrams

Theorem 1 (Olson–Sagan '18)

For *Young diagrams*, there always exists x, y :

$$\frac{1}{3} \leq P[x \preceq y] \leq \frac{2}{3}.$$

We sketch an alternative proof for Young diagrams using [Naruse hook-length formulas](#).

Hook-length formulas


Number of standard Young tableau of shape λ is

$$f^\lambda := \frac{n!}{\prod_{x \in \lambda} h_\lambda(x)}.$$

7	6	4	1
5	4	2	
4	3	1	
2	1		

$$f^\lambda = \frac{12!}{764154243121} = 2970$$

Skew Young diagrams




Skew Young diagram of shape λ/μ ,
 $\lambda = (5, 3, 3, 1)$ and $\mu = (2, 1)$.

We write n for **number of cells** in λ ,
and m for **number of cells** in μ .

Excited diagrams

Black boxes can move on SouthEast direction.




Naruse hook-length formulas

Theorem (Naruse '14, Morales-Pak-Panova '17)

Number of skew Young tableau of shape λ/μ is

$$f^{\lambda/\mu} := f^\lambda \frac{(n-m)!}{n!} \sum_{\substack{\text{excited} \\ \text{diagrams } B}} \prod_{\substack{\text{black cells} \\ x \in B}} h_\lambda(x).$$


Naruse hook-length formulas



The number of SYT of shape λ/μ is equal to

$$2970 \frac{9!}{12!} (7 \cdot 6 \cdot 5 + 7 \cdot 5 \cdot 2 + 7 \cdot 2 \cdot 3 + 7 \cdot 6 \cdot 3 + 4 \cdot 2 \cdot 3)$$
$$= 1062.$$

Proof of Theorem Olson–Sagan




$$P[x \preceq y_1] = \underbrace{\quad\quad\quad}_{0 \quad\quad\quad 1}$$

The **jump probabilities** are

$$p_i := P[y_i \preceq x \preceq y_{i+1}]$$

Proof of Theorem Olson–Sagan





$$P[x \preceq y_2] = \int_0^1 \mathbb{1}_{[0, p_1]}(t) dt$$

A diagram illustrating the integral representation of the probability. A horizontal line segment represents the interval from 0 to 1. The left endpoint is labeled '0' and the right endpoint is labeled '1'. A red shaded rectangle is drawn below the line, starting at 0 and ending at a point labeled p_1 . A double-headed arrow below the line indicates the length of this interval is p_1 . Dashed vertical lines connect the endpoints 0 and 1 to the top of the red rectangle.

The **jump probabilities** are

$$p_i := P[y_i \preceq x \preceq y_{i+1}]$$


Proof of Theorem Olson–Sagan



The **jump probabilities** are

$$p_i := P[y_i \preceq x \preceq y_{i+1}]$$

Proof of Theorem Olson–Sagan




$$P[x \preceq y_4] = \int_0^1 \mathbb{1}_{[0, p_1]}(x) \mathbb{1}_{[p_1, p_1+p_2]}(x) \mathbb{1}_{[p_1+p_2, p_1+p_2+p_3]}(x) dx$$

The **jump probabilities** are

$$p_i := P[y_i \preceq x \preceq y_{i+1}]$$

Proof of Theorem Olson–Sagan



$$P[x \preceq y_5] =$$


A horizontal bar representing the interval $[0, 1]$ on the real line. The bar is divided into four segments of decreasing length from left to right, shaded in increasing lightness from dark red to light pink. The segments are labeled p_1 , p_2 , p_3 , and p_4 from left to right. The endpoints are labeled 0 and 1.

The **jump probabilities** are

$$p_i := P[y_i \preceq x \preceq y_{i+1}]$$

Linial-type argument

Suppose that p_1, p_2, \dots, p_ℓ are all $< \frac{1}{3}$.



Look at when the probability exceeds $\frac{1}{3}$. Then

$$\frac{1}{3} \leq \mathbf{P}[x \preceq y_{i+1}] \leq \frac{2}{3}.$$


Proof of $p_1 < \frac{1}{3}$

Suppose to the contrary that $p_1 \geq \frac{1}{3}$. Then

- If $\frac{1}{3} \leq p_1 \leq \frac{2}{3}$, then

$$\frac{1}{3} \leq p_1 = P[x \preceq y_2] \leq \frac{2}{3}.$$


- If $p_1 > \frac{2}{3}$, then **conjugate** to get $p_1 < \frac{1}{3}$.




Skew diagrams enter the scene

It suffices to show $p_1 \geq p_2 \geq \dots \geq p_\ell$.

$$p_1 = P[y_1 \preceq x \preceq y_2] = \frac{\# \text{ of SYTs of } f^\lambda}{f^\lambda}$$




$$p_2 = P[y_2 \preceq x \preceq y_3] = \frac{\# \text{ of SYTs of } f^\lambda}{f^\lambda}$$




Skew diagrams enter the scene

It suffices to show $p_1 \geq p_2 \geq \dots \geq p_\ell$.

$$p_1 = P[y_1 \preceq x \preceq y_2] = \frac{\# \text{ of SYTs of } f^\lambda}{f^\lambda}$$




$$p_2 = P[y_2 \preceq x \preceq y_3] = \frac{\# \text{ of SYTs of } f^\lambda}{f^\lambda}$$




We can now use **NHLF**.


Proof of $p_1 \geq p_2$




$$(10!)(7)(6)$$




$$(10!)(7)(2)$$




$$(10!)(4)(2)$$




$$(9!)(7)(6)(5)$$




$$(9!)(7)(6)(3)$$



$$(9!)(7)(2)(5)$$



$$(9!)(7)(2)(3)$$



$$(9!)(4)(2)(3)$$

$$p_1 = \frac{(10! \cdot 7 \cdot 6 + 10! \cdot 7 \cdot 2 + 10! \cdot 4 \cdot 2)}{12!} = \frac{9!}{12!} 640.$$

$$p_2 = \frac{(9! \cdot 7 \cdot 6 \cdot 8 + 9! \cdot 7 \cdot 2 \cdot 8 + 9! \cdot 4 \cdot 2 \cdot 3)}{12!} = \frac{9!}{12!} 472.$$

Thus we complete the proof of this theorem.


Theorem (Olson–Sagan '18)

There always exists x, y :

$$\frac{1}{3} \leq P[x \preceq y] \leq \frac{2}{3},$$

for poset P_λ of Young diagram of shape λ .

Back to previous example



Comparison probability for this Young diagram is

$$P[x \preceq y] = \frac{16}{33} \approx 0.4848,$$

which is closer to $\frac{1}{2}$ than $\frac{1}{3}$, $\frac{2}{3}$.

What we will do next

Previously, we want to find x, y :

$$\frac{1}{3} \leq \mathbb{P}[x \preceq y] \leq \frac{2}{3},$$

Now, we want to find x, y :

$$\frac{1}{2} - \delta \leq \mathbb{P}[x \preceq y] \leq \frac{1}{2} + \delta,$$

Sorting probability

Sorting probability of a poset P is

$$\delta(P) := \min_{\text{distinct } x, y} |P[x \prec y] - P[y \prec x]|.$$

In particular, there exists x, y :

$$\frac{1}{2} - \frac{\delta(P)}{2} \leq P[x \preceq y] \leq \frac{1}{2} + \frac{\delta(P)}{2}.$$

Kahn–Saks Conjecture

Conjecture (Kahn-Saks '84)

For every finite poset,

$$\delta(P) \rightarrow 0 \quad \text{as} \quad \text{width}(P) \rightarrow \infty.$$

Here $\text{width}(P)$ is the largest size of anti-chains in P .

Komlós '90 proved such a result for posets with $\Omega\left(\frac{n}{\log \log \log n}\right)$ minimal elements.

Our results


First result

Theorem (C.-Pak-Panova '20+)

Let $\lambda_1 \geq \dots \geq \lambda_d \geq \varepsilon n$. For poset P_λ of *Young diagram* of λ ,


$$\delta(P_\lambda) \leq \frac{C}{\sqrt{n}},$$

for some $C = C(d, \varepsilon) > 0$.




Where is the improvement?

Before: x is 2nd element in 1st row, y is in 1st column.




Now: x is middle element in 1st row, y is in 2nd row.




Where is the improvement?

Before: x is 2nd element in 1st row, y is in 1st column.




Now: x is middle element in 1st row, y is in 2nd row.




Where is the improvement?

Before: x is 2nd element in 1st row, y is in 1st column.




Now: x is middle element in 1st row, y is in 2nd row.




Where is the improvement?

Before: x is 2nd element in 1st row, y is in 1st column.




Now: x is middle element in 1st row, y is in 2nd row.



Sketch of proof

After reductions using [Hoeffding's inequality](#),

$$\delta(P_\lambda) \leq \sum_{\mu} \frac{\text{SYTs of } \mu}{f^\lambda}$$



$$\text{with } \mu \approx \left(\frac{\lambda_1}{2} \pm \sqrt{n}, \dots, \frac{\lambda_d}{2} \pm \sqrt{n} \right).$$

Right side is then upper-bounded via [NHLF](#).

Back to first result

Theorem (C.-Pak-Panova '20+)

Let $\lambda_1 \geq \dots \geq \lambda_d \geq \varepsilon n$. For poset P_λ of *Young diagram* of λ ,


$$\delta(P_\lambda) \leq \frac{C}{\sqrt{n}},$$

for some $C = C(d, \varepsilon) > 0$.


Next: better bound for Catalan posets.

Catalan posets, $\lambda = \left(\frac{n}{2}, \frac{n}{2}\right)$

Young diagram is rectangle with 2 rows and n cells.



1	2	4	7
3	5	6	8



Second result


Theorem (C.-Pak-Panova '21)

For Catalan posets with n cells,

$$\delta(P_\lambda) \leq Cn^{-\frac{5}{4}},$$

for some $C > 0$.

How good is this bound?



Open Problem

Show that

$$\limsup_{n \rightarrow \infty} \frac{\log \delta(P_\lambda)}{n} = -\frac{5}{4}; \quad \liminf_{n \rightarrow \infty} \frac{\log \delta(P_\lambda)}{n} < -\frac{5}{4}.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
y								

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := \left| \mathbb{P}[x \prec y(x)] - \mathbb{P}[y(x) \prec x] \right|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
	y							

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := \left| \mathbb{P}[x \prec y(x)] - \mathbb{P}[y(x) \prec x] \right|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		y						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := \left| \mathbb{P}[x \prec y(x)] - \mathbb{P}[y(x) \prec x] \right|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
			y					

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
				y				

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

x								
$y(x)$								

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

	x							
$y(x)$								

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

		x						
	$y(x)$							

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

				x				
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

					x			
			$y(x)$					

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

						x		
				$y(x)$				

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

							x	
						$y(x)$		

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						

Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

								x
							$y(x)$	

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$

Where is the improvement?

Before: x is **fixed** at midpoint, only y is **optimized**.

				x				
		$y(x)$						


Now: **Optimize** $y = y(x)$ for each x , then **optimize** x .

			x					
		$y(x)$						

For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := |P[x \prec y(x)] - P[y(x) \prec x]|.$$


Location of the optimizer $y(x)$ for $n = 2000$



For each x , $y(x)$ is the element that minimizes

$$\delta(x, y(x)) := \left| \mathbb{P}[x \prec y(x)] - \mathbb{P}[y(x) \prec x] \right|.$$

Sorting probability $\delta(P)$ for $n = 2000$



$$\delta(x, y(x)) := \left| \mathbb{P}[x \prec y(x)] - \mathbb{P}[y(x) \prec x] \right|.$$

Back to second result

Theorem (C.-Pak-Panova '21)

For Catalan posets with n cells,

$$\delta(P_\lambda) \leq Cn^{-\frac{5}{4}},$$

for some $C > 0$.

Important: Estimates are not done by NHLF,
but by direct computation.

Better upper bound for general Young diagrams
remain open.

What is next?

Theorem (C.-Pak-Panova '20+)

Let $\lambda_1 \geq \dots \geq \lambda_d \geq \varepsilon n$. For poset P_λ of *Young diagram* of λ , there exists x, y :

$$\delta(P_\lambda) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Open Problem

Prove same result for other *families of posets*, e.g., *k-dimensional Young diagrams* and *periodic posets*.

arXiv preprints: [2005.08390](#) and [2005.13686](#).

Webpage: <http://math.ucla.edu/~sweehong/>

THANK YOU!

arXiv preprints: [2005.08390](#) and [2005.13686](#).

Webpage: <http://math.ucla.edu/~sweehong/>