

Math 350: Linear Algebra

Wolmer V. Vasconcelos

Set 7

Spring 2010

Outline

- 1 Inner Products Spaces
- 2 HomeWork #7
- 3 Gram-Schmidt Orthogonalization
- 4 HomeWork#8
- 5 The Adjoint of a Linear Operator
- 6 Least Squares Approximation
- 7 Old Hourly #2 Questions
- 8 HomeQuiz #7
- 9 Normal Operators
- 10 Unitary Operators
- 11 Goodies
- 12 HomeQuiz #8
- 13 HomeWork #9

Last Class... and Today ...

- Eigenvectors and eigenvalues
- Diagonalization
- Inner products spaces
- Norms

Metric properties of vector spaces

Let \mathbf{V} be a vector space over the field \mathbf{F} . We want to develop a geometry for \mathbf{V} . For that, it is helpful to have a notion of **distance**, or **length**. We will transport and then extend numerous constructions of ordinary geometry and their calculus.

We will restrict ourselves to the cases of $\mathbf{F} = \mathbb{R}$, or $\mathbf{F} = \mathbb{C}$. In the case of \mathbb{C} , we use the standard notation for the **complex conjugate** of the complex number $z = a + bi$

$$\bar{z} = a - bi.$$

Some of its properties are:

$$\begin{aligned} z\bar{z} &= a^2 + b^2 \\ \overline{z_1 + z_2} &= \bar{z}_1 + \bar{z}_2 \\ \overline{z_1 \cdot z_2} &= \bar{z}_1 \cdot \bar{z}_2 \\ \frac{1}{z} &= \frac{\bar{z}}{z \cdot \bar{z}}, \quad z \neq 0 \end{aligned}$$

For certain operations, like solving polynomial equations, the polar representation of complex numbers

$$a + bi = r(\cos \theta + i \sin \theta), \quad r = \sqrt{a^2 + b^2}, \quad \tan \theta = \frac{a}{b}$$

is useful. For instance,

$$\sqrt{i} = \pm(\cos \pi/2 + i \sin \pi/2)^{1/2} = \pm(\cos \pi/4 + i \sin \pi/4) = \pm \frac{\sqrt{2}}{2}(1 + i).$$

Inner product space

An **inner product vector space** \mathbf{V} is a V.S. over \mathbb{R} or \mathbb{C} with a mapping

$$\mathbf{V} \times \mathbf{V} \rightarrow \mathbf{F}, \quad (u, v) \rightarrow \langle u, v \rangle = u \cdot v \in \mathbf{F}$$

satisfying certain conditions. Let us give an example to guide us in what is needed. Let $\mathbf{V} = \mathbb{R}^n$ and define

$$\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix} = a_1 b_1 + \cdots + a_n b_n = \sum_{i=1}^n a_i b_i$$

Note the properties: **bi-additive** ; $v \cdot v$ is a non-negative real number, so we can use $\sqrt{v \cdot v}$ to define the **magnitude** of v .

Question: Could we use the same formula to define an inner product for \mathbb{C}^n ? Well... $(i) \cdot (i)$ would be -1 . Of course the formula still defines a nice bilinear mapping but would not meet our need.

Dot product

Definition

An inner product vector space is a vector space with a mapping

$$\mathbf{V} \times \mathbf{V} \rightarrow \mathbf{F}, \quad (u, v) \rightarrow u \cdot v \in \mathbf{F}$$

satisfying:

- 1 $(u_1 + u_2) \cdot v = u_1 \cdot v + u_2 \cdot v$
- 2 $(cu) \cdot v = c(u \cdot v)$
- 3 $\overline{u \cdot v} = v \cdot u$
- 4 $u \cdot u > 0$ if $u \neq 0$

The better notation for this product is

$$u \cdot v = \langle u, v \rangle$$

Examples

Of course, the example above of \mathbb{R}^n is the grandmother of all examples. Let us modify it a bit to get an example for \mathbb{C}^n :

$$\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix} = a_1 \bar{b}_1 + \cdots + a_n \bar{b}_n = \sum_{i=1}^n a_i \bar{b}_i.$$

Note the properties: **additive** ; $v \cdot v$ is a non-negative real number

$$v \cdot v = \sum_{i=1}^n a_i \bar{a}_i$$

so we can use $\sqrt{v \cdot v}$ to define the **magnitude** of v . Note the lack of full symmetry.

Example of Function Space

Let us give an example from left field: Let \mathbf{V} be the vector space of all real continuous functions on the interval $[a, b]$, and define for $f(t), g(t) \in \mathbf{V}$,

$$\langle f(t), g(t) \rangle = f(t) \cdot g(t) = \int_a^b f(t)g(t)dt.$$

An important case: If m, n are integers,

$$\langle \sin nt, \cos mt \rangle = \int_0^{2\pi} \sin nt \cos mt dt = 0$$

$$\langle \sin nt, \sin mt \rangle = \int_0^{2\pi} \sin nt \sin mt dt = 0, \quad m \neq n$$

$$\langle \cos nt, \cos mt \rangle = \int_0^{2\pi} \cos nt \cos mt dt = 0, \quad m \neq n$$

$$\langle \sin nt, \sin nt \rangle = \int_0^{2\pi} \sin^2 nt dt = \pi, \quad n \neq 0$$

Example: $M_n(\mathbf{F})$

Let $\mathbf{V} = M_n(\mathbf{F})$ be the V.S. of all n -by- n matrices. For any such matrix $\mathbf{A} = [a_{ij}]$ define the **adjoint** of \mathbf{A} (unfortunately we have already used the word for a very different notion!) to be the matrix

$$\mathbf{A}^* = [\bar{a}_{ji}],$$

that is, we transpose \mathbf{A} and take the complex conjugate of each entry. Define the product (Frobenius product)

$$\langle \mathbf{A}, \mathbf{B} \rangle = \text{trace}(\mathbf{AB}^*) = \sum_i (\mathbf{AB}^*)_{ii}.$$

It is clear that this product has the properties of an inner product. We just check the positivity condition:

$$\begin{aligned} \langle \mathbf{A}, \mathbf{A} \rangle &= \text{trace}(\mathbf{AA}^*) = \sum_i (\mathbf{AA}^*)_{ii} \\ &= \sum_i \sum_j a_{ij} \bar{a}_{ij} = \sum_i |a_{ij}|^2 \geq 0 \end{aligned}$$

Proposition

If \mathbf{V} is an inner product space, the following hold:

- 1 $\langle u, v + w \rangle = \langle u, v \rangle + \langle u, w \rangle$
- 2 $\langle u, cv \rangle = \bar{c} \langle u, v \rangle$
- 3 $\langle u, O \rangle = \langle O, v \rangle = 0$
- 4 $\langle u, u \rangle = 0$ iff $u = O$
- 5 $\langle u, v \rangle = \langle u, w \rangle$ for all $u \in \mathbf{V}$ then $v = w$

Proof of 1: Note

$$\begin{aligned} \langle u, v + w \rangle &= \overline{\langle v + w, u \rangle} = \overline{\langle v, u \rangle + \langle w, u \rangle} \\ &= \overline{\langle v, u \rangle} + \overline{\langle w, u \rangle} = \langle u, v \rangle + \langle u, w \rangle \end{aligned}$$

Length of a vector

Definition

Let \mathbf{V} , $\langle \cdot, \cdot \rangle$ be an inner product space. If $v \in \mathbf{V}$, the **length** or **norm** of v is the real number $\|v\| = \sqrt{\langle v, v \rangle}$.

If $\mathbf{V} = \mathbb{C}^n$, $v = (a_1, \dots, a_n)$,

$$\|v\| = \left[\sum_{i=1}^n |a_i|^2 \right]^{1/2}$$

If \mathbf{V} is the space of real continuous functions on $[0, 1]$ and inner product is that we defined previously,

$$\|f(t)\|^2 = \int_0^1 f(t)^2 dt.$$

Framework for Geometry

The following assertions permits the construction of 'recognizable' objects in any inner product space:

Theorem

If \mathbf{V} is an inner product space, then for all $u, v \in \mathbf{V}$

① [Cauchy-Schwarz Inequality]

$$|\langle u, v \rangle| \leq \|u\| \cdot \|v\|$$

② [Triangle Inequality]

$$\|u + v\| \leq \|u\| + \|v\|.$$

The **Cauchy-Schwarz Inequality** will allow the introduction of **angles** and its **trigonometry** in \mathbf{V} , while the **Triangle Inequality** will lead to many constructions extending those we are familiar with in 2- and 3-space.

Proofs of CSI and Δ -Inequality

To prove Cauchy-Schwarz Inequality: Note that for ANY $c \in \mathbf{F}$, $v \neq 0$

$$\begin{aligned} 0 \leq \|u - cv\|^2 &= \langle u - cv, u - cv \rangle = \langle u, u - cu \rangle - c \langle v, u - cv \rangle \\ &= \langle u, u \rangle - \bar{c} \langle u, v \rangle - c \langle v, u \rangle + c\bar{c} \langle v, v \rangle \end{aligned}$$

If we set $c = \frac{\langle u, v \rangle}{\langle v, v \rangle}$ the inequality becomes

$$0 \leq \langle u, u \rangle - \frac{|\langle u, v \rangle|^2}{\|v\|^2},$$

which proves the assertion.

For the Δ -inequality: Consider

$$\begin{aligned}
 \|u + v\|^2 &= \langle u + v, u + v \rangle = \langle u, u \rangle + \langle u, v \rangle + \langle v, u \rangle + \langle v, v \rangle \\
 &= \|u\|^2 + (\langle u, v \rangle + \overline{\langle u, v \rangle}) + \|v\|^2 = \|u\|^2 + 2\Re\langle u, v \rangle + \|v\|^2 \\
 &\leq \|u\|^2 + 2|\langle u, v \rangle| + \|v\|^2 \\
 &\leq \|u\|^2 + 2\|u\| \cdot \|v\| + \|v\|^2 \quad \text{by C-S inequality} \\
 &= (\|u\| + \|v\|)^2.
 \end{aligned}$$

We used that for any complex number $z = a + bi$, its real part $\Re z = a \leq |z| = \sqrt{a^2 + b^2}$.

Example

To illustrate the power of the axiomatic method, compare the proof above [which holds for ALL examples] with the work needed to check the inequalities just the case of the following example:

$$\left| \sum_{i=1}^n a_i \bar{b}_i \right| \leq \left[\sum_{i=1}^n |a_i|^2 \right]^{1/2} \left[\sum_{i=1}^n |b_i|^2 \right]^{1/2}$$

$$\left[\sum_{i=1}^n |a_i + b_i|^2 \right]^{1/2} \leq \left[\sum_{i=1}^n |a_i|^2 \right]^{1/2} + \left[\sum_{i=1}^n |b_i|^2 \right]^{1/2}$$

Angles and Distances

Equipped with these results, we can define angles and distances, with many of the usual properties, in any inner product space. For example, for a real inner product space, the Cauchy-Schwarz inequality says that for any two [will assume nonzero] vectors u, v ,

$$\langle u, v \rangle \leq \|u\| \cdot \|v\|,$$

that is

$$-1 \leq \frac{\langle u, v \rangle}{\|u\| \cdot \|v\|} \leq 1$$

This means that the ratio can be identified to the cosine, $\cos \alpha$, of a unique angle $0 \leq \alpha \leq \pi$: So we can write

$$\langle u, v \rangle = \|u\| \cdot \|v\| \cos \alpha$$

and say that α is the **angle** between the vectors u and v .

An important relationship between two vectors u, v is when $\langle u, v \rangle = 0$: We then say that u and v are **orthogonal** or **perpendicular**. One notation for this situation is:

$$u \perp v$$

The **distance** between the vectors u, v is defined by

$$\text{dist}(u, v) = \|u - v\| = \langle u - v, u - v \rangle^{1/2}$$

One of its properties follow from the triangle inequality: If u, v, w are three vectors

$$\text{dist}(u, w) \leq \text{dist}(u, v) + \text{dist}(v, w).$$

Properties

These notions have numerous consequences. Let us begin with:

Proposition

Let v_1, \dots, v_n be nonzero vectors of the inner product space \mathbf{V} . If $v_i \perp v_j$ for $i \neq j$, then these vectors are linearly independent.

Proof.

Suppose we have a linear combination

$$c_1 v_1 + c_2 v_2 + \cdots + c_n v_n = O.$$

We claim all $c_i = 0$. To prove, say $c_1 = 0$, take the inner product of the linear combination with v_1 :

$$c_1 \underbrace{\langle v_1, v_1 \rangle}_{\neq 0} + c_2 \underbrace{\langle v_2, v_1 \rangle}_{=0} + \cdots + c_n \underbrace{\langle v_n, v_1 \rangle}_{=0} = \langle O, v_1 \rangle = 0.$$

A vector v of length $\|v\| = 1$ is called a **unit** vector. They are easy to find: given a nonzero vector u , $v = \frac{u}{\|u\|}$ is a unit vector.

A set of vectors v_1, \dots, v_n is said to be **orthonormal** if $v_i \perp v_j$, for $i \neq j$ and $\|v_i\| = 1$ for any i . Of course, a good example are the ordinary coordinate vectors of 3-space.

Proposition

Let \mathbf{V} be an inner product space with an *orthonormal* basis v_1, \dots, v_n . Then for any $v \in \mathbf{V}$,

$$v = c_1 v_1 + \dots + c_n v_n,$$

where $c_i = \langle v, v_i \rangle$. The c_i are called the *Fourier coefficients* of v relative to the basis.

Proof.

To get c_i , it suffices to form the inner product of v with v_i :

$$\langle v, v_i \rangle = c_i \langle v_i, v_i \rangle = c_i,$$

since $\langle v_i, v_i \rangle = 1$ and all other $\langle v_j, v_i \rangle = 0$. □

Matrix representation

Orthonormal bases are also useful in finding the matrix representation of a L.T. $\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$:

Let $\mathcal{A} = \{v_1, \dots, v_n\}$ be such a basis. Then $[\mathbf{T}]_{\mathcal{A}} = [a_{ij}]$ where a_{ij} are the coefficients in the expression

$$\mathbf{T}(v_j) = a_{1j}v_1 + \cdots + a_{ij}v_i + \cdots + a_{nj}v_n$$

To select a_{ij} it suffices to 'dot' with v_i

$$\langle \mathbf{T}(v_j), v_i \rangle = a_{1j} \underbrace{\langle v_1, v_i \rangle}_{=0} + \cdots + a_{ij} \underbrace{\langle v_i, v_i \rangle}_{=1} + \cdots + a_{nj} \underbrace{\langle v_n, v_i \rangle}_{=0}$$

$$[\mathbf{T}]_{\mathcal{A}} = [\langle \mathbf{T}(v_j), v_i \rangle]$$

Parallelogram Law

Exercise: If u, v are vectors of an inner product space \mathbf{V} , verify the parallelogram law:

$$\|u + v\|^2 + \|u - v\|^2 = 2(\|u\|^2 + \|v\|^2).$$

Draw a picture to illustrate this equality.

HomeWork #7

Section 6.1: 2, 5, 9, 10, 11, 18, 27 (challenge)

Things to come

- 1 We will prove that every finite-dimensional vector space \mathbf{W} of an inner product space \mathbf{V} has an orthonormal basis.
- 2 This will allow us to express the distance from a vector $v \in \mathbf{V}$ to the subspace \mathbf{W} . For instance, if

$$\mathbf{Ax} = \mathbf{b}$$

is a consistent system of linear equations, that is, if there is some solution $\mathbf{Ax}_0 = \mathbf{b}$, we know that the solution set is the set

$$\mathbf{x}_0 + N(\mathbf{A}),$$

where $N(\mathbf{A})$ is the nullspace of \mathbf{A} . Now we will be able to find the solution of smallest length, if need be.

Let us show how to obtain an orthonormal basis of a vector space from an arbitrary basis $\mathcal{A} = \{u_1, \dots, u_n\}$.

If $n = 1$, $w_1 = \frac{u_1}{\|u_1\|}$ is the answer.

Assume now that we have a basis of two vectors u_1, u_2 . We need to find two nonzero vectors v_1, v_2 in the span of u_1, u_2 so that $v_1 \perp v_2$. We use a projection trick: we set $v_1 = u_1$ and look for c so that


$$v_2 = u_2 - cu_1 \perp v_1,$$

that is

$$\langle v_2, v_1 \rangle = \langle u_2, v_1 \rangle - c\langle u_1, v_1 \rangle = 0$$

$$c = \frac{\langle u_2, v_1 \rangle}{\langle v_1, v_1 \rangle}$$

Observe that v_1, v_2 have same span as u_1, u_2 . Now replace v_i by $v_i/\|v_i\|$.



$w = \text{Projection of } v \text{ along } u$

Projection formula

If \mathbf{L} is a line defined by the vector $u \neq O$ and v is another vector,

$$w = \frac{\langle v, u \rangle}{\langle u, u \rangle} u$$

is the **projection** of v along \mathbf{L} or u .

Proposition

$v - w$ is perpendicular to \mathbf{L} and the smallest distance from v to any vector of \mathbf{L} is $\|v - w\|$.

Proof.

We have already seen that $v - w \perp v$. If cu is a vector of \mathbf{L} , the square distance from v to cu is $(v - w \perp \mathbf{L}$, so will use Pythagorean Theorem)

$$\|v - cu\|^2 = \|(v - w) + (w + cu)\|^2 = \|v - w\|^2 + \underbrace{\|w + cu\|^2}_{\geq 0}.$$

Gram-Schmidt Algorithm

The routine to obtain a basis that is orthogonal from another basis [Gram-Schmidt process]:

- 1 Input: $\mathcal{A} = \{u_1, \dots, u_n\}$ given basis
- 2 Set $v_1 = u_1$
- 3 Compute v_2, \dots, v_n successively, one at a time, by

$$v_i = u_i - \underbrace{\left(\frac{u_i \cdot v_1}{v_1 \cdot v_1} v_1 + \frac{u_i \cdot v_2}{v_2 \cdot v_2} v_2 + \dots + \frac{u_i \cdot v_{i-1}}{v_{i-1} \cdot v_{i-1}} v_{i-1} \right)}_{\text{projection of } u_i \text{ onto span}\{v_1, \dots, v_{i-1}\}}$$

- 4 Set $w_i = \frac{v_i}{\|v_i\|}$
- 5 Output: $\mathcal{B} = \{w_1, \dots, w_n\}$ is an orthonormal basis.

Hadamard's Inequality

Let \mathbf{A} be a matrix whose columns form a basis $\{u_1, u_2, \dots, u_n\}$ of \mathbb{R}^n (put $n = 3$ for simplicity)

$$\mathbf{A} = [u_1 \mid u_2 \mid u_3]$$

Now consider the matrix

$$\mathbf{B} = [v_1 \mid v_2 \mid v_3] = [u_1 \mid u_2 - a_1 u_1 \mid u_3 - b_1 u_1 - b_2 u_2]$$

where the coefficients are chosen for that the v_i 's are perpendicular to one another. Note that \mathbf{B} is obtained from \mathbf{A} by adding scalar multiples of columns to another, so

$$\det(\mathbf{A}) = \det(\mathbf{B}).$$

Furthermore, for each i

$$\|v_i\| \leq \|u_i\|$$

by the projection formula.

Let us calculate $\det(\mathbf{A})^2$:

$$\begin{aligned}
 \det(\mathbf{A})^2 &= \det(\mathbf{B})^2 = \det(\mathbf{B}) \det(\mathbf{B}^t) \\
 &= \det[v_1 \mid v_2 \mid v_3] \det[v_1 \mid v_2 \mid v_3]^t \\
 &= \begin{bmatrix} \langle v_1, v_1 \rangle & 0 & 0 \\ 0 & \langle v_2, v_2 \rangle & 0 \\ 0 & 0 & \langle v_3, v_3 \rangle \end{bmatrix} \\
 &= \prod \langle v_i, v_i \rangle
 \end{aligned}$$

Theorem (Hadamard)

For any square real matrix $\mathbf{A} = [u_1, \dots, u_n]$,

$$|\det(\mathbf{A})|^2 \leq \prod_{i=1}^n \langle u_i, u_i \rangle.$$

For instance, if \mathbf{A} is a 4×4 whose entries are $0, 1, -1$, its column vectors have length at most 2, so that $\det(\mathbf{A}) \leq 16$. According to Joe, there is a such a matrix.

General Projection Formula

Proposition

Let \mathbf{W} be a subspace with an orthonormal basis $\mathcal{A} = \{u_1, \dots, u_n\}$. For any vector v , the vector of \mathbf{W}

$$w = \text{proj}_{\mathbf{W}}(v) = \langle v, u_1 \rangle u_1 \cdots + \langle v, u_n \rangle u_n$$

is the **projection** of v onto \mathbf{W} . It has the following properties

- 1 $v - w$ is perpendicular to any vector of \mathbf{W} . (We say that it is perpendicular to \mathbf{W})
- 2 $\|v - w\|$ is the shortest distance from v to \mathbf{W} .

The proof is like above.

Orthogonal Complement

If \mathbf{W} is a subspace of an inner product space \mathbf{V} , its **orthogonal complement** \mathbf{W}^\perp is the set of all vectors v that are perpendicular to each vector w of \mathbf{W} . In ordinary 3-space \mathbb{R}^3 , the z -axis is the orthogonal complement of the xy -plane.

Proposition

\mathbf{W}^\perp is a subspace of \mathbf{V} .

Proof.

Clearly $0 \in \mathbf{W}^\perp$. If $v_1, v_2 \in \mathbf{W}^\perp$, for any vector $w \in \mathbf{W}$

$$\langle c_1 v_1 + c_2 v_2, w \rangle = c_1 \langle v_1, w \rangle + c_2 \langle v_2, w \rangle = 0,$$

so \mathbf{W}^\perp passes the subspace test. □

Example

Let \mathbf{A} be an $m \times n$ real matrix. The nullspace of \mathbf{A} is the set of all n -tuples \mathbf{x} such that

$$\mathbf{Ax} = \mathbf{0}.$$

This means that the nullspace is the orthogonal complement of the row space of \mathbf{A} :

$$N(\mathbf{A}) = \text{row space}^\perp.$$

Similarly, the **left** nullspace of \mathbf{A} , left $N(\mathbf{A})$, are the m -tuples \mathbf{y} such that

$$\mathbf{yA} = \mathbf{0}$$

that is the orthogonal complement of the column space of \mathbf{A} .

These observations suggest several properties of the \perp operation:

- ① Let \mathbf{V} be a vector space with a basis e_1, \dots, e_n . If \mathbf{W} is spanned by u_1, \dots, u_m , \mathbf{W}^\perp is the set of all vectors $x_1 e_1 + \dots + x_n e_n$ such that

$$x_1 \langle e_1, u_i \rangle + \dots + x_n \langle e_n, u_i \rangle = 0, \quad i = 1, \dots, m.$$

Thus we find \mathbf{W} by solving a system of linear equations.

- ② $\mathbf{W} \cap \mathbf{W}^\perp = \{O\}$.
- ③ $\underbrace{\dim \mathbf{W} + \dim \mathbf{W}^\perp}_{= \dim \mathbf{V}}$
- ④ $(\mathbf{W}^\perp)^\perp = \mathbf{W}$

Proposition

$$\dim \mathbf{W} + \dim \mathbf{W}^\perp = \dim \mathbf{V}.$$

Proof.

Let u_1, \dots, u_m be an orthonormal basis of \mathbf{W} . We define a mapping $\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$ as follows

$$\mathbf{T}(v) = \langle v, u_1 \rangle u_1 + \cdots + \langle v, u_m \rangle u_m.$$

\mathbf{T} is clearly a linear transformation: This is the orthogonal projection of \mathbf{V} onto \mathbf{W} . Its range $R(\mathbf{T})$ is \mathbf{W} . Its nullspace $N(\mathbf{T})$ is the set of vectors v such that $\langle v, u_j \rangle = 0$ for each u_j . This is precisely \mathbf{W}^\perp . From the dimension formula

$$\dim \mathbf{V} = \dim R(\mathbf{T}) + \dim N(\mathbf{T}) = \dim \mathbf{W} + \dim \mathbf{W}^\perp.$$



HomeWork #8

Section 6.2: 2a, 4, 9, 15, 22 (too laborious)

If \mathbf{V} is a vector space over the field \mathbf{F} , a linear functional is a linear transformation

$$\mathbf{f} : \mathbf{V} \longrightarrow \mathbf{F}.$$

For example, if $\mathbf{V} = \mathbf{F}^n$ and $\mathbf{a} = [a_1, \dots, a_n]$ is a matrix, then for every column vector $v \in \mathbf{F}^n$, the function

$$v \longrightarrow \mathbf{a} \cdot v$$

is a linear functional. In fact, every linear functional \mathbf{f} has this description.

Inner product spaces, finite/infinite dimensional have a natural method to define linear functionals. Let us exploit it.

Let \mathbf{V} be an inner product space. If $u \in \mathbf{V}$, the mapping

$$\mathbf{f} : \mathbf{V} \rightarrow \mathbf{F}, \quad \mathbf{f}(v) = \langle v, u \rangle$$

is a linear functional. Observe that if $\langle v, u \rangle = \langle v, w \rangle$, for all v , then $\langle v, u - w \rangle = 0$ and therefore $u = w$.

Proposition

If \mathbf{V} is a finite-dimensional inner product space, for every linear functional \mathbf{f} on \mathbf{V} , there is a unique vector u such that $\mathbf{f}(v) = \langle v, u \rangle$ for all $v \in \mathbf{V}$.

Proof.

Let v_1, \dots, v_n be an orthonormal basis of \mathbf{V} , and let

$$u = \overline{\mathbf{f}(v_1)}v_1 + \cdots + \overline{\mathbf{f}(v_n)}v_n.$$

Note that for each v_j , $\langle v_j, u \rangle = \overline{\overline{\mathbf{f}(v_j)}} = \mathbf{f}(v_j)$, so the functionals defined by u and \mathbf{f} agree on each basis vector, so are equal. □

Adjoint of a Linear Transformation

Let \mathbf{T} be a L.T. of the inner product space \mathbf{V} . We are going to build another L.T. associated to \mathbf{T} , which will be called the **adjoint** of \mathbf{T} . It is the parent [or child] of the transpose!

Fix the vector $u \in \mathbf{V}$. Consider the mapping $v \rightarrow \langle \mathbf{T}(v), u \rangle$. This is a linear functional. According to the previous Proposition, there is a unique w such that

$$\langle \mathbf{T}(v), u \rangle = \langle v, w \rangle, \quad \forall v \in \mathbf{V}.$$

We set $w = \mathbf{S}(u)$. This gives a function $\mathbf{S} : \mathbf{V} \rightarrow \mathbf{V}$. It is routine to check that if $w_1 = \mathbf{S}(u_1)$ and $w_2 = \mathbf{S}(u_2)$, then $\mathbf{S}(u_1 + u_2) = w_1 + w_2$, and also $\mathbf{S}(cu) = c\mathbf{S}(u)$. This L.T. is denoted \mathbf{T}^* and termed the adjoint of \mathbf{T} .

Proposition

Let \mathbf{T} be a L.T. and let $\mathbf{A} = [a_{ij}]$ be its matrix representation relative to the orthonormal basis v_1, \dots, v_n . Then the matrix representation of the adjoint \mathbf{T}^* is $\overline{\mathbf{A}}^t = [\overline{a_{ji}}]$, the conjugate transpose of \mathbf{A} .

Proof.


To find the matrix representation $[b_{ij}]$ of \mathbf{T}^* we write $\mathbf{T}^*(v_j) = \sum_i b_{ij} v_i$, so that

$$\overline{b_{ij}} = \langle v_i, \mathbf{T}^*(v_j) \rangle = \langle \mathbf{T}(v_i), v_j \rangle = a_{ji},$$

as desired. □

Problem

Given 3 (or more) points $P_1 = (x_1, y_1)$, $P_2 = (x_2, y_2)$, $P_3 = (x_3, y_3)$ in \mathbb{R}^2 , find the best fit line (what does this mean?):



$$Y = at + b, \quad Y_i = at_i + b, \quad \text{error} = |Y_i - y_i|$$

t	y	Y
t_1	y_1	Y_1
\vdots	\vdots	\vdots
t_n	y_n	Y_n

$$\mathbf{E} = \text{Square Error} = \sum_{i=1}^n |Y_i - y_i|^2 = \sum_{i=1}^n |at_i + b - y_i|^2$$

Problem: Find a and b so that the square error is as small as possible. To answer, we first write the problem in vector notation.

$$\mathbf{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}, \quad \mathbf{A} = \begin{bmatrix} t_1 & 1 \\ \vdots & \vdots \\ t_m & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} a \\ b \end{bmatrix}$$

$$\mathbf{E} = \|\mathbf{y} - \mathbf{Ax}\|^2$$

We are going to do much better: Given a $m \times n$ matrix \mathbf{A} and a vector $\mathbf{y} \in \mathbf{F}^m$, we are going to find a vector $\mathbf{x}_0 \in \mathbf{F}^n$ such that

$$\|\mathbf{y} - \mathbf{Ax}_0\|^2 \leq \|\mathbf{y} - \mathbf{Ax}\|^2$$

for all $\mathbf{x} \in \mathbf{F}^n$

We know that the answer to this will be affirmative: Let \mathbf{W} be the range of \mathbf{A} , that is the set of all vectors \mathbf{Ax} , for $\mathbf{x} \in \mathbf{F}^n$. There is a vector $w \in \mathbf{W}$, that is $w = \mathbf{Ax}_0$ such that

$$\|\mathbf{y} - \mathbf{Ax}_0\|^2 \leq \|\mathbf{y} - \mathbf{Ax}\|^2.$$

The issue is how to find \mathbf{x}_0 more explicitly. For this we use the notion of the adjoint of a linear transformation:

$$\mathbf{T} : \mathbf{F}^n \rightarrow \mathbf{F}^m, \quad \mathbf{T}^* : \mathbf{F}^m \rightarrow \mathbf{F}^n$$

$$\langle \mathbf{T}(u), v \rangle_m = \langle u, \mathbf{T}^*(v) \rangle_n$$

To derive the desired formula (known as the projection formula) we need two properties of \mathbf{T}^* .

Proposition

Let \mathbf{A} be an $m \times n$ complex matrix and \mathbf{A}^* its adjoint (conjugate transpose). Then

- 1 $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^* \mathbf{A})$.
- 2 If $\text{rank}(\mathbf{A}) = n$ then $\mathbf{A}^* \mathbf{A}$ is invertible.

Proof.

It will suffice to show that \mathbf{A} and $\mathbf{A}^* \mathbf{A}$ have the same nullspace. Why? If $\mathbf{A}^* \mathbf{A}(\mathbf{x}) = \mathbf{0}$, then for all $\mathbf{z} \in \mathbf{F}^n$

$$0 = \langle \mathbf{A}^* \mathbf{A}(\mathbf{x}), \mathbf{z} \rangle_n = \langle \mathbf{A}\mathbf{x}, (\mathbf{A}^*)^* \mathbf{z} \rangle_m = \langle \mathbf{A}\mathbf{x}, \mathbf{A}\mathbf{z} \rangle_m =$$

so $\mathbf{A}\mathbf{x} = \mathbf{0}$ by choosing $\mathbf{z} = \mathbf{x}$.

The second assertion now follows: Since $\mathbf{A}^* \mathbf{A}$ is an $n \times n$ matrix of rank n , it is invertible. □

Projection Formula

Theorem

Let \mathbf{A} be an $m \times n$ complex matrix and let $\mathbf{y} \in \mathbf{F}^m$. Then there exists $\mathbf{x}_0 \in \mathbf{F}^n$ such that $\mathbf{A}^* \mathbf{A}(\mathbf{x}_0) = \mathbf{A}^* \mathbf{y}$ and $\|\mathbf{A}\mathbf{x}_0 - \mathbf{y}\| \leq \|\mathbf{A}\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x} \in \mathbf{F}^n$. If \mathbf{A} has rank n then

$$\mathbf{x}_0 = (\mathbf{A}^* \mathbf{A})^{-1} \mathbf{A}^* \mathbf{y}.$$

Proof.

Since $\mathbf{A}\mathbf{x}_0 - \mathbf{y}$ is perpendicular to the range of \mathbf{A} ,

$$0 = \langle \mathbf{A}\mathbf{x}, \mathbf{A}\mathbf{x}_0 - \mathbf{y} \rangle_m = \langle \mathbf{x}, \mathbf{A}^*(\mathbf{A}\mathbf{x}_0 - \mathbf{y}) \rangle = \langle \mathbf{x}, ((\mathbf{A}^* \mathbf{A})\mathbf{x}_0 - \mathbf{A}^* \mathbf{y}) \rangle$$

for all $\mathbf{x} \in \mathbf{F}^n$. Thus $(\mathbf{A}^* \mathbf{A})\mathbf{x}_0 - \mathbf{A}^* \mathbf{y} = \mathbf{0}$ and therefore

$$\mathbf{x}_0 = (\mathbf{A}^* \mathbf{A})^{-1} \mathbf{A}^* \mathbf{y},$$

that completes the proof □

Illustration

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \end{bmatrix}, \quad \text{rank}(\mathbf{A}) = 2, \quad \mathbf{y} = \begin{bmatrix} 2 \\ 3 \\ 5 \\ 7 \end{bmatrix}$$

$$\mathbf{A}^* \mathbf{A} = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \end{bmatrix} = \begin{bmatrix} 30 & 10 \\ 10 & 4 \end{bmatrix}$$

$$(\mathbf{A}^* \mathbf{A})^{-1} = \frac{1}{20} \begin{bmatrix} 4 & -10 \\ -10 & 30 \end{bmatrix}$$

$$\mathbf{x}_0 = \begin{bmatrix} a \\ b \end{bmatrix} = \frac{1}{20} \begin{bmatrix} 4 & -10 \\ -10 & 30 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 & 4 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 5 \\ 7 \end{bmatrix} = \begin{bmatrix} 1.7 \\ 0 \end{bmatrix}$$

Answer: The least squares line is

$$y = 1.7t$$

The error is

$$\mathbf{E} = \|\mathbf{Ax}_0 - \mathbf{y}\|^2 = 0.3$$

The method is very general: Suppose we are given a number of points and we want to fit a quadratic polynomial

$$Y = at^2 + bt + c$$

to the data.

$$\mathbf{A} = \begin{bmatrix} t_1^2 & t_1 & 1 \\ \vdots & \vdots & \vdots \\ t_n^2 & t_n & 1 \end{bmatrix} \quad \mathbf{x}_0 = \begin{bmatrix} a \\ b \\ c \end{bmatrix}, \quad \mathbf{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$$

Now $\text{rank}(\mathbf{A}) = 3$ if there are 3 distinct values of t .

Shortest solution

We are going to find the **shortest** solution of a consistent system of equations ($m \times n$)


$$\mathbf{Ax} = \mathbf{b}.$$

This will be a solution u such that $\|u\|$ is minimal. The argument will also show that u is unique.

Let \mathbf{x}_0 be a special solution and denote by $N(\mathbf{A})$ the **nullspace** of \mathbf{A} . The solution set is

$$\mathbf{x}_0 + N(\mathbf{A}) = \{\mathbf{x}_0 + \mathbf{v}, \quad \mathbf{v} \in N(\mathbf{A})\}.$$

To pick out of this set the vector $\mathbf{x}_0 + \mathbf{v}$ of smallest length, note that $\|\mathbf{x}_0 + \mathbf{v}\|$ is the distance from \mathbf{x}_0 to $-\mathbf{v}$. So we have our answer: Pick for $-\mathbf{v}$ the projection w of \mathbf{x}_0 into $N(\mathbf{A})$. Then $s = \mathbf{x}_0 - w$ is the desired solution:



$w = \text{Projection of } \mathbf{x}_0 \text{ along } N(\mathbf{A})$

One algorithm for the shortest solution

- 1 Find an orthonormal basis u_1, \dots, u_r for $N(\mathbf{A})$
- 2 Determine the projection w of \mathbf{x}_0 onto $N(\mathbf{A})$:

$$w = \sum_{i=1}^r \langle \mathbf{x}_0, u_i \rangle u_i$$

- 3 $\mathbf{x}_0 - w$ is the shortest solution of $\mathbf{Ax} = \mathbf{b}$

This solution requires the calculation of the projection of \mathbf{x}_0 into $N(\mathbf{A})$. Let us discuss another, more direct, approach. If $\mathbf{v} \in N(\mathbf{A})$, $\mathbf{A}(\mathbf{v}) = \mathbf{0}$,

$$0 = \langle \mathbf{x}, \mathbf{A}(\mathbf{v}) \rangle = \langle \mathbf{A}^*(\mathbf{x}), \mathbf{v} \rangle$$

which means $\mathbf{v} \perp \mathbf{A}^*(\mathbf{x}) = 0$ for all \mathbf{x} . This means that the range of \mathbf{A}^* , $R(\mathbf{A}^*)$, is contained in the orthogonal complement $N(\mathbf{A})^\perp$ of $N(\mathbf{A})$. By the dimension formula we have $N(\mathbf{A})^\perp = R(\mathbf{A}^*)$.

Summary: The minimal vector \mathbf{s} satisfies

$$\mathbf{A}\mathbf{s} = \mathbf{b}, \quad \mathbf{s} \in R(\mathbf{A}^*)$$

That is, pick any solution of

$$\mathbf{A}\mathbf{A}^*\mathbf{y} = \mathbf{b},$$

and set

$$\mathbf{s} = \mathbf{A}^*\mathbf{y}.$$

Old Hourly #2 Questions

- (20 pts) Give proofs of the following facts:
 - If the 2×2 matrix A has nonzero nullspace and $A^2 = 2A$, then it is diagonalizable.
 - If the nullspace of a $n \times n$ matrix B is nonzero then $\det B = 0$.
- (20 pts) Let W be the subspace of \mathbb{R}^4 spanned by $v_1 = (1, 0, 1, 0)$, $v_2 = (1, 1, 0, 0)$.
 - Find an orthonormal basis for W .
 - Find the projection of $v = (1, 2, 3, 5)$ onto W .
 - Explain why the projection is a linear transformation and has determinant zero.

3. (20 pts) Let T be the linear transformation of $V = M_{2 \times 2}(\mathbb{C})$

$$T\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = \begin{bmatrix} c & a \\ d & b \end{bmatrix}$$

(a) Decide whether T is normal, hermitian, or neither.

(b) If T is diagonalizable, find a basis of eigenvectors.

4. (15 pts) Argue the following:

(a) If the characteristic polynomial of a linear transformation T splits into distinct linear factors then T is diagonalizable.

(b) There are nonzero matrices with some repeated eigenvalues that are diagonalizable [Give example]

5. (10 pts) Explain the meaning of every underlined keyword in the following statement:

Theorem: If T is a normal operator of a complex inner vector space V , then there is an orthonormal basis of eigenvectors of T .

6. (15 pts) If V is an inner product space,

(a) What is the meaning of the **triangle inequality** and of the **Cauchy-Schwarz inequality**?

(b) Give a proof of one of them.

1. (15 pts) Let $\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$ be a L.T. of the vector space \mathbf{V} over the field \mathbf{F} . Respond succinctly:

- 1 What is an eigenvector of \mathbf{T} ?
- 2 What are the eigenspaces of \mathbf{T} and what are their roles in deciding whether \mathbf{T} is diagonalizable?
- 3 Prove or disprove: All 2×2 complex matrices are diagonalizable.

2. (15 pts) Let $\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$ be a L.T. of the vector space \mathbf{V} over the field \mathbf{F} .

- 1 What is a \mathbf{T} -invariant subspace \mathbf{W} ?
- 2 If $v \in \mathbf{V}$ and \mathbf{W} is the span of the set of vectors $\{\mathbf{T}^n(v), n \geq 0\}$, prove that \mathbf{W} is \mathbf{T} -invariant.
- 3 Indicate the kind of matrix representation one gets for the restriction map $\mathbf{T}_{\mathbf{W}}$.

3. (12 pts) Let $\mathbf{A}, \mathbf{B} \in M_n(\mathbb{R})$.

- 1 What is $e^{\mathbf{A}}$? Argue that if \mathbf{A} is upper triangular then $e^{\mathbf{A}}$ is also upper triangular.
- 2 Prove that if $\mathbf{AB} = \mathbf{BA}$, then $e^{\mathbf{A+B}} = e^{\mathbf{A}}e^{\mathbf{B}}$.

3. (12 pts) Find the eigenvalues and corresponding eigenspaces of the linear transformation

$$\mathbf{A} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 10 & 3 \\ 0 & 3 & 2 \end{bmatrix}.$$

4. (20 pts) Let \mathbf{V} be the set of all real 2×2 matrices. If \mathbf{T} is the mapping

$$\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}, \quad \mathbf{T}(\mathbf{A}) = \mathbf{A} - (1/2)\text{trace}(\mathbf{A})\mathbf{I}$$

- 1 Prove that \mathbf{T} is a linear transformation.
 - 2 Prove that $\mathbf{T}^2 = \mathbf{T}$.
 - 3 Explain why maps such that $\mathbf{T}^2 = \mathbf{T}$ are always diagonalizable.
5. (13 pts) Let u, v_1 and v_2 be the following vectors of \mathbb{R}^4 , $(1, 2, 3, 4)$, $(1, 1, 1, 1)$ and $(2, -3, -3, 2)$.
- 1 Find an orthonormal basis of the subspace \mathbf{W} spanned by v_1, v_2 .
 - 2 Find the vector in \mathbf{W} closest to u ?

6. (15 pts)

- 1 What is an inner product space?
- 2 Argue that the Pythagorean theorem holds in such spaces.
- 3 If \mathbf{V} is the space of real continuous functions on $[0, 1]$, prove that $\int_0^1 f(t) \cdot g(t) dt$ defines an inner product on \mathbf{V} .

7. (10 pts) Let v_1, v_2, \dots, v_n a set of pairwise orthogonal vectors of the inner product space \mathbf{V} .

- 1 Prove that they are linearly independent.
- 2 Prove that

$$\|v_1 + v_2 + \dots + v_n\| = \sqrt{\sum_{i=1}^n \|v_i\|^2}.$$

3. (12 pts) Find the FULL set of solutions of the system of equations

$$\begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & 1 \\ 7 & 8 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -3 \\ 1 \\ -7 \end{bmatrix}.$$

3. (12 pts) Let A be a 3×3 matrix with determinant equal to 2.

(a) Explain carefully why A is invertible.

(b) If A is diagonalizable, explain carefully why A^{-1} is diagonalizable.

(c) What is the determinant of the matrix of cofactors of A ?

8. (6 pts) Let A be a 3×3 matrix with 3 nonzero entries of 2, 3 and 6. The other 6 entries are 0. Find and explain all the possible values for the determinant such matrices.
9. (8 pts) Let A be a 3×3 matrix whose columns are the vectors v_1 , v_2 and v_3 .
- (a) If a matrix B has for columns the vectors $2v_2 + v_3$, $3v_3 + v_1$ and v_1 , respectively, how are the determinants of A and B related?
- (b) Suppose further that v_1 , v_2 , v_3 are perpendicular to each other and satisfy

$$v_1 \cdot v_1 = 2, \quad v_2 \cdot v_2 = 6, \quad v_3 \cdot v_3 = 3.$$

Argue that the determinant of A is ± 6 . (Hint: multiply A by its transpose and take determinants.)

10. (9 pts) If A is a 3×3 matrix and $\det A = 2$, find the determinant of B if

(a) $B = 2A^2$ (careful, this is not $(2A)^2$)

(b) B is derived from A as follows: The first row of A is moved to the second row, the second row to the third row and the third row to the first row.

(c) $B = A^T \cdot A^{-1}$.

HomeQuiz #7

Section 6.3: 3a, 6, 10, 13, 18, 22a, 23

Today

- 1 Normal Operators ($\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T}$): real symmetric/skew symmetric
- 2 Hermitian Operator
- 3 Unitary Operator ($\mathbf{T}\mathbf{T}^* = \mathbf{I} = \mathbf{T}^*\mathbf{T}$): Orthogonal
- 4 Spectral Theorem
- 5 Goodies: Applications

Interesting diagonalizable operators

We are going to show a class of linear transformations that are diagonalizable. It will include the class represented by real symmetric matrices.

Let $\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$ be a L.T. of a complex inner product space. We have defined the **adjoint** \mathbf{T}^* of \mathbf{T} as the L.T. with the property

$$\langle \mathbf{T}(u), v \rangle = \langle u, \mathbf{T}^*(v) \rangle, \quad \forall u, v \in \mathbf{V}.$$

Let us compare the eigenvalues and eigenvectors of \mathbf{T} and \mathbf{T}^* :

Proposition

If λ is an eigenvalue of \mathbf{T} then $\bar{\lambda}$ is an eigenvalue of \mathbf{T}^ .*

Proof: Suppose $\mathbf{T}(u) = \lambda u$, $u \neq 0$. Then for any $v \in \mathbf{V}$,

$$\begin{aligned} 0 = \langle 0, v \rangle &= \langle (\mathbf{T} - \lambda \mathbf{I})(u), v \rangle = \langle u, (\mathbf{T} - \lambda \mathbf{I})^*(v) \rangle \\ &= \langle u, (\mathbf{T}^* - \bar{\lambda} \mathbf{I})(v) \rangle \end{aligned}$$

This says that $0 \neq u \perp \text{range}(\mathbf{T}^* - \bar{\lambda} \mathbf{I})$, so the range of $\mathbf{T}^* - \bar{\lambda} \mathbf{I}$ is not the whole of \mathbf{V} , which implies nullspace of $\mathbf{T}^* - \bar{\lambda} \mathbf{I} \neq 0$. This means that $\bar{\lambda}$ is an eigenvalue of \mathbf{T}^* .

Let us use this result to decide when a L.T. \mathbf{T} of an inner product space \mathbf{V} admits a basis \mathcal{A} such that

$$[\mathbf{T}]_{\mathcal{A}} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix},$$

that is, \mathbf{T} admits a matrix representation that is upper triangular.

Note that the characteristic polynomial has all of its roots in the field

$$\det(\mathbf{T} - x\mathbf{I}) = (a_{11} - x)(a_{22} - x) \cdots (a_{nn} - x),$$

that is the characteristic polynomial splits. Recall that this is always the case when the field is \mathbb{C} .

Theorem (Schur)

Let \mathbf{T} be a L.T. of the inner product space \mathbf{V} . If the characteristic polynomial of \mathbf{T} splits, then \mathbf{V} admits an orthonormal basis \mathcal{A} such that $[\mathbf{T}]_{\mathcal{A}}$ is upper triangular.

Proof: We will argue by induction on $\dim \mathbf{V} = n$. If $n = 1$, the assertion is obvious. Let us assume that the assertion holds for dimension $n - 1$. By the Proposition above, we know that \mathbf{T}^* has one eigenvalue λ . Let u be a unit vector so that $\mathbf{T}^*(u) = \lambda u$, and set \mathbf{W} for the subspace spanned by u . We claim that \mathbf{W}^{\perp} is \mathbf{T} -invariant: If $v \in \mathbf{W}^{\perp}$

$$\begin{aligned} \langle \mathbf{T}(v), u \rangle &= \langle v, \mathbf{T}^*(u) \rangle = \langle v, \lambda u \rangle \\ &= \bar{\lambda} \langle v, u \rangle = 0 \end{aligned}$$

So $\mathbf{T}(v) \in \mathbf{W}^{\perp}$.

We also have $\dim W + \dim \mathbf{W}^\perp = \dim \mathbf{V} = n$, so $\dim \mathbf{W}^\perp = n - 1$. Now we apply the induction hypothesis to the restriction of \mathbf{T} to \mathbf{W}^\perp : Let v_1, \dots, v_{n-1} be an orthonormal basis of \mathbf{W}^\perp for which the restriction of \mathbf{T} is upper triangular. If we add to the v_j the vector u , we get the orthonormal basis $\mathcal{A} = v_1, \dots, v_{n-1}, u$. The matrix representation

$$[\mathbf{T}]_{\mathcal{A}} = \begin{bmatrix} & & & a_{1n} \\ & [\mathbf{T}]_{\mathbf{W}^\perp} & & \vdots \\ & & & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix},$$

which has the desired form.

Normal operator

Observe that if there is an orthonormal basis \mathcal{A} of eigenvectors of \mathbf{T} , $[\mathbf{T}]_{\mathcal{A}}$ is a diagonal matrix, and since $[\mathbf{T}^*]_{\mathcal{A}} = [\mathbf{T}]_{\mathcal{A}}^*$, this matrix is also diagonal. Since diagonal matrices commute, we have $\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T}$.

Definition

A linear transformation \mathbf{T} of an inner product space is **normal** if $\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T}$.

Example: If \mathbf{A} is a symmetric real matrix, $\mathbf{A}^* = \mathbf{A}^t = \mathbf{A}$, so \mathbf{A} commutes with itself! Skew-symmetric real matrices, $\mathbf{A}^* = -\mathbf{A}$, are also normal.

Theorem

If \mathbf{T} is a normal operator ($\mathbf{T}\mathbf{T}^ = \mathbf{T}^*\mathbf{T}$) of a complex inner vector space \mathbf{V} , then there is an orthonormal basis of eigenvectors of \mathbf{T} . (The converse was proved already so this is a characterization of normal operators.)*

This is an important result, it has many useful consequences. To prove it we shall need some properties of normal operators.

Proposition

Let \mathbf{T} be a normal operator ($\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T}$) of the inner vector space \mathbf{V} .
Then:

- 1 $\|\mathbf{T}(u)\| = \|\mathbf{T}^*(u)\|$ for every $u \in \mathbf{V}$.
- 2 $\mathbf{T} - c\mathbf{I}$ is normal for every $c \in \mathbf{F}$.
- 3 If $\mathbf{T}(u) = \lambda u$ then $\mathbf{T}^*(u) = \bar{\lambda}u$.
- 4 If λ_1 and λ_2 are distinct eigenvalues of \mathbf{T} with corresponding eigenvectors u_1 and u_2 , then $u_1 \perp u_2$.

Proof: 1. For any vector $u \in \mathbf{V}$,

$$\begin{aligned} \|\mathbf{T}(u)\|^2 &= \langle \mathbf{T}(u), \mathbf{T}(u) \rangle = \langle \mathbf{T}^*\mathbf{T}(u), u \rangle = \langle \mathbf{T}\mathbf{T}^*(u), u \rangle \\ &= \langle \mathbf{T}^*(u), \mathbf{T}^*(u) \rangle = \|\mathbf{T}^*(u)\|^2 \end{aligned}$$

2. $(\mathbf{T} - c\mathbf{I})(\mathbf{T}^* - \bar{c}\mathbf{I}) = (\mathbf{T}^* - \bar{c}\mathbf{I})(\mathbf{T} - c\mathbf{I})$: check

3. Suppose $\mathbf{T}(u) = \lambda u$. Let $\mathbf{U} = \mathbf{T} - \lambda \mathbf{I}$. Then $\mathbf{U}(u) = 0$ so by 2. \mathbf{U} is normal and by 1. $\mathbf{U}^*(u) = 0$. That is $\mathbf{T}^*(u) = \bar{\lambda}u$.
4. Let λ_1 and λ_2 be distinct eigenvalues of \mathbf{T} with corresponding eigenvectors u_1 and u_2 . Then by 3.

$$\begin{aligned}\lambda_1 \langle u_1, u_2 \rangle &= \langle \lambda_1 u_1, u_2 \rangle = \langle \mathbf{T}(u_1), u_2 \rangle = \langle u_1, \mathbf{T}^*(u_2) \rangle \\ &= \langle u_1, \bar{\lambda}_2 u_2 \rangle = \lambda_2 \langle u_1, u_2 \rangle.\end{aligned}$$

Since $\lambda_1 \neq \lambda_2$, $\langle u_1, u_2 \rangle = 0$.

We are now in position to prove that a normal operator \mathbf{T} admits an orthonormal basis v_1, v_2, \dots, v_n of eigenvectors. We already know, by Schur theorem, that there is an orthonormal basis for which the matrix representation is upper triangular

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix}$$

We want to show that the off-diagonal elements are 0, that is, all the v_i are eigenvectors. [For simplicity we take $n = 3$] Note that $\mathbf{T}(v_1) = a_{11}v_1$, so v_1 is an eigenvector. To show v_2 is an eigenvector notice that

$$\mathbf{T}(v_2) = a_{12}v_1 + a_{22}v_2$$

We must show $a_{12} = 0$.

$$\mathbf{T}(v_2) = a_{12}v_1 + a_{22}v_2$$

We must show $a_{12} = 0$:

$$a_{12} = \langle \mathbf{T}(v_2), v_1 \rangle = \langle v_2, \mathbf{T}^*(v_1) \rangle = \langle v_2, \overline{a_{11}}v_1 \rangle = a_{11}\langle v_2, v_1 \rangle = 0$$

as desired. Now with v_1, v_2 eigenvectors, we show that $a_{13} = a_{23} = 0$. We consider

$$\mathbf{T}(v_3) = a_{13}v_1 + a_{23}v_2 + a_{33}v_3$$

The proof is similar: For instance

$$a_{23} = \langle \mathbf{T}(v_3), v_2 \rangle = \langle v_3, \mathbf{T}^*(v_2) \rangle = \langle v_3, \overline{a_{22}}v_2 \rangle = a_{22}\langle v_3, v_2 \rangle = 0$$

We have already remarked that real symmetric matrices, $\mathbf{A} = \mathbf{A}^t$, are normal. It turns out that **complex** symmetric matrices are not always normal. Truly the complex cousins of real symmetric matrices are called:

Definition

Let \mathbf{T} be a linear operator of the inner product space \mathbf{V} . \mathbf{T} is called **self-adjoint (Hermitian)** if $\mathbf{T} = \mathbf{T}^*$.

$$\mathbf{A} = \begin{bmatrix} 2 & 3 + 5i \\ 3 - 5i & 6 \end{bmatrix}$$

Lemma

Let \mathbf{T} be a self-adjoint linear operator of the inner product space \mathbf{V} .
Then

- ① Every eigenvalue is real.
- ② If \mathbf{V} is a real vector space then the characteristic polynomial splits.

Proof: 1. Suppose $\mathbf{T}(u) = \lambda u$, $u \neq 0$. By a previous result,
 $\mathbf{T}^*(u) = \bar{\lambda}u$. Since $\mathbf{T} = \mathbf{T}^*$, λ is real.

2. Let $n = \dim \mathbf{V}$, \mathcal{B} an orthonormal basis of \mathbf{V} and $\mathbf{A} = [\mathbf{T}]_{\mathcal{B}}$. Then \mathbf{A} is self-adjoint. Let $\mathbf{T}_{\mathbf{A}}$ be the linear operator of \mathbb{C}^n defined by $\mathbf{T}_{\mathbf{A}}(u) = \mathbf{A}u$ for all $u \in \mathbb{C}^n$.

Note that \mathbf{T}_A is self-adjoint because $[\mathbf{T}_A]_{\mathcal{C}} = \mathbf{A}$, where \mathcal{C} is the standard (orthonormal) basis of \mathbb{C}^n . So the eigenvalues of \mathbf{T}_A are real. Since the characteristic polynomial of \mathbf{T}_A is equal to the characteristic polynomial of \mathbf{A} , which is equal to the characteristic of \mathbf{T} , the characteristic polynomial of \mathbf{T} splits.

What we are saying is the following: Let \mathbf{A} be a $n \times n$ symmetric real matrix and employ it to define a L.T. of the **complex** vector space \mathbb{C}^n

$$\mathbf{T} = \mathbf{T}_A : \mathbb{C}^n \rightarrow \mathbb{C}^n, \quad \mathbf{T}(u) = \mathbf{A}(u).$$

Note $\det(\mathbf{T} - x\mathbf{I}) = \det(\mathbf{A} - x\mathbf{I})$.

First Main Theorem of the Course

Theorem

Let \mathbf{T} be a linear operator on the finite-dimensional inner product space \mathbf{V} . Then \mathbf{T} is self-adjoint if and only if there exists an orthonormal basis of \mathbf{V} consisting of eigenvectors of \mathbf{T} .

Unitary Operators

Definition

A linear operator \mathbf{T} of the inner product space \mathbf{V} is called **unitary** if $\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T} = \mathbf{I}$. If \mathbf{V} is a real inner product space, \mathbf{T} is called **orthogonal**.

The rotation operator

$$\mathbf{T}(x, y) = (x \cos \alpha + y \sin \alpha, -x \sin \alpha + y \cos \alpha)$$

is a major example.

If \mathbf{A} is a complex n -by- n matrix and $\mathbf{A}\mathbf{A}^* = \mathbf{A}^*\mathbf{A} = \mathbf{I}$, the column vectors of \mathbf{A} form an orthonormal basis of \mathbb{C}^n .

We now develop quickly some basic properties of these operators.

Theorem

Let \mathbf{T} be a linear operator of the finite-dimensional inner product space \mathbf{V} . TFAE:

- 1 \mathbf{T} is an unitary operator: $\mathbf{T}\mathbf{T}^* = \mathbf{T}^*\mathbf{T} = \mathbf{I}$.
- 2 $\langle \mathbf{T}(u), \mathbf{T}(v) \rangle = \langle u, v \rangle$ for all $u, v \in \mathbf{V}$.
- 3 For every orthonormal basis $\mathcal{B} = v_1, \dots, v_n$ of \mathbf{V} , $\mathbf{T}(v_1), \dots, \mathbf{T}(v_n)$ is also an orthonormal basis of \mathbf{V} .
- 4 For some orthonormal basis $\mathcal{B} = v_1, \dots, v_n$ of \mathbf{V} , $\mathbf{T}(v_1), \dots, \mathbf{T}(v_n)$ is also an orthonormal basis of \mathbf{V} .
- 5 $\|\mathbf{T}(u)\| = \|u\|$ for every $u \in \mathbf{V}$.

Proof. 1 \Rightarrow 2, 3, 4, 5: (Other \Rightarrow LTR)

$$\langle u, v \rangle = \langle \mathbf{T}^*\mathbf{T}(u), v \rangle = \langle \mathbf{T}(u), (\mathbf{T}^*)^*(v) \rangle = \langle \mathbf{T}(u), \mathbf{T}(v) \rangle.$$

$$\delta_{ij} = \langle v_i, v_j \rangle = \langle \mathbf{T}(v_i), \mathbf{T}(v_j) \rangle.$$

Properties of unitary operators

Let \mathbf{T} be an unitary operator of the inner product space \mathbf{V} .

- ① The eigenvalues of \mathbf{T} have length 1: If $\mathbf{T}(u) = \lambda u$,

$$\langle u, u \rangle = \langle \mathbf{T}(u), \mathbf{T}(u) \rangle = \langle \lambda u, \lambda u \rangle = \bar{\lambda} \lambda \langle u, u \rangle$$

and thus $\bar{\lambda} \lambda = 1$.

- ② If \mathbf{A} is a matrix representation of \mathbf{T} ,
 $|\det(\mathbf{A})| = 1: \det(\mathbf{A}) \det(\mathbf{A}^*) = 1$
- ③ If \mathbf{T} is orthogonal, $\det(\mathbf{A}) = \pm 1$.
- ④ If \mathbf{T} and \mathbf{U} are unitary operators, then \mathbf{T}^* and $\mathbf{T} \circ \mathbf{U}$ are also unitary operators.

Orthogonal operators of \mathbb{R}^2

We have already mentioned rotations, R_α . Let us analyze the possibilities. Let

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = [v_1 | v_2] \quad \|v_1\| = \|v_2\| = 1, \quad v_1 \perp v_2$$

be an orthogonal matrix. This means

$$a^2 + c^2 = 1, \quad b^2 + d^2 = 1, \quad ab + cd = 0$$

We can set $a = \cos \alpha$, $c = \sin \alpha$ and $b = \cos \beta$, $d = \sin \beta$ so that

$$ab + cd = \cos \alpha \cos \beta + \sin \alpha \sin \beta = \cos(\alpha - \beta) = 0.$$

This means that $\alpha - \beta = \pm\pi/2$. The two possibilities lead to

$$R_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}, \quad \mathbf{T} = \begin{bmatrix} \cos \beta & \sin \beta \\ \sin \beta & -\cos \beta \end{bmatrix}$$

To analyze

$$\mathbf{T} = \begin{bmatrix} \cos \beta & \sin \beta \\ \sin \beta & -\cos \beta \end{bmatrix}$$

we look at its eigenvalues:

$$\det(\mathbf{T} - x\mathbf{I}) = \begin{bmatrix} \cos \beta - x & \sin \beta \\ \sin \beta & -\cos \beta - x \end{bmatrix} = x^2 - 1$$

So $\lambda = \pm 1$. This means we have an orthonormal basis v_1, v_2 , and $\mathbf{T}(v_1) = v_1$, $\mathbf{T}(v_2) = -v_2$.

Thus the line $\mathbb{R}v_1$ is fixed under \mathbf{T} , and the perpendicular line $\mathbb{R}v_2$ is flipped about $\mathbb{R}v_1$. These transformations are called **reflections**.

Summary: If \mathbf{A} is an orthogonal 2-by-2 matrix, then if $\det \mathbf{A} = 1$, it is a rotation, and if $\det \mathbf{A} = -1$, it is a reflection.

Matrix product and dot product

Let u and v be two vectors of \mathbb{R}^n . Their **dot product**

$$u \cdot v = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

can be expressed as a **matrix product**

$$u^t v = \begin{bmatrix} a_1 & \cdots & a_n \end{bmatrix} \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

Keep in mind

$$u^t v = u \cdot v$$

Spectral Decomposition

Let \mathbf{A} be a n -by- n symmetric real matrix, $\mathbf{P} = [v_1 | \cdots | v_n]$ a matrix whose columns form an orthonormal basis of eigenvectors of \mathbf{A} :

$$\mathbf{A} = \mathbf{PDP}^t = [v_1 | \cdots | v_n] \cdot \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{bmatrix} \cdot \begin{bmatrix} \frac{v_1^t}{\|v_1\|} \\ \vdots \\ \frac{v_n^t}{\|v_n\|} \end{bmatrix}$$

Instead of this representation of \mathbf{A} as a product of 3 matrices, we are going to express \mathbf{A} as a **sum** of simple matrices of rank 1.

Expanding we get

$$\begin{aligned}
 \mathbf{A} &= \mathbf{PDP}^t = [v_1 | \cdots | v_n] \cdot \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{bmatrix} \cdot \begin{bmatrix} \frac{v_1^t}{} \\ \vdots \\ \frac{v_n^t}{} \end{bmatrix} \\
 &= [\lambda_1 v_1 | \cdots | \lambda_n v_n] \cdot \begin{bmatrix} \frac{v_1^t}{} \\ \vdots \\ \frac{v_n^t}{} \end{bmatrix} \\
 &= \lambda_1 v_1 v_1^t + \cdots + \lambda_n v_n v_n^t \\
 &= \sum \lambda_i \mathbf{P}_i, \quad \mathbf{P}_i = v_i v_i^t.
 \end{aligned}$$

Let us examine the matrices \mathbf{P}_i .

- 1 \mathbf{P}_i has rank 1 and is symmetric

$$\mathbf{P}_i = v_i v_i^t, \quad \mathbf{P}_i^t = (v_i v_i^t)^t = (v_i^t)^t v_i^t = \mathbf{P}_i$$

- 2 \mathbf{P}_i is a projection

$$\mathbf{P}_i \mathbf{P}_i = (v_i v_i^t)(v_i v_i^t) = v_i (v_i^t v_i) v_i^t = v_i v_i^t = \mathbf{P}_i$$

since $v_i^t v_i = \langle v_i, v_i \rangle = 1$

- 3 $\mathbf{P}_i \mathbf{P}_j = \mathbf{O}$ for $i \neq j$

$$\mathbf{P}_i \mathbf{P}_j = (v_i v_i^t)(v_j v_j^t) = v_i (v_i^t v_j) v_j^t = \mathbf{O}$$

since $v_i^t v_j = \langle v_i, v_j \rangle = 0$

The equality

$$\mathbf{A} = \sum \lambda_i \mathbf{P}_i, \mathbf{P}_i = v_i v_i^t$$

is called the **spectral decomposition** of \mathbf{A} .

Example: Let $\mathbf{A} = \begin{bmatrix} 3 & -4 \\ -4 & -3 \end{bmatrix}$

The eigenvalues are 5 and -5 , with corresponding [normalized] eigenvectors

$$v_1 = \frac{1}{\sqrt{5}} \begin{bmatrix} -2 \\ 1 \end{bmatrix}, \quad v_2 = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\mathbf{P}_1 = v_1 v_1^t = \begin{bmatrix} 4/5 & -2/5 \\ -2/5 & 1/5 \end{bmatrix}, \quad \mathbf{P}_2 = v_2 v_2^t = \begin{bmatrix} 1/5 & 2/5 \\ 2/5 & 4/5 \end{bmatrix}$$

Exercise:

Let \mathbf{A} be a real symmetric matrix. Prove that there is a symmetric matrix \mathbf{B} such that $\mathbf{B}^3 = \mathbf{A}$.

We know that there is an orthonormal basis v_1, \dots, v_n of eigenvectors of \mathbf{A} . The matrix $\mathbf{P} = [v_1 | \dots | v_n]$ is orthogonal [i.e. $\mathbf{P}^{-1} = \mathbf{P}^t$] and

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}$$

is a real diagonal matrix. Let \mathbf{E} be a real 'cubic root' of \mathbf{D} (if a diagonal entry of \mathbf{D} is d_{ii} , the corresponding entry of \mathbf{E} is the real root $d_{ii}^{1/3}$).

Set $\mathbf{B} = \mathbf{P}^{-1}\mathbf{E}\mathbf{P}$. Note

$$\mathbf{B}^t = (\mathbf{P}^{-1}\mathbf{E}\mathbf{P})^t = \mathbf{P}^t\mathbf{E}^t(\mathbf{P}^{-1})^t = \mathbf{P}^{-1}\mathbf{E}\mathbf{P} = \mathbf{B}, \quad \mathbf{B}^3 = \mathbf{P}^{-1}\mathbf{E}^3\mathbf{P} = \mathbf{A}.$$

Exercise: Let \mathbf{A} be skew-symmetric matrix. Prove that $\det \mathbf{A} \geq 0$. *Hint:* Recall that \mathbf{A} is normal, then pair up the complex eigenvalues of \mathbf{A} . Moreover, show that if \mathbf{A} has integer entries, then $\det \mathbf{A}$ is the square of an integer.

Real quadratic forms

A real **quadratic form** in n variables is a polynomial

$$\mathbf{q}(\mathbf{x}) = \sum_{i,j} a_{ij}x_i x_j.$$

They occur in the elementary theory of conic sections—e.g. what is $10x^2 + 6xy + 2y^2 = 5$, an ellipse, a parabola, or a hyperbola?— but also in the theory of max and min of functions $\mathbf{f}(x_1, \dots, x_n)$ of several variables. In both endeavors, a solution arises after an appropriate change of variables, $\mathbf{x} = \mathbf{P}(\mathbf{y})$,

$$\mathbf{q}(\mathbf{x}) = \mathbf{q}(\mathbf{P}(\mathbf{y})) = \sum_i d_i y_i^2.$$

Let us see how this comes about:

Let us begin with $Ax^2 + Bxy + Cy^2$, which we write as $ax^2 + 2bxy + cy^2$. (For general fields this would require $2 \neq 0$.) Now look:

$$\begin{aligned} ax^2 + 2bxy + cy^2 &= x(ax + by) + y(bx + cy) \\ &= \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} a & b \\ b & c \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \\ &= \mathbf{x}^t \mathbf{Q} \mathbf{x} \end{aligned}$$

where $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$ and \mathbf{Q} is a symmetric matrix.

It is routine to verify that every quadratic form $\mathbf{q}(\mathbf{x})$ has such a representation,

$$\mathbf{q}(\mathbf{x}) = \mathbf{x}^t \mathbf{Q} \mathbf{x}, \quad \mathbf{Q} = \mathbf{Q}^t$$

Now we can apply to \mathbf{Q} the spectral theorem we have developed.

Since \mathbf{Q} is (orthogonally) diagonalizable, there is an orthogonal matrix \mathbf{P} (formed by an orthonormal basis of eigenvectors of \mathbf{Q}) such that

$$\mathbf{P}^{-1}\mathbf{Q}\mathbf{P} = \mathbf{D} = \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{bmatrix}$$

This means that in $\mathbf{q}(\mathbf{x}) = \mathbf{x}^t\mathbf{Q}\mathbf{x}$, if we change the variables by the rule $\mathbf{x} = \mathbf{P}\mathbf{y}$,

$$\mathbf{q}(\mathbf{x}) = \mathbf{x}^t\mathbf{Q}\mathbf{x} = \mathbf{y}^t\mathbf{P}^{-1}\mathbf{Q}\mathbf{P}\mathbf{y} = \mathbf{y}^t\mathbf{D}\mathbf{y} = \sum_i \lambda_i y_i^2.$$

Some applications

Among the potential applications, we mentioned the identification of conics. For example, $10x_1^2 + 6x_1x_2 + 2x_2^2 = 5$: The matrix

$$\mathbf{Q} = \begin{bmatrix} 10 & 3 \\ 3 & 2 \end{bmatrix}$$

has for eigenvalues 11, 1 with

$$\mathbf{P} = \frac{1}{\sqrt{10}} \begin{bmatrix} 1 & -3 \\ 3 & 1 \end{bmatrix}$$

The change of variables $\mathbf{x} = \mathbf{P}\mathbf{y}$ gives

$$11y_1^2 + y_2^2 = 5,$$

the equation of an ellipse.

Another application, to the theory of max and min appears as follows: If \mathbf{a} is a critical point of the function $\mathbf{f}(\mathbf{x})$ —that is all the partial derivatives vanish at $\mathbf{x} = \mathbf{a}$, $\frac{\partial \mathbf{f}}{\partial x_i}(\mathbf{a}) = 0$, Taylor's expansion of \mathbf{f} in a neighborhood of \mathbf{a} gives

$$\mathbf{f}(\mathbf{x}) = \mathbf{f}(\mathbf{a}) + \mathbf{q}(\mathbf{h}) + \text{error}$$

where \mathbf{q} is a quadratic polynomial on the vector $\mathbf{h} = \mathbf{x} - \mathbf{a}$. The corresponding symmetric matrix is

$$\mathbf{Q} = \left[\frac{\partial^2 \mathbf{f}(\mathbf{x})}{\partial x_i \partial x_j}(\mathbf{a}) \right]$$

If all the eigenvalues of \mathbf{Q} are positive [negative], $\mathbf{q}(\mathbf{h}) \geq 0$ Then $\mathbf{f}(\mathbf{x}) \geq \mathbf{f}(\mathbf{a})$ in a neighborhood of \mathbf{a} : local max [local min] . The other cases are saddle points [the higher dimensional analogues of inflection points]

Rigid Motion

A **rigid motion** on the inner product space \mathbf{V} is a mapping

$$\mathbf{T} : \mathbf{V} \rightarrow \mathbf{V}$$

with the property

$$\|\mathbf{T}(u) - \mathbf{T}(v)\| = \|u - v\|, \quad \forall u, v \in \mathbf{V}.$$

That is, \mathbf{T} preserves distance of the images. A simple example is a translation: If \mathbf{a} is a fixed vector, the function

$$\mathbf{T}(v) := \mathbf{a} + v$$

is obviously a rigid motion. What else? We have seen that orthogonal transformations \mathbf{S} , $\mathbf{S}\mathbf{S}^t = \mathbf{I}$, preserve distances. Another such motion is obtained by composition: following a translation with an orthogonal mapping. What else? That is it!

Theorem

Any rigid motion \mathbf{T} of \mathbf{V} decomposes into $\mathbf{T} = \mathbf{S} \circ \mathbf{U}$, where \mathbf{S} is an orthogonal transformation and \mathbf{U} is a translation.

Proof: Set $\mathbf{a} = \mathbf{T}(O)$. Then the function $\mathbf{F}(u) = \mathbf{T}(u) - \mathbf{a}$ is a rigid motion and $\mathbf{F}(O) = O$. It is enough to prove that \mathbf{F} is orthogonal. Note that

$$\|\mathbf{F}(u) - \mathbf{F}(O)\| = \|u - O\|,$$

so \mathbf{F} preserves lengths, which is the key property of orthogonal transformations. BUT we are NOT assuming that \mathbf{F} is linear, we must prove it.

We first prove that \mathbf{F} preserves dot products: $\langle \mathbf{F}(u), \mathbf{F}(v) \rangle = \langle u, v \rangle$: We start from the equality and expand both sides

$$\begin{aligned}
\|\mathbf{F}(u) - \mathbf{F}(v)\|^2 &= \|u - v\|^2 \\
(\mathbf{F}(u) - \mathbf{F}(v)) \cdot (\mathbf{F}(u) - \mathbf{F}(v)) &= (u - v) \cdot (u - v) \\
\underbrace{\|\mathbf{F}(u)\|^2}_* - 2\langle \mathbf{F}(u), \mathbf{F}(v) \rangle + \underbrace{\|\mathbf{F}(v)\|^2}_{**} &= \underbrace{\|u\|^2}_* - 2\langle u, v \rangle + \underbrace{\|v\|^2}_{**}
\end{aligned}$$

Thus proving

$$\langle \mathbf{F}(u), \mathbf{F}(v) \rangle = \langle u, v \rangle.$$

Now we are going to prove that \mathbf{F} is a linear function by first showing that it is additive:

$$\begin{aligned}
\|\mathbf{F}(u+v) - \mathbf{F}(u) - \mathbf{F}(v)\|^2 &\stackrel{?}{=} 0 \\
\|\mathbf{F}(u+v)\|^2 + \|\mathbf{F}(u)\|^2 + \|\mathbf{F}(v)\|^2 - &= \|u+v\|^2 + \|u\|^2 + \|v\|^2 - \\
2\langle \mathbf{F}(u+v), \mathbf{F}(u) \rangle - 2\langle \mathbf{F}(u+v), \mathbf{F}(v) \rangle &= 2\langle (u+v), u \rangle - 2\langle (u+v), v \rangle \\
+ 2\langle \mathbf{F}(u), \mathbf{F}(v) \rangle &= +2\langle u, v \rangle \\
&= \|(u+v) - u - v\|^2 = 0.
\end{aligned}$$

Scaling, that $\mathbf{F}(cu) = c\mathbf{F}(u)$ for any $c \in \mathbb{R}$, has a similar proof: Expand

$$\|\mathbf{F}(cu) - c\mathbf{F}(u)\|^2$$

HomeQuiz #8

Section 6.4: 2f, 4, 6, 12, 13, 15

Section 6.5: 6, 10, 11, 17, 27a

Homework #9

- 1 Section 6.5, Problem 27d
- 2 Let \mathbf{A} be a 3×3 orthogonal matrix. Prove that \mathbf{A} is similar to a matrix of the form

$$\begin{bmatrix} \mathbf{R} & \mathbf{O} \\ \mathbf{O} & \pm 1 \end{bmatrix}$$

where \mathbf{R} is a 2×2 orthogonal matrix.

- 3 Section 6.3, Problem 22c
- 4 Let \mathbf{A} be a skew-symmetric real matrix. If \mathbf{A} diagonalizable, prove that $\mathbf{A} = \mathbf{O}$.