

Answers to review problems for Exam #2, Math 151, Sections 13, 14, 15

1. i) $\frac{5 \cos(5x) + 2e^{2x}}{\sin(5x) + e^{2x}}$. **Hint:** Use the chain rule.
 ii) $\frac{6x - 5}{1 + (3x^2 - 5x + 2)^2}$. **Hint:** Use the chain rule.
 iii) $-e^{\cos x} \sin x \sin(e^x) + e^{\cos x} \cos(e^x)e^x$. **Hint:** Use the product rule and the chain rule.
2. $y = 8x - 2$. **Hint:** Use the implicit differentiation to find the slope.
3. $f(4) \geq 16$. **Hint:** Use the mean value theorem: $f(4) - f(1) = f'(c)(4 - 1)$ for some c satisfying $1 \leq c \leq 4$. Since $f'(c) \geq 2$, $f'(c)(4 - 1) \geq 6$. So $f(4) \geq 16$.
4. (a) $h'(x) = 10x\sqrt{44 - 35x^2}$ and $h'(1) = 30$. **Hint:** Use the chain rule.
 (b) $h(0.95) \simeq 1.5$. **Hint:** Use $h(0.95) \simeq h(1) + h'(1)(0.95 - 1)$, $h(1) = f(2) = 3$ and $h'(1) = 30$.
 (c) Likely to be greater than the true value of $h(0.95)$. Reason: Since $h''(1) = -\frac{260}{3}$ is negative, the graph of $h(x)$ is concave downward near $x = 1$. So the tangent line is above the graph. Thus the linear approximation is greater than the true value.
5. (a) $1 - \frac{x}{2}$. **Hint:** Use the linear approximation formula: $f(x) \simeq f(a) + f'(a)(x - a)$.
 (b) 0.995. **Hint:** Take x to be 0.01 in the linear approximation above.
6. i) 0. **Hint:** Use L'Hôpital's rule.
 ii) $\frac{2}{\pi^2}$. **Hint:** Use L'Hôpital's rule.
 iii) 0. **Hint:** Use L'Hôpital's rule.
 iv) $\frac{3}{2}$. **Hint:** Use $x - \sqrt{x^2 - 3x} = \frac{3x}{x + \sqrt{x^2 - 3x}}$ and then use either the method of compute limits of algebraic functions or L'Hôpital's rule.
 v) e^2 . **Hint:** Take the logarithm of $(1 + 2x)^{\frac{1}{x}}$ and then use L'Hôpital's rule.
7. (a) We have $f(0) = 1$ and $f(1) = -\sin 3$. Since $0 < 3 < \pi$, we know that $\sin 3 > 0$. So $f(1) = -\sin 3 < 0$. Thus 0 is between $f(0)$ and $f(1)$. By the intermediate value theorem, there is c between 0 and 1 such that $f(c) = 0$, that is, $f(x)$ has a root in $[0, 1]$.
 (b) $g(x) = x + \frac{1 - x - \sin(3x)}{1 + 3 \cos(3x)}$. **Hint:** Use the formula $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$.

(c) $x_1 = 0.5 + \frac{0.5 - \sin(1.5)}{1 + 3 \cos(1.5)}$ and

$$x_2 = 0.5 + \frac{0.5 - \sin(1.5)}{1 + 3 \cos(1.5)} + \frac{0.5 - \frac{0.5 - \sin(1.5)}{1 + 3 \cos(1.5)} - \sin(1.5 + 3 \frac{0.5 - \sin(1.5)}{1 + 3 \cos(1.5)})}{1 + 3 \cos(1.5 + 3 \frac{0.5 - \sin(1.5)}{1 + 3 \cos(1.5)})}.$$

8. (a) $\lim_{x \rightarrow \infty} f(x) = \infty$ and $\lim_{x \rightarrow -\infty} f(x) = 0$. **Hint:** Since $\lim_{x \rightarrow \infty} (x^2 - 1) = \infty$ and $\lim_{x \rightarrow \infty} e^x = \infty$, $\lim_{x \rightarrow \infty} f(x) = \infty$. For $\lim_{x \rightarrow -\infty} f(x)$, use L'Hôpital's rule.
- (b) $f'(x) = (x^2 + 2x - 1)e^x$. The exact solutions of $f'(x) = 0$ are $-1 \pm \sqrt{2}$. When $x \leq -1 - \sqrt{2}$ or $x > -1 + \sqrt{2}$, $f'(x) > 0$. When $-1 - \sqrt{2} < x < -1 + \sqrt{2}$, $f'(x) < 0$.
- (c) $f''(x) = (x^2 + 4x + 1)e^x$. The exact solutions of $f''(x) = 0$ are $-2 \pm \sqrt{3}$. When $x \leq -2 - \sqrt{3}$ or $x > -2 + \sqrt{3}$, $f''(x) > 0$. When $-2 - \sqrt{3} < x < -2 + \sqrt{3}$, $f''(x) < 0$.
- (d) See the picture.
9. (a) Local minimum: 0. Local maximum: 2. **Hint:** Use the information on $f'(x)$.
- (b) Inflection points: $-1, 1, 3$. **Hint:** Use the information on $f''(x)$.
- (c) See the picture.
- (d) No. For $x > 4$, $f''(x) > 0$. So $f'(x)$ is increasing when $x > 4$. But $f'(4) = 0$. So $f'(x) \geq 0$ for $x > 4$.
10. (a) $\theta = \tan^{-1} \frac{a}{b}$. **Hint:** Draw a picture.
- (b) $\theta = \tan^{-1} 2$ and $\frac{d\theta}{dt} = -\frac{1}{50}$. **Hint:** θ can be obtained by just substitute the values of a and b . $\frac{d\theta}{dt}$ is obtained by using chain rules (both a and b are functions of t) and then using $a = 10$, $b = 5$, $a' = 0.3$ and $b' = 0.4$.
11. 5062.5. **Hint:** Let y be the length of the sides with parallel fencing inside and x the length of the other two sides. Then the area $A = xy$ and we have the constraint $2x + 5y = 450$. So $A = 90x - \frac{2}{5}x^2$. The domain is $[0, 225]$. A reaches its maximum at $x = 112.5$. Then calculate the area at this x .
12. $f(\frac{\pi}{6}) = -\frac{325\pi^3}{108} + \frac{5\pi}{6} - \frac{\sqrt{3}}{2} + 1$. **Hint:** Find $f'(x) = -6x^2 + \sin x + 6\pi^2 - 1$ first. Then find $f(x) = -2x^3 - \cos x + (6\pi^2 - 1)x + 1 - 4\pi^3 + \pi$.