

Dr. Z's Math251 Handout #15.4a (2nd ed.) [Double Integrals in Polar Coordinates]

By Doron Zeilberger

**Problem Type 15.4aa:** Evaluate the integral

$$\int \int_D F(x, y) dA \quad ,$$

where  $D$  is a region best described in polar coordinates,

$$D = \{ (r, \theta) \mid \alpha \leq \theta \leq \beta, h_1(\theta) \leq r \leq h_2(\theta) \} \quad .$$

**Example Problem 15.4aa:** Evaluate the integral

$$\int \int_D e^{-x^2-y^2} dA \quad ,$$

where  $D$  is the region bounded by the semi-circle  $x = \sqrt{25 - y^2}$  and the  $y$ -axis.

---

**Steps**

**1.** Draw the region and express it, if possible and convenient, as

$$D =$$

$$\{ (r, \theta) \mid \alpha \leq \theta \leq \beta, h_1(\theta) \leq r \leq h_2(\theta) \} \quad .$$

Of course, in many problems, the  $h_1(\theta)$  and/or  $h_2(\theta)$  may be plain numbers (i.e. not involve  $\theta$ ).

**Example**

**1.** This is a **semi**-circle, i.e. **half** a circle, center origin, radius 5, and since it is bounded by the  $y$ -axis, and  $x \geq 0$ , it is the **right** half

[Had it been  $x = -\sqrt{25 - y^2}$  it would have been the left-half. Had it been  $y = \sqrt{25 - x^2}$  it would have been the upper-half. Had it been  $y = -\sqrt{25 - x^2}$  it would have been the lower-half.]

Since it is the right-half,  $\theta$  ranges from  $\theta = -\pi/2$  (the downwards direction) to  $\theta = \pi/2$  (the upwards direction). For each ray  $\theta = \theta_0$ ,  $r$ , the distance from the origin, ranges from  $r = 0$  to  $r = 5$  (and indeed does not depend on  $\theta$  in this problem). So our region phrased in **polar coordinates** is:

$$D = \{ (r, \theta) \mid -\pi/2 \leq \theta \leq \pi/2, 0 \leq r \leq 5 \} \quad .$$

2. Rewrite the area integral

$$\int \int_D F(x, y) dA \quad ,$$

in **polar** coordinates by replacing

$x$  by  $r \cos \theta$ ,  $y$  by  $r \sin \theta$ ,  $dA$  by  $r dr d\theta$ .

[**shortcut:** Whenever you see  $x^2 + y^2$  you can replace it by  $r^2$ .]

Write it as an iterated integral

$$\int \int_D F(x, y) dA = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} F(r \cos \theta, r \sin \theta) r dr d\theta \quad ,$$

with the  $\theta$ -integral being at the **outside** and the  $r$ -integral being in the **inside**.

3. Evaluate this iterated integral by first doing the inner-integral (possibly getting an expression in  $\theta$ , or just a number), and then the outer integral.

2.

$$\begin{aligned} & \int \int_D e^{-x^2-y^2} dA \\ &= \int_{-\pi/2}^{\pi/2} \int_0^5 e^{-r^2} r dr d\theta \quad . \end{aligned}$$

3. The inside integral is (do the change-of-variable  $u = -r^2$ ):

$$\int_0^5 e^{-r^2} r dr = (-1/2)e^{-r^2} \Big|_0^5 = (1-e^{-25})/2 \quad ,$$

and the whole double-integral is

$$\begin{aligned} & \int_{-\pi/2}^{\pi/2} \int_0^5 e^{-r^2} r dr d\theta \\ &= \int_{-\pi/2}^{\pi/2} \left[ \int_0^5 e^{-r^2} r dr \right] d\theta \\ &= \int_{-\pi/2}^{\pi/2} [(1-e^{-25})/2] d\theta = (1-e^{-25})/2 \int_{-\pi/2}^{\pi/2} d\theta = \\ & [(1-e^{-25})/2][\pi/2 - (-\pi/2)] = \pi(1-e^{-25})/2 \quad . \end{aligned}$$

**Ans.:**  $\pi(1 - e^{-25})/2$  .

**Problem Type 15.4ab:** Find the volume of the solid above the surface  $z = f(x, y)$  and below the surface  $z = g(x, y)$ .

**Example Problem 15.4ab:** Find the volume of the solid above the cone  $z = \sqrt{x^2 + y^2}$  and below the sphere  $x^2 + y^2 + z^2 = 2$ .

### Steps

1. Find the “floor”, let’s call it  $D$ , by setting  $f(x, y) = g(x, y)$  (or if convenient already convert to polar coordinates).

2. The volume is the area integral of TOP-BOTTOM

$$\int \int_D [f(x, y) - g(x, y)] dA$$

Set it up. Then convert it to polar-coordinates.

### Example

1. In polar coordinates, the two surfaces are  $z = r$  and  $z = \sqrt{2 - r^2}$ . Setting them equal gives  $r = \sqrt{2 - r^2}$ . Squaring both sides gives  $r^2 = 2 - r^2$ , which gives  $2r^2 = 2$ , which gives  $r^2 = 1$  and so  $r = \pm 1$ . But  $r$  is never negative, so  $r = -1$  is nonsense. Hence the “floor”,  $D$ , is the region bounded by the circle  $r = 1$ , or, if you wish, the disk  $r \leq 1$ .

So

$$D = \{(r, \theta) | 0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$$

2. The bottom is  $z = \sqrt{x^2 + y^2}$ , and in polar  $z = r$ , and the top is  $x^2 + y^2 + z^2 = 2$  which is  $z = \sqrt{2 - x^2 - y^2}$  and in polar  $z = \sqrt{2 - r^2}$ . So the volume in polar coordinates is

$$\int_0^{2\pi} \int_0^1 [\sqrt{2 - r^2} - r] r dr d\theta$$

**3.** Evaluate the iterated integral. First do the inner integral (w.r.t. to  $r$ ) getting an expression in  $\theta$  (or just a number), and then do the outer integral.

**3.** The inner integral is

$$\begin{aligned}\int_0^1 [\sqrt{2-r^2}-r] r \, dr &= \int_0^1 [r\sqrt{2-r^2}-r^2] \, dr \\&= \int_0^1 r(2-r^2)^{1/2} \, dr - \int_0^1 r^2 \, dr \\&= -(1/3)(2-r^2)^{3/2} \Big|_0^1 - r^3/3 \Big|_0^1 \\&= -(1/3)(2-r^2)^{3/2} \Big|_0^1 - r^3/3 \Big|_0^1 \\&= -(1/3)[(2-1^2)^{3/2} - (2-0^2)^{3/2}] - 1/3 \\&= [2^{3/2} - 2]/3 = (2\sqrt{2} - 1)/3 \quad .\end{aligned}$$

The whole integral is thus:

$$\begin{aligned}\int_0^{2\pi} \int_0^1 [\sqrt{2-r^2}-r] r \, dr \, d\theta \\&= \int_0^{2\pi} \left[ \int_0^1 [\sqrt{2-r^2}-r] r \, dr \right] d\theta \\&= \int_0^{2\pi} (2\sqrt{2}-1)/3 \, d\theta \\&= 2\pi(2\sqrt{2}-1)/3 \quad .\end{aligned}$$

**Ans.:** The volume is  $2\pi(2\sqrt{2}-1)/3$ .

**Problem Type 15.4ac:** Evaluate the iterated integral by converting to polar coordinates.

$$\int_a^b \int_{f_1(y)}^{f_2(y)} F(x, y) \, dx \, dy$$

**Example Problem 15.4ac:** Evaluate the iterated integral by converting to polar coordinates.

$$\int_0^3 \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} x^2 y \, dx \, dy$$

**Steps**

**Example**

1. By looking at the limits of integration of the outer and inner integral signs, figure out the region  $D$ .

$$D = \{(x, y) \mid a \leq y \leq b, f_1(y) \leq x \leq f_2(y)\}$$

Draw this region, and express it in polar coordinates

$$D = \{(r, \theta) \mid \alpha \leq \theta \leq \beta, g_1(\theta) \leq r \leq g_2(\theta)\}$$

1. Our region is:

$$D = \{(x, y) \mid 0 \leq y \leq 3, -\sqrt{9-y^2} \leq x \leq \sqrt{9-y^2}\} \quad .$$

. Drawing it (do it!), we see that this is the upper-half of the circle whose center is the origin and whose radius is 3. In polar coordinates it is:

$$D = \{(r, \theta) \mid 0 \leq \theta \leq \pi, 0 \leq r \leq 3\} \quad .$$

2. Write the iterated integral as an area integral, then convert it to an iterated integral in polar coordinates. Use the “dictionary”  $x = r \cos \theta$   $y = r \sin \theta$   $dx dy = r dr d\theta$ .

2.

$$\begin{aligned} & \int_0^3 \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} x^2 y \, dx \, dy \\ &= \int_0^\pi \int_0^3 (r \cos \theta)^2 (r \sin \theta) r \, dr \, d\theta \\ &= \int_0^\pi \int_0^3 r^4 \sin \theta \cos^2 \theta \, dr \, d\theta \quad . \end{aligned}$$

**3.** Evaluate that iterated integral by doing the inner integral first, and then the outer integral.

**3.** The inner integral is

$$\begin{aligned}\int_0^3 r^4 \sin \theta \cos^2 \theta \, dr &= \sin \theta \cos^2 \theta \int_0^3 r^4 \, dr \\ &= \sin \theta \cos^2 \theta \left[ \frac{r^5}{5} \right]_0^3 \\ &= \frac{243}{5} \sin \theta \cos^2 \theta \quad .\end{aligned}$$

The outer integral is:

$$\begin{aligned}\int_0^\pi \int_0^3 r^4 \sin \theta \cos^2 \theta \, dr \, d\theta &= \int_0^\pi \left[ \int_0^3 r^4 \sin \theta \cos^2 \theta \, dr \right] d\theta \\ &= \int_0^\pi \frac{243}{5} \cos^2 \theta \sin \theta \, d\theta = \frac{243}{5} \int_0^\pi \cos^2 \theta \sin \theta \, d\theta \\ &= \frac{243}{5} \cdot \left[ -\frac{\cos^3 \theta}{3} \right]_0^\pi = \frac{81}{5} \cdot (-\cos^3(\pi) - (-\cos^3(0))) = \frac{162}{5} \quad .\end{aligned}$$

**Ans.:**  $\frac{162}{5}$ .

### A Problem from a Previous Final

Use polar coordinates to compute the double integral

$$\int \int_D xy \, dA \quad ,$$

where

$$D = \{(x, y) \mid x^2 + y^2 \leq 4, x \geq 0, y \geq 0\} \quad .$$

(Hint: recall the trig identity  $\sin 2\theta = 2 \sin \theta \cos \theta$ )

**Ans.:** 2.

### Another Problem from a Previous Final

Evaluate the iterated integral by converting it to polar coordinates

$$\int_0^1 \int_0^{\sqrt{1-x^2}} e^{x^2+y^2} \, dy \, dx \quad .$$

**Ans.:**  $\pi(e-1)/4$ .