

**Dr. Z's Math251 Handout #17.2 (2nd ed.) [Stokes' Theorem]**

By Doron Zeilberger

**Problem Type 17.2a:** Use Stokes' Theorem to evaluate  $\int \int_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ , where

$$\mathbf{F}(x, y, z) = P(x, y, z)\mathbf{i} + Q(x, y, z)\mathbf{j} + R(x, y, z)\mathbf{k} \quad ,$$

and  $S$  is some surface with a given orientation (that should boil down to either outwards or inwards).

**Example Problem 17.2a:** Use Stokes' Theorem to evaluate  $\int \int_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ , where

$$\mathbf{F}(x, y, z) = x^2 e^{2yz}\mathbf{i} + y^2 e^{3xz}\mathbf{j} + z^2 e^{4xy}\mathbf{k} \quad ,$$

and  $S$  is the hemisphere  $x^2 + y^2 + z^2 = 9$ ,  $z \geq 0$ , oriented upwards.

---

**Steps**

**Example**

1. You are going to use Stokes' Theorem

$$\int \int \text{curl } \mathbf{F} \cdot d\mathbf{S} = \int_C \mathbf{F} \cdot d\mathbf{r} \quad .$$

The challenge is to find the bounding curve  $C$ .

For hemispheres  $x^2 + y^2 + z^2 = R^2$ ,  $z \geq 0$ , like in this problem, the bounding curve is simply the circle  $x^2 + y^2 = R^2$ , on the  $xy$ -plane  $z = 0$  and the parametric representation is

$$x = R \cos t, y = R \sin t, z = 0 \quad .$$

If it is the hemisphere  $z \leq 0$  then it is the same but in the negative direction. If it is the hemisphere  $x^2 + y^2 + z^2 = R^2$ ,  $y \leq 0$ , then the parametric representation is

$$x = R \cos t, z = R \sin t, y = 0 \quad ,$$

etc.

If it is the part of a surface  $z = g(x, y)$  that lies above a plane  $z = a$ , oriented outwards, then  $C$  is obtained by solving  $g(x, y) = a$  and representing  $g(x, y) = a$  in parametric notation and adding to it  $z = a$ . The orientation of  $C$  is such as to obey the right-hand rule.

At the end you need to represent  $C$  in parametric form

$$\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle, a \leq t \leq b \quad ,$$

for some expressions  $x(t), y(t), z(t)$  of  $t$  and some numbers  $a$  and  $b$ .

1. Here  $C$  is simply the circle  $x^2 + y^2 = 3^2$  that lives in the  $xy$ -plane, and its parametric representation is

$$x = 3 \cos t, y = 3 \sin t, z = 0 \quad .$$

So

$$\mathbf{r}(t) = 3 \cos t \mathbf{i} + 3 \sin t \mathbf{j} + 0 \mathbf{k} \quad ,$$

$$0 \leq t \leq 2\pi \quad .$$

**2.** Plug-in the expressions for  $x, y, z$  in terms of  $t$  in  $\mathbf{F}$ , in order to express it in terms of the parameter  $t$ . Also figure out  $d\mathbf{r} = \mathbf{r}'(t) dt$ .

**2.**

$$\begin{aligned}\mathbf{F}(x, y, z) &= (3 \cos t)^2 e^0 \mathbf{i} + (3 \sin t)^2 e^0 \mathbf{j} + 0 \mathbf{k} \\ &= 9 \cos^2 t \mathbf{i} + 9 \sin^2 t \mathbf{j} + 0 \mathbf{k} \quad , \\ d\mathbf{r}(t) &= (-3 \sin t \mathbf{i} + 3 \cos t \mathbf{j} + 0 \mathbf{k}) dt \quad .\end{aligned}$$

**3.** Find the dot product  $\mathbf{F} \cdot d\mathbf{r}$  and integrate from  $t = a$  to  $t = b$ .

**3.**

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= ((9 \cos^2 t) \cdot (-3 \sin t) + (9 \sin^2 t) \cdot (3 \cos t) + 0 \cdot 0) dt = \\ &= (-27 \cos^2 t \sin t + 27 \sin^2 t \cos t) dt \quad .\end{aligned}$$

Finally, integrating from  $t = 0$  to  $t = 2\pi$ , we get

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^{2\pi} (-27 \cos^2 t \sin t + 27 \sin^2 t \cos t) dt \\ &= 9 \cos^3 t + 9 \sin^3 t \Big|_0^{2\pi} \\ &= (9 \cos^3(2\pi) + 9 \sin^3(2\pi)) - (9 \cos^3(0) + 9 \sin^3(0)) \\ &= 9 - 9 = 0 \quad .\end{aligned}$$

**Ans.:** 0.

**Problem Type 17.2b:** Use Stokes' Theorem to evaluate  $\int \int_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ , if

$$\mathbf{F} = P(x, y, z) \mathbf{i} + Q(x, y, z) \mathbf{j} + R(x, y, z) \mathbf{k} \quad ,$$

and  $S$  consists of the top and four sides (but not the bottom) of the cube with vertices  $(\pm A, \pm A, \pm A)$ .

**Example Problem 17.2b:** Use Stokes' Theorem to evaluate  $\int \int_S \text{curl } \mathbf{F} \cdot d\mathbf{S}$ , if

$$\mathbf{F} = x^2 y^2 \mathbf{i} + y^2 z \mathbf{j} + y z^2 \mathbf{k} \quad ,$$

and  $S$  consists of the top and four sides (but not the bottom) of the cube with vertices  $(\pm 2, \pm 2, \pm 2)$ .

**Steps**

**Example**

1. In this problem, it is possible to find the bounding curve  $C$ , and use Stokes's Theorem directly, **but**, in this case  $C$  is a square with four sides and we would have to do four integrals, and it is a pain. We will use Stokes's theorem **indirectly** by finding **another** surface with the same bounding curve. Naturally for a box in which the given surface consists of the top and the four walls, the bottom is such a surface.

2. Find  $\text{curl } \mathbf{F}$ .

3. Plug-in  $z = -A$  and note that  $dS = dx dy \mathbf{k}$ , and the region of integration is

$$\{(x, y) \mid -A \leq x \leq A, -A \leq y \leq A\} \quad .$$

Do the integration

1. The bottom face is

$$-2 \leq x, y \leq 2 \quad , \quad z = -2 \quad .$$

2.

$$\text{curl } \mathbf{F} = (z^2 - y^2) \mathbf{i} + 0 \mathbf{j} - 2x^2 y \mathbf{k}$$

(You do it!)

3. When  $z = -2$ ,

$$\text{curl } \mathbf{F} = (4 - y^2) \mathbf{i} + 0 \mathbf{j} - 2x^2 y \mathbf{k} \quad .$$

So

$$\text{curl } \mathbf{F} \cdot d\mathbf{S} = ((4 - y^2) \mathbf{i} + 0 \mathbf{j} - 2x^2 y \mathbf{k}) \cdot \mathbf{k} = -2x^2 y \quad .$$

Finally,

$$\begin{aligned} \iint_S \text{curl } \mathbf{F} \cdot d\mathbf{S} &= \int_{-2}^2 \int_{-2}^2 -2x^2 y \, dx \, dy \\ &= \int_{-2}^2 \left[ \int_{-2}^2 -2x^2 y \, dx \right] dy = \int_{-2}^2 (-2y) \left[ \frac{x^3}{3} \right]_{-2}^2 dy \\ &= \int_{-2}^2 (-2y) \frac{16}{3} dy \\ &= \frac{-32}{3} \int_{-2}^2 y \, dy = \frac{-32}{3} \cdot \left[ \frac{y^2}{2} \right]_{-2}^2 = \frac{-32}{3} \cdot 0 = 0 \quad . \end{aligned}$$

**Ans.:** 0 .

**Problem Type 17.2c:** Use Stokes' Theorem to evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where

$$\mathbf{F}(x, y, z) = P(x, y, z)\mathbf{i} + Q(x, y, z)\mathbf{j} + R(x, y, z)\mathbf{k} \quad ,$$

and  $C$  is a curve that bounds some surface (that you have to figure out!),  $z = g(x, y)$ , above the region  $\{(x, y) | (x, y) \in D\}$  (that you have to find!).  $C$  is oriented counterclockwise as viewed from above.

**Example Problem 17.2c:** Use Stokes' Theorem to evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where

$$\mathbf{F}(x, y, z) = (2x + y^2)\mathbf{i} + (2y + z^2)\mathbf{j} + (3z + x^2)\mathbf{k} \quad ,$$

and  $C$  is the triangle with vertices  $(2, 0, 0), (0, 2, 0), (0, 0, 2)$ , and is oriented counterclockwise as viewed from above.

---

### Steps

**1.** Find a convenient surface that our curve bounds, and express it in terms of  $z = g(x, y)$ . Also figure out its projection on the  $xy$ -plane.

**2.** Find  $\text{curl } \mathbf{F}$ .

### Example

**1.** The three vertices of our triangle lie on the plane  $x + y + z = 2$  (**you do it!**), so  $z = 2 - x - y$ , and  $g(x, y) = 2 - x - y$ . Also the projection of the triangle on the  $xy$  plane is bounded by the line  $x + y = 2$  and the axes, so it is the type I region

$$D = \{(x, y) | 0 \leq x \leq 2, 0 \leq y \leq 2 - x\} \quad .$$

**2.**

$$\text{curl } \mathbf{F} = -2z\mathbf{i} - 2x\mathbf{j} - 2y\mathbf{k} \quad .$$

(You do it!)

3. You have to use Stokes' Theorem

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_S \text{curl } \mathbf{F} \cdot d\mathbf{S} \quad .$$

Convert the surface integral into an area integral using the formula from 16.7

$$\begin{aligned} \int \int_S \mathbf{F} \cdot d\mathbf{S} = \\ \int \int_D \left( -P \frac{\partial g}{\partial x} - Q \frac{\partial g}{\partial y} + R \right) dA \end{aligned}$$

Be also sure to replace  $z$  by  $g(x, y)$ . **Note:** The  $\mathbf{F}$  from this formula is **not** the same as our  $\mathbf{F}$ , it is rather our  $\text{curl } \mathbf{F}$ , so use  $\mathbf{F}$  as a *local variable*.

3.

$$\int_S \text{curl } \mathbf{F} \cdot d\mathbf{S} = \int_S (-2z \mathbf{i} - 2x \mathbf{j} - 2y \mathbf{k}) \cdot d\mathbf{S}$$

where the surface is the one from step 1, i.e.  $z = 2 - x - y$  over  $D$  given above. Here we have  $P = -2z$ ,  $Q = -2x$ ,  $R = -2y$ ,  $g = 2 - x - y$ , so

$$\int \int_D (-(-2z)(-1) - (-2x)(-1) + (-2y)) dA \quad .$$

Now we have to replace  $z$  by  $2 - x - y$ , so this equals

$$\begin{aligned} \int \int_D (-2(2-x-y) - 2x - 2y) dA &= \int \int_D -4 dA \\ &= -4 \int \int_D dA \quad , \end{aligned}$$

where  $D$  is the triangle

$$D = \{(x, y) \mid 0 \leq x \leq 2, 0 \leq y \leq 2 - x\} \quad .$$

In this case the area integral is simply the area of the triangle ( $2 \cdot 2 / 2 = 2$ ) times  $-4$ , so the answer is  $-8$ , but of course you are welcome to do it without the shortcut:

$$\int \int_D (-4) dA = \int_0^2 \int_0^{2-x} (-4) dx dy = -8 \quad .$$

**Ans.:**  $-8$ .

### A Problem from a Previous Final

By using Stokes' Theorem, or otherwise, evaluate  $\int_C \mathbf{F} \cdot d\mathbf{r}$ , where

$$F(x, y, z) = yz^2 \mathbf{i} + xz^2 \mathbf{j} + 2xyz \mathbf{k} \quad ,$$

where  $C$  is the curve of intersection of the plane  $x + y + z = 1$  and the cylinder  $x^2 + y^2 = 9$ , oriented counterclockwise as viewed from above. Be sure to explain everything.

**Ans.:**  $0$ .

### Another Problem from a Previous Final

By using Stokes' theorem, or otherwise, evaluate the line integral

$$\int_C \mathbf{F} \cdot d\mathbf{r} \quad ,$$

where  $\mathbf{F}$  is the vector field

$$\mathbf{F}(x, y, z) = \langle 2xy^2z^2, 2x^2yz^2, 2x^2y^2z \rangle \quad ,$$

and  $C$  is the **closed** curve going from  $(1, 0, 1)$  to  $(3, 4, 9)$ , and then from  $(3, 4, 9)$  to  $(-1, 4, 11)$ , and then from  $(-1, 4, 11)$  to  $(5, 2, 11)$  and finally from  $(5, 2, 11)$  back to the starting point  $(1, 0, 1)$ . Explain everything!

**Ans.:** 0.