A Maple One-Line Proof of George Andrews's Formula that Says that the Number of Triangles with Integer Sides Whose Perimeter is n Equals $\{\frac{n^2}{12}\} - \lfloor\frac{n}{4}\rfloor\lfloor\frac{n+2}{4}\rfloor$

Shalosh B. EKHAD [1]

```
evalb(seq(coeff(taylor(q^3/(1-q^2)/(1-q^3)/(1-q^4),q=0,37),q,i),i=0..36)
=seq(round(n^2/12)-trunc(n/4)*trunc((n+2)/4),n=0..36));
```

Comments by Doron Zeilberger

1. The succinct formula of the title was discovered and first proved by George Andrews[A1], in a less-than-one-page cute note, improving a four-page note by Jordan et. al. [JWW]. Andrews's note, while short and sweet, is not self-contained, using results from his classic [A2].

2. Here is a clarification of the above Maple one-line proof. Arrange the lengths of the sides of a typical integer-side triangle in non-increasing order, and write them as $[a+b+c+1, b+c+1, c+1]$ for $a, b, c \geq 0$. By [E] I.20, $(b+c+1)+(c+1) > a+b+c+1$, so $c \geq a$, so $c = a+t$ for $t \geq 0$. So a generic integer-side triangle can be written as $[a+b+(a+t)+1, b+(a+t)+1, a+t+1] = [2a+b+t+1, a+b+t+1, a+t+1]$, and hence the number of triangles with integer sides whose perimeter equals n is the coefficient of q^n in

$$\sum_{a,b,t\geq 0} q^{(2a+b+t+1)+(a+b+t+1)+(a+t+1)} = \sum_{a,b,t\geq 0} q^{4a+3t+2b+3} = q^3\left(\sum_{a\geq 0} q^{4a}\right)\left(\sum_{t\geq 0} q^{3t}\right)\left(\sum_{b\geq 0} q^{2b}\right) =$$

$$\frac{q^3}{(1-q^4)(1-q^3)(1-q^2)} \quad .$$

The coefficient of q^n is obviously a *quasi-polynomial* of degree 2 and "period" $lcm(2,3,4) = 12$, but so is $\{\frac{n^2}{12}\} - \lfloor\frac{n}{4}\rfloor\lfloor\frac{n+2}{4}\rfloor$, hence it suffices to check that the first $(2+1)\cdot 12 + 1 = 37$ values match. $\square$

3. This is yet another example where "physical" (as opposed to "mathematical") induction, i.e. checking **finitely** (and not that many!) special cases, constitutes a *perfectly rigorous proof*! No need for the "simple" argument by "mathematical" induction alluded to by Andrews in his note. In this case we were in the *quasi-polynomial* **ansatz**. See [Z1][Z2], as well as the future classic [KP].

References

[A1] G. E. Andrews, *A note on partitions and triangles with integer sides*, Amer. Math. Monthly **86** (1979),477-478.

[A2] G. E. Andrews, *"The Theory of Partitions"*, Addison-Wesley, 1976. Reprinted by Cambridge University Press, 1984. First paperback edition, 1998.

[1] `ShaloshBEkhad@gmail.com` .

Exclusively published in: `http://www.math.rutgers.edu/~zeilberg/pj.html` and `http://arxiv.org/` .

Feb. 4 2012.

[E] Euclid, *"The Elements"*, Alexandria University Press, ca. 300 BC.

[JWW] J.H. Jordan, R. Walch, and R.J. Wisner, *Triangles with integer sides*, Amer. Math. Monthly **86** (1979), 686-689.

[KP] M. Kauers and P. Paule, *"The Concrete Tetrahedron"*, Springer, 2011
`http://www.springer.com/mathematics/analysis/book/978-3-7091-0444-6` .

[Z1] D. Zeilberger, *Enumerative and Algebraic Combinatorics*, in: "Princeton Companion to Mathematics" , (Timothy Gowers, ed.), Princeton University Press, pp. 550-561, 2008.
`http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimPDF/enu.pdf`

[Z2] D. Zeilberger, *An Enquiry Concerning Human (and Computer!) [Mathematical] Understanding*, in: C.S. Calude, ed., "Randomness & Complexity, from Leibniz to Chaitin" World Scientific, Singapore, 2007.
`http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimPDF/enquiry.pdf`