

## **INFORMATION TO USERS**

**This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.**

**The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.**

**In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.**

**Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each original is also photographed in one exposure and is included in reduced form at the back of the book.**

**Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.**

# **U·M·I**

University Microfilms International  
A Bell & Howell Information Company  
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA  
313/761-4700 800/521-0600



**Order Number 9215703**

**The Jacobian conjecture and the degree of field extension**

**Zhang, Yitang, Ph.D.**

**Purdue University, 1991**

**U·M·I**  
300 N. Zeeb Rd.  
Ann Arbor, MI 48106



**PURDUE UNIVERSITY  
GRADUATE SCHOOL  
Thesis Acceptance**

This is to certify that the thesis prepared

By Yitang Zhang

Entitled

The Jacobian Conjecture And  
The Degree of Field Extension

Complies with University regulations and meets the standards of the Graduate School for  
originality and quality

For the degree of Doctor of Philosophy

Signed by the final examining committee:

W. W. Mosh, chair  
Louis de Branges  
William J. Harvey  
J. Larson

Approved by:

J. Larson Department Head      9/18/91 Date

This thesis  is  is not to be regarded as confidential

W. W. Mosh  
Major Professor



39189

9/18/91

**THE JACOBIAN CONJECTURE AND  
THE DEGREE OF FIELD EXTENSION**

**A Thesis**

**Submitted to the Faculty**

**of**

**Purdue University**

**by**

**Yitang Zhang**

**In Partial Fulfillment of the  
Requirements for the Degree**

**of**

**Doctor of Philosophy**

**December 1991**

To my grandmother and grandfather and my parents

### ACKNOWLEDGEMENTS

I wish to express my sincere thanks to my major advisor, Professor Tzuong-Tsieng Moh for his kind encouragement and inspiring motivation.

I would like to thank Professors Louis de Branges, William Heinzer and Joseph Lipman for their support and serving on my committee.

## TABLE OF CONTENTS

	Page
ABSTRACT . . . . .	v
INTRODUCTION . . . . .	1
CHAPTER 1. $\pi$ -APPROXIMATE ROOTS OF A POLYNOMIAL . . . . .	4
CHAPTER 2. DEGREE OF FIELD EXTENSION . . . . .	7
CHAPTER 3. PROOF OF THEOREM . . . . .	11
BIBLIOGRAPHY . . . . .	16
VITA . . . . .	17

## ABSTRACT

Zhang, Yitang. Ph.D., Purdue University, December 1991. The Jacobian Conjecture and the degree of Field Extension. Major Professor: Tzuong-Tsieng Moh

Let  $k$  be an algebraically closed field of characteristic zero. If two polynomials  $f(x, y)$  and  $g(x, y)$  satisfy the Jacobian condition  $f_x g_y - f_y g_x \in k^*$ , then the degree of the field extension of  $k(x, y)$  over  $k(f, g)$ ,  $[k(x, y) : k(f, g)]$ , is less than or equal to the minimum of  $\deg f$  and  $\deg g$ .

## INTRODUCTION

The well-known Jacobian conjecture states "let  $k$  be an algebraically closed field of characteristic zero and  $f(x, y), g(x, y)$  two polynomials over  $k$ . If

$$(1) \quad f_x g_y - f_y g_x \in k^*$$

then  $k[x, y] = k[f, g]$ ." In other words a polynomial map from  $A^2$  to  $A^2$  is one-one and onto if and only if the Jacobian of the map is a nonzero constant. It was proved by Professor T.T.Moh among numerous other results [3] that this conjecture holds if

$$\max\{\deg f, \deg g\} \leq 100.$$

If two polynomials  $f(x, y)$  and  $g(x, y)$  satisfy the Jacobian condition (1), then it is easy to see that  $f(x, y)$  and  $g(x, y)$  are algebraically independent over  $k$ . Hence the field  $k(x, y)$  is a finite extension of the field  $k(f, g)$ . In fact it is well-known [1] that if  $f(x, y)$  and  $g(x, y)$  satisfy the Jacobian condition (1), and  $k(x, y) = k(f, g)$ , then  $k[x, y] = k[f, g]$ .

In the present paper, by means of the decomposition of polynomials over a Puiseux field, explicitly investigated by Professor T.T.Moh [3], we will prove the following

**THEOREM.** *Let  $k$  be an algebraically closed field of characteristic zero. Let  $f(x, y)$  and  $g(x, y)$  be two polynomials over  $k$ . If*

$$f_x g_y - f_y g_x \in k^*,$$

then the degree of field extension of  $k(x, y)$  over  $k(f, g)$ ,  $[k(x, y) : k(f, g)]$ , satisfies

$$[k(x, y) : k(f, g)] \leq \min\{\deg f, \deg g\}.$$

Throughout this paper, the ground field  $k$  is algebraically closed and of characteristic zero. The second and third sections are preparations for the proof of the theorem, where we do not assume that the polynomials satisfy the Jacobian condition.

In Chapter 1, we will give the definition and basic properties of  $\pi$ -approximate roots of a polynomial over the Puiseux field, which was introduced by Professor T.T.Moh [3]. We will also give a sufficient condition for a polynomial  $g(x, y)$  satisfying the condition that the field  $k(g)$  is relatively algebraically closed in  $k(x, y)$ , by the mean of  $\pi$ -approximate roots.

In Chapter 2, we will give a general result which states that if  $f(x, y)$  and  $g(x, y)$  are algebraically independent over  $k$  and monic in  $y$ , satisfying

- (i)  $k(g)$  is relatively algebraically closed in  $k(x, y)$ ,
- (ii)  $k(x, f, g) = k(x, y)$ ,

then there exists a constant  $c \in k$  such that  $g(x, y) - c$  is irreducible and such that

$$[k(x, y) : k(f, g)] = [k(\bar{x}, \bar{y}) : k(f(\bar{x}, \bar{y}))]$$

where  $\bar{x}$  and  $\bar{y}$  are the images of  $x$  and  $y$  under the quotient map

$$k[x, y] \longrightarrow k[x, y]/(g(x, y) - c).$$

In the last chapter, we will assume that  $f(x, y)$  and  $g(x, y)$  satisfy the Jacobian condition (1). Suppose that  $f(x, y)$  and  $g(x, y)$  are monic in  $y$  and  $\deg_y g(x, y) \leq \deg_y f(x, y)$ . Then it is not difficult to prove that the above conditions (i) and (ii)

are satisfied by  $f(x, y)$  and  $g(x, y)$ . Hence, with the same notations as above and by a result in algebraic function theory, we have

$$[k(x, y) : k(f, g)] = - \sum_{v(f(\bar{x}, \bar{y})) < 0} v(f(\bar{x}, \bar{y}))$$

where the summation is taken over all valuations  $v$  on  $k(\bar{x}, \bar{y})/k$  such that  $v(f(\bar{x}, \bar{y})) < 0$ . Now the key point of the proof of our theorem is that under the Jacobian condition (1) we can prove

$$\begin{aligned} - \sum_{v(f(\bar{x}, \bar{y})) < 0} v(f(\bar{x}, \bar{y})) &\leq - \sum_{v(\bar{x}) < 0} v(\bar{x}) \\ &= \deg_y g. \end{aligned}$$

Clearly the theorem then follows.

CHAPTER 1  
 $\pi$ -APPROXIMATE ROOTS OF A POLYNOMIAL

Following Professor Moh, for a polynomial  $g(x, y)$  over  $k$ , we put  $x = t^{-1}$  and regard  $g$  as a polynomial in the variable  $y$  over the Puiseux field  $k \ll t \gg$ . If  $g$  is monic in  $y$ , then  $g(t^{-1}, y)$  has the following decomposition over  $k \ll t \gg$ .

$$(2) \quad g(t^{-1}, y) = \prod_{i=1}^n (y - \tau_i)$$

with  $\tau_i \in k \ll t \gg$ .

DEFINITION 1. ([3]) Let  $\pi$  be a symbol, and let  $\sigma = \sum_{j < \delta} a_j t^j + \pi t^\delta \in k[\pi] \ll t \gg$  with  $a_j \in k$ . Then  $\sigma$  is said to be a  $\pi$ -approximate root (abbr.  $\pi$ -root) of a polynomial  $g(y)$  over  $k \ll t \gg$  if in the expression in  $k[\pi] \ll t \gg$

$$g(\sigma) = f_\sigma(\pi)t^\lambda + \text{higher terms in } t$$

we have  $1 \leq \deg_\pi g_\sigma(\pi) < \infty$ . The multiplicity of  $\sigma$  as a  $\pi$ -root of  $g(y)$  is defined to be  $\deg_\pi f_\sigma(\pi)$  as a polynomial in  $\pi$ .

The following properties are obvious.

(I) For every root  $\tau$  of the equation  $g(y) = 0$ , there exists a  $\pi$ -root  $\sigma$  of  $g(y)$  such that

$$\text{ord}_t g(\sigma) = 0$$

and

$$\sigma - \tau = (\pi - c)t^t + \text{higher terms in } t$$

for some  $c \in k$  and some rational number  $\iota$ . In this case we call  $\sigma$  a  $\pi$ -root associated to the root  $\tau$ .

On the other hand, for every  $\pi$ -root  $\sigma$  of  $g(y)$ , there exists a root  $\tau$  of the equation  $g(y) = 0$  such that  $\sigma$  is associated to  $\tau$ .

(II) Let  $\tau$  be a root of the equation  $g(y) = 0$  and let  $\sigma$  be a  $\pi$ -root of  $g(y)$  associated to  $\tau$ . Then for every  $\tau^* \in k \ll t \gg$ , we have

$$\text{ord}_t(\tau - \tau^*) \geq \text{ord}_t(\sigma - \tau^*),$$

and therefore

$$\text{ord}_t f(\tau) \geq \text{ord}_t f(\sigma)$$

for every polynomial  $f(y)$  over  $k \ll t \gg$ .

LEMMA 1.. Let  $g(x, y)$  be a polynomial monic in  $y$ . If  $g(t^{-1}, y)$  has a  $\pi$ -root  $\sigma$  of multiplicity one such that

$$\text{ord}_t g(t^{-1}, \sigma) = 0,$$

then the field  $k(g)$  is relatively algebraically closed in  $k(x, y)$ .

Proof.: Let  $\overline{k(g)}$  be the algebraic closure of  $k(g)$  in  $k(x, y)$ . We want to prove  $\overline{k(g)} = k(g)$ . Let  $R$  denote the integral closure of  $k[g]$  in  $\overline{k(g)}$ . Since every element in  $R$  is integral over  $k[x, y]$  which is integrally closed, we have  $R \subset k[x, y]$ . Since  $\overline{k(g)}$  is the fraction field of  $R$ , it suffices to show that  $R = k[g]$ . Let  $h$  be a non-zero polynomial in  $R$ . Then  $h$  satisfies an equation

$$h^s + G_1(g)h^{s-1} + \cdots + G_s(g) = 0$$

with  $G_s(g) \neq 0$ . By substituting  $x = t^{-1}$  and  $y = \sigma$ , it easily follows from the above equation that

$$(3) \quad \text{ord}_t h(t^{-1}, \sigma) = 0.$$

Write

$$h(t^{-1}, \sigma) = h_\sigma(\pi) + \text{higher terms in } t$$

. Since  $\deg g_\sigma(\pi) = 1$ , there exists a polynomial  $H$  in one variable such that  $h_\sigma(\pi) = H(g_\sigma(\pi))$ , and therefore

$$(4) \quad \text{ord}_t(h(t^{-1}, \sigma) - H(g(t^{-1}, \sigma))) > 0.$$

Since (3) holds for every non-zero polynomial in  $R$  and  $h - H(g) \in R$ , by (4) we have

$$h - H(g) = 0.$$

It follows that

$$R = k[g].$$

Lemma 1 is proved. ■

CHAPTER 2  
DEGREE OF FIELD EXTENSION

The main result of this section is the following

LEMMA 2. . Let  $f(x, y)$  and  $g(x, y)$  be two polynomials over  $k$  such that

- (i)  $f(x, y)$  and  $g(x, y)$  are algebraically independent over  $k$  and monic in  $y$ ,
- (ii) the field  $k(g)$  is relatively algebraically closed in  $k(x, y)$ , and
- (iii)  $k(x, f, g) = k(x, y)$ .

Then there exists a constant  $c \in k$  such that

$$(5) \quad [k(x, y) : k(f, g)] = [k(\bar{x}, \bar{y}) : k(f(\bar{x}, \bar{y}))]$$

where  $\bar{x}$  and  $\bar{y}$  are the images of  $x$  and  $y$  under the quotient map

$$k[x, y] \longrightarrow k[x, y]/(g(x, y) - c).$$

*Proof:* . Let

$$F(x, f, g) = 0$$

be a defining equation of  $x$ ,  $f(x, y)$  and  $g(x, y)$  over  $k$ . The polynomial  $F(X, Y, Z)$  with variables  $X, Y$  and  $Z$  is uniquely determined up to a constant factor. Since  $g(x, y)$  is monic in  $y$ ,  $f(x, y)$  is integral over  $k[x, g(x, y)]$ , and, in view of condition (iii),

$$[k(x, f, g) : k(x, g)] = \deg_y g.$$

Let  $n = \deg_y g$ . It follows from the above discussion that the polynomial  $F(X, Y, Z)$  can be taken as the form

$$F(X, Y, Z) = Y^n + F_1(X, Z)Y^{n-1} + \cdots + F_n(X, Z)$$

with

$$\max_{0 < i \leq n} \deg_X F_i(X, Z) = [k(x, y) : k(f, g)].$$

By condition (iii), there exists a non-zero polynomial  $G(X, Y, Z)$  such that

$$yG(x, f, g) \in k[x, f, g].$$

It is obvious that for all but finitely many constants  $c \in k$

$$(a) \quad G(x, f(x, y), g(x, y)) \not\equiv 0 \pmod{g(x, y) - c}$$

and

$$(b) \quad \max_{0 < i \leq n} \deg_X F_i(X, c) = [k(x, y) : k(f, g)].$$

On the other hand, by condition (ii) and the second Bertini theorem (see [4]), for all but finitely many constants  $c \in k$  we have

$$(c) \quad g(x, y) - c \quad \text{is irreducible.}$$

It follows that there exists a constant  $c$  satisfying (a), (b) and (c). Fix such a constant  $c$ . Let  $\bar{x}$  and  $\bar{y}$  be the images of  $x$  and  $y$  under the quotient map

$$k[x, y] \longrightarrow k[x, y]/(g(x, y) - c).$$

Notice that  $k[x, y]/(g(x, y) - c)$  is an integral domain and  $k(\bar{x}, \bar{y})$  is its fraction field. We claim that in the present case (5) holds. It follows from (a) that  $G(\bar{x}, f(\bar{x}, \bar{y}), c) \neq 0$ , and therefore  $\bar{y} \in k(\bar{x}, f(\bar{x}, \bar{y}))$ . Hence

$$(6) \quad [k(\bar{x}, f(\bar{x}, \bar{y})) : k(\bar{x})] = n = \deg_Y F(X, Y, c).$$

Clearly we have

$$(7) \quad F(\bar{x}, f(\bar{x}, \bar{y}), c) = 0.$$

If the polynomial  $F(X, Y, c)$  is reducible, then  $f(\bar{x}, \bar{y})$  satisfies an equation over  $k(\bar{x})$  whose degree is less than  $n$ . This contradicts (6). Hence  $F(X, Y, c)$  is irreducible and (7) is a defining equation of  $\bar{x}$  and  $f(\bar{x}, \bar{y})$  over  $k$ . By (b), it follows that

$$[k(\bar{x}, \bar{y}) : k(f(\bar{x}, \bar{y}))] = [k(\bar{x}, f(\bar{x}, \bar{y})) : k(f(\bar{x}, \bar{y}))] = \deg_X F(X, Y, c) = [k(x, y) : k(f, g)],$$

as claimed. ■

To end this chapter we state some basic terminologies and results in algebraic function theory. Let  $K$  denote a function field over  $k$ , that is,  $K$  is a finitely generated field of transcendence degree one over  $k$ . A valuation on  $K/k$  is a map  $v : K \rightarrow \mathbf{Z}$  such that

$$v(\alpha\beta) = v(\alpha) + v(\beta) \quad \text{for all } \alpha, \beta \in K,$$

$$v(c) = 0 \quad \text{for all } c \in k^*.$$

Here, as usual, we define  $v(0) = +\infty$ . We assume that every valuation on  $K/k$  is normalized, i.e., its valuation group is exactly  $\mathbf{Z}$ . The following results are well-known.

(I) Let  $\alpha \in K$ . If  $v(\alpha) \geq 0$  for every valuation  $v$  on  $K/k$ , then  $\alpha \in k$ .

(II) Let  $\alpha \in K, \alpha \notin k$ . Then  $K$  is a finite extension of  $k(\alpha)$ , and

$$[K : k(\alpha)] = - \sum_{v(\alpha) < 0} v(\alpha)$$

where the sum is taken over all valuations  $v$  on  $K/k$  satisfying  $v(\alpha) < 0$ .

(III) In the case that  $K$  is generated by  $\bar{x}$  and  $\bar{y}$ , let

$$g(\bar{x}, \bar{y}) = 0$$

be a defining equation of  $\bar{x}$  and  $\bar{y}$  over  $k$ . Then for every valuation  $v$  on  $K/k$  satisfying  $v(\bar{x}) < 0$ , there exists a root  $\tau$  of the equation  $g(t^{-1}, y) = 0$  in  $k \ll t \gg$  and a positive integer  $a$  such that

$$v(h(\bar{x}, \bar{y})) = a\{\text{ord}_t(h(t^{-1}, \tau))\}$$

for every polynomial  $h(x, y)$ .

CHAPTER 3  
PROOF OF THEOREM

From now on, we will assume that  $f(x, y)$  and  $g(x, y)$  are two polynomials monic in  $y$  satisfying

- (i)  $f_x g_y - f_y g_x \in k^*$ ,  
(ii)  $\deg_y g(x, y) \leq \deg_y f(x, y)$ .

Let  $n = \deg_y g(x, y)$ . To prove the theorem, it suffices to show that under the above conditions

$$[k(x, y) : k(f, g)] \leq n.$$

For notational simplicity we will write  $f(y) = f(t^{-1}, y)$ ,  $g(y) = g(t^{-1}, y)$ . Let  $\sigma = \sum_{j < \delta} a_j t^j + \pi t^\delta$  be a  $\pi$ -root of  $g(y)$  such that  $\text{ord}_t g(\sigma) = 0$ . Put  $\lambda = \text{ord}_t f(\sigma)$  and write

$$\begin{aligned} g(\sigma) &= g_\sigma(\pi) + \text{higher terms in } t, \\ f(\sigma) &= f_\sigma(\pi)t^\lambda + \text{higher terms in } t \end{aligned}$$

with  $\deg_\pi g_\sigma(\pi) \geq 1$ ,  $f_\sigma(\pi) \neq 0$ .

LEMMA 3. . Let  $\sigma, \delta, \lambda, g_\sigma(\pi)$  and  $f_\sigma(\pi)$  be as above. Then we have

- (i)  $\delta \geq 0$ ,  $\lambda \geq -1$ ,  
(ii) if  $\lambda \neq 0$ , then  $f_\sigma(\pi)g'_\sigma(\pi) \in k^*$ .

*Proof:* . (i). It is obvious that for every  $\tau \in k \ll t \gg$

$$\text{ord}_t(\sigma - \tau) \leq \delta.$$

Hence, in view of (2), we have

$$(8) \quad 0 = \text{ord}_t g(\sigma) \leq n\delta.$$

It is easy to see that the Jacobian of  $f(\sigma)$  and  $g(\sigma)$  with respect to  $t$  and  $\pi$  is of the form

$$(9) \quad J_{t,\pi}(f(\sigma), g(\sigma)) = \lambda f_\sigma(\pi) g'_\sigma(\pi) t^{\lambda-1} + \text{higher terms in } t.$$

On the other hand, by the chain rule and the Jacobian condition we get

$$(10) \quad J_{t,\pi}(f(\sigma), g(\sigma)) = J_{x,y}(f, g) J_{t,\pi}(t^{-1}, \sigma) = c_1 t^{-2+\delta}$$

for some  $c_1 \in k^*$ . To prove  $\lambda \geq -1$ , we may assume  $\lambda \neq 0$ . Comparing the right-hand sides of (9) and (10) we get

$$(11) \quad \lambda = -1 + \delta.$$

By (8) and (11) we get (i).

(ii). If  $\lambda \neq 0$ , then by (9) and (10) we get

$$f_\sigma(\pi) g'_\sigma(\pi) = \lambda^{-1} c_1 \in k^*.$$

Lemma 3 is proved. ■

LEMMA 4. . The field  $k(g)$  is relatively algebraically closed in  $k(x, y)$ .

Proof: . By Lemma 1, it suffices to show that  $g(y)$  has a  $\pi$ -root  $\sigma$  with multiplicity one such that

$$(12) \quad \text{ord}_t g(\sigma) = 0.$$

First we claim that  $g(y)$  has a  $\pi$ -root  $\sigma$  satisfying (12) and

$$(13) \quad \text{ord}_t f(\sigma) < 0.$$

On the contrary let us assume that (13) is false. Then for every  $\pi$ -root  $\sigma$  of  $g(y)$  satisfying (12) we have

On the contrary let us assume that (13) is false. Then for every  $\pi$ -root  $\sigma$  of  $g(y)$  satisfying (12) we have

$$\text{ord}_t f(\sigma) \geq 0.$$

: 0 we

have

$$(14) \quad \text{ord}_t f(\tau) \geq 0.$$

Let  $g_1$  be an irreducible factor of  $g$  and let  $K$  be the fraction field of the quotient ring  $k[x, y]/(g_1(x, y))$ . As we did in Section 3, we let  $\bar{x}$  and  $\bar{y}$  be the images of  $x$  and  $y$  under the quotient map  $k[x, y] \rightarrow k[x, y]/(g_1(x, y))$ . For every valuation  $v$  on  $K/k$ , if  $v(\bar{x}) \geq 0$ , then we have  $v(f(\bar{x}, \bar{y})) \geq 0$ . If  $v(\bar{x}) < 0$ , then by (14) and Section 3,(III) we also get  $v(f(\bar{x}, \bar{y})) \geq 0$ . It therefore follows from Section 3,(I) that

$$f(\bar{x}, \bar{y}) \in k.$$

Thus, there exists a constant  $c \in k$  such that

$$(15) \quad f(x, y) \equiv c \pmod{g_1(x, y)}.$$

Since  $g_1(x, y)$  is a factor of  $g(x, y)$ , by (15) and a simple computation we get

$$f_x g_y - f_y g_x \equiv 0 \pmod{g_1}.$$

A contradiction. Hence  $g(y)$  must have a  $\pi$ -root  $\sigma$  satisfying both (12) and (13), as claimed. In addition, for such a  $\pi$ -root  $\sigma$  of  $g(y)$ , by Lemma 3 we get  $g'_\sigma(\pi) \in k^*$ .

Hence the multiplicity of this  $\pi$ -root  $\sigma$  is equal to one. ■

REMARK. The decomposition of  $f(y)$  and  $g(y)$  over  $k \ll t \gg$  was explicitly investigated by Professor Moh [3] where much deeper results were obtained. In fact, the existence of a  $\pi$ -root  $\sigma$  of  $g(y)$  with multiplicity one and satisfying (12) can also be deduced from [3], Section 5, and the result in Lemma 3 is contained in [3], Section 4.

LEMMA 5. . The polynomials  $f(x, y)$  and  $g(x, y)$  satisfy

$$k(x, f, g) = k(x, y).$$

Proof: . See [3], Section 2. ■

It is well-known that the Jacobian condition implies that  $f(x, y)$  and  $g(x, y)$  are algebraically independent over  $k$ . Hence, by Lemma 4 and 5,  $f(x, y)$  and  $g(x, y)$  satisfy all conditions in Lemma 2. It now follows from Lemma 2 that there exists a constant  $c \in k$  such that  $g(x, y) - c$  is irreducible and such that

$$(16) \quad [k(x, y) : k(f, g)] = [k(\bar{x}, \bar{y}) : k(f(\bar{x}, \bar{y}))]$$

where  $\bar{x}$  and  $\bar{y}$  are images of  $x$  and  $y$  respectively under the quotient map  $k[x, y] \rightarrow k[x, y]/(g(x, y) - c)$ . We will simply write  $\bar{f} = f(\bar{x}, \bar{y})$ , and, without loss of generality, assume  $c = 0$ . By Section 3,(II) we get

$$(17) \quad [k(\bar{x}, \bar{y}) : k(\bar{f})] = - \sum_{v(\bar{f}) < 0} v(\bar{f})$$

where the summation is taken over all valuations  $v$  on  $k(\bar{x}, \bar{y})/k$  satisfying  $v(\bar{f}) < 0$ .

Proof of Theorem: . Since  $g(x, y)$  is monic in  $y$  and  $g(\bar{x}, \bar{y}) = 0$ , for every valuation  $v$  on  $k(\bar{x}, \bar{y})/k$ , if  $v(\bar{x}) \geq 0$ , then  $v(\bar{y}) \geq 0$ , and therefore  $v(\bar{f}) \geq 0$ . Hence, in the case  $v(\bar{f}) < 0$ , we have  $v(\bar{x}) < 0$ . Now let us fix a valuation  $v$  on

$k(\bar{x}, \bar{y})/k$  satisfying  $v(\bar{f}) < 0$ . By Chapter 3, (III), there exists a root  $\tau$  of the equation  $g(y) = 0$  and a positive integer  $a$  such that

$$(18) \quad v(\bar{f}) = a\{\text{ord}_t f(\tau)\}, \quad v(\bar{x}) = a\{\text{ord}_t(t^{-1})\} = -a.$$

By Section 2, (I), there exists a  $\pi$ -root  $\sigma$  associated to  $\tau$  such that  $\text{ord}_t g(\sigma) = 0$ .

It follows from Lemma 3 that

$$\text{ord}_t f(\sigma) \geq -1.$$

Since  $\sigma$  is associated to  $\tau$ , by (18) and Section 2, (II) we get

$$v(\bar{f}) \geq -a = v(\bar{x}).$$

From the above discussion we conclude that

$$(19) \quad \sum_{v(\bar{f}) < 0} v(\bar{f}) \geq \sum_{v(\bar{f}) < 0} v(\bar{x}) \geq \sum_{v(\bar{x}) < 0} v(\bar{x}).$$

By Section 3, (II) we get

$$(20) \quad \sum_{v(\bar{x}) < 0} v(\bar{x}) = -[k(\bar{x}, \bar{y}) : k(\bar{x})] = -\text{deg}_y g.$$

By (16), (17), (19) and (20), we complete the proof of the theorem. ■

**BIBLIOGRAPHY**

## BIBLIOGRAPHY

- [1] S.S.ABHYANKAR, *Expansion Techniques in Algebraic Geometry*, TATA Institute, Bombay, 1977.
- [2] C.CHEVALLEY, *Introduction to the Theory of Algebraic Functions of One Variable*, Mathematical Surveys VI, Am. Math. Society, 1951.
- [3] T.T.MOH, *On the Jacobian conjecture and the configurations of roots*, J. reine angew. Math., 340 (1983), pp. 140-212.
- [4] O.ZARISKI, *The theorem of Bertini on the variable singular points of a linear system*, Trans. Amer. Math. Soc., 56 (1944), pp. 130-140.

VITA

## VITA

Yitang Zhang was born on February 5, 1955 in Shanghai, China. He attended Peking University at Beijing, China where he received a B.A. degree in mathematics in 1982. He became a graduate student at the same university in the same year, and received a M.S. degree in mathematics in 1985. He began working towards his Ph.D. degree in mathematics in June 1985 at Purdue University.